

## Book Chapter

# Safety and Efficacy of Erythrocyte Encapsulated Thymidine Phosphorylase in Mitochondrial Neurogastrointestinal Encephalomyopathy

Michelle Levene<sup>1</sup>, Murray D Bain<sup>1</sup>, Nicholas F Moran<sup>2</sup>, Niran Nirmalanathan<sup>3</sup>, Joanna Poulton<sup>4</sup>, Mauro Scarpelli<sup>5</sup>, Massimiliano Filosto<sup>6</sup>, Hanna Mandel<sup>7</sup>, Andrew D MacKinnon<sup>3</sup>, Lynette Fairbanks<sup>8</sup>, Dario Pacitti<sup>9</sup> and Bridget E Bax<sup>1\*</sup>

<sup>1</sup>Molecular and Clinical Sciences, St. George's, University of London, UK

<sup>2</sup>Department of Neuroscience, East Kent Hospitals Foundation Trust, UK

<sup>3</sup>Departments of Neurology and Neuroradiology, St George's University Hospitals NHS Foundation Trust, UK

<sup>4</sup>Nuffield Department of Obstetrics and Gynaecology, University of Oxford, UK

<sup>5</sup>Institute of Neurology, University of Verona, Italy

<sup>6</sup>Centre for Neuromuscular Diseases, University Hospital "Spedali Civili", Italy

<sup>7</sup>Galilee Medical Center, Israel

<sup>8</sup>The Purine Research Laboratory, St Thomas' Hospital, UK

<sup>9</sup>Medical School, University of Exeter, UK

**\*Corresponding Author:** Bridget E Bax, Molecular and Clinical Sciences, St. George's, University of London, London, UK

Published **March 30, 2020**

This Book Chapter is a republication of an article published by Bridget E Bax, et al. at Journal of Clinical Medicine in April 2019. (Levene, M.; Bain, M.D.; Moran, N.F.; Nirmalanathan, N.; Poulton, J.; Scarpelli, M.; Filosto, M.; Mandel, H.; MacKinnon, A.D.; Fairbanks, L.; Pacitti, D.; Bax, B.E. Safety

and Efficacy of Erythrocyte Encapsulated Thymidine Phosphorylase in Mitochondrial Neurogastrointestinal Encephalomyopathy. *J. Clin. Med.* 2019, 8, 457.)

**How to cite this book chapter:** Michelle Levene, Murray D Bain, Nicholas F Moran, Niran Nirmalananthan, Joanna Poulton, Mauro Scarpelli, Massimiliano Filosto, Hanna Mandel, Andrew D MacKinnon, Lynette Fairbanks, Dario Pacitti, Bridget E Bax. Safety and Efficacy of Erythrocyte Encapsulated Thymidine Phosphorylase in Mitochondrial Neurogastrointestinal Encephalomyopathy. In: *Prime Archives in Medicine*. Hyderabad, India: Vide Leaf. 2020.

© The Author(s) 2020. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Author Contributions:** Conceptualization, B.E.B; methodology, B.E.B., M.L; investigation, M.L., M.D.B., N.F.M., N.N., J.P., M.S., M.F., H.M., AD.M., L.F., D.P., B.E.B; formal analysis, B.E.B; M.D.B, N.F.M; A.D.M; writing—original draft preparation, B.E.B; writing—review and editing, M.L., M.D.B., N.F.M., N.N., J.P., M.S., M.F., H.M., A.D.M.,L.F., D.P., B.E.B. ; supervision, B.E.B.; project administration, B.E.B.; funding acquisition, B.E.B.

**Funding:** This research was funded by grants awarded by the United Mitochondrial Disease Foundation (08-004) and the Purine Metabolic Patients Association. The supply of thymidine phosphorylase was supported by Medical Research Council Development Pathway Funding Scheme grants GO90217 and K025406.

**Acknowledgments:** The authors are thankful to the patients for their participation in this study.

**Conflicts of Interest:** B.E.B. serves on the scientific advisory board for Erytech Pharma, has received a travel grant from

Orphan technologies for the purpose of attending a rare disease conference in Israel and a license fee payment from Orphan Technologies. She has received research support from the Purine Metabolic Patients Association, Medical Research Council, United Mitochondrial Disease Foundation and the Higher Education Funding Council of England. M.D.B. has received a license fee payment from Orphan Technologies.

## Abstract

Mitochondrial neurogastrointestinal encephalomyopathy (MNGIE) is an ultra-rare autosomal recessive disorder of nucleoside metabolism caused by mutations in the nuclear *TYMP* gene, encoding for the enzyme thymidine phosphorylase. There are currently no approved treatments for MNGIE. The aim of this study was to investigate the safety, tolerability and efficacy of an enzyme replacement therapy for the treatment of MNGIE. In this single centre study three adult patients with MNGIE received intravenous escalating doses of erythrocyte encapsulated thymidine phosphorylase (EE-TP; dose range: 4 to 108 U/kg/4 weeks). EE-TP was well tolerated and reductions in the disease-associated plasma metabolites, thymidine and deoxyuridine were observed in all three patients. Clinical improvements, including weight gain and improved disease scores were observed in two patients, suggesting that EE-TP is able to reverse some aspects of the disease pathology. Transient, non-serious adverse events were observed in two of the three patients; these did not lead to therapy discontinuation and were managed with pre-medication prior to infusion of EE-TP. To conclude, enzyme replacement therapy with EE-TP demonstrated biochemical and clinical therapeutic efficacy with an acceptable clinical safety profile.

## Keywords

Mitochondrial Neurogastrointestinal Encephalomyopathy; MNGIE; *TYMP*; Enzyme Replacement; Thymidine Phosphorylase; Mitochondrial Disease; Rare Disease; Orphan Disease

## Introduction

MNGIE is an ultra-rare autosomal recessive metabolic disorder (Online Mendelian inheritance in Man #603041, Genome Database accession #9835128) that is almost universally fatal [1]. It causes relentless and progressive morbidity, followed by premature death at an average age of 37.6 years [2]. MNGIE is caused by mutations in the nuclear thymidine phosphorylase gene (*TYMP*) and a subsequent deficiency in thymidine phosphorylase activity (EC 2.4.2.4) [3,4]. The thymidine phosphorylase enzyme plays a pivotal role in the deoxyribonucleoside salvage metabolic pathway through catalysing the reversible phosphorylation of thymidine (also known as deoxythymidine) and deoxyuridine to 2-deoxyribose 1-phosphate and their respective bases, thymine and uracil [3]. In the absence of thymidine phosphorylase activity there is a systemic accumulation of thymidine and deoxyuridine which then generate imbalances within the mitochondrial deoxyribonucleotide pools, causing mitochondrial DNA (mtDNA) point mutations, depletion and deletion abnormalities, and ultimately mitochondrial dysfunction [5–9]. The clinical picture of MNGIE is mainly typified by gastrointestinal dysfunction with frequent intestinal sub-occlusive episodes, malnutrition, cachexia and neurological involvement including ptosis and ophthalmoparesis, peripheral neuropathy, myopathy, symmetric distal weakness which prominently affects the inferior extremities, and leukoencephalopathy [3,5,10].

No specific therapies for patients with MNGIE have been established in clinical trials. However, a number of experimental therapeutic approaches have been investigated which either directly remove the deoxyribonucleosides (haemodialysis and peritoneal dialysis), or introduce the deficient thymidine phosphorylase (platelet transfusions, allogeneic haematopoietic stem cell transplantation (AHSCT), autologous erythrocyte-encapsulated thymidine phosphorylase (EE-TP) and orthotopic liver transplantation) [6,11–20].

EE-TP is under clinical development by us and is formulated by the *ex vivo* encapsulation of *Escherichia coli* (*E.coli*) thymidine

phosphorylase within the patient's autologous erythrocytes using a reversible hypo-osmotic dialysis process. The enzyme-loaded erythrocytes are then returned to the patient. The rationale for this therapeutic approach is based on the facilitated diffusion of thymidine and deoxyuridine across the erythrocyte membrane via the equilibrative nucleoside transporter, ENT1. Once inside the erythrocyte, the encapsulated enzyme catalyses the metabolism of the deoxyribonucleosides to the normal products, thymine and uracil, which then exit the erythrocyte and enter the normal metabolic pathways. EE-TP aims to normalise the systemic accumulations of thymidine and deoxyuridine, and thereby ameliorate mitochondrial deoxyribonucleotide pool imbalances and slow disease progression. EE-TP has the pharmacological advantages of prolonging the circulatory half-life of the enzyme and potentially minimising immunogenic reactions which are frequently observed in enzyme replacement therapies administered by the conventional route [18,19,21].

Here, we describe three adult patients with MNGIE (Patients 1 to 3) who were treated with EE-TP under a compassionate use programme. Clinical and metabolic improvements were previously reported in Patient 2 who was treated with EE-TP for 27 months [19]; here we provide an update on the subsequent 49 months of treatment in this patient. EE-TP was shown to decrease the plasma metabolites in all three patients and clinical improvements, including weight gain and improved disease scores were observed in two patients.

## Experimental Section


*Study participants.* Female and male patients  $\geq 18$  years and of all ethnic backgrounds, on a world-wide basis were considered for study inclusion based on the following defined inclusion criteria: i) a definitive diagnosis of MNGIE due to thymidine phosphorylase deficiency established by the confirmation of a pathogenic mutation in *TYMP*,  $< 18\%$  of normal thymidine phosphorylase activity in the buffy coat, and plasma thymidine and deoxyuridine concentrations greater than  $3\mu\text{mol/L}$  and  $5\mu\text{mol/L}$ , respectively; ii) a whole blood haematocrit  $\geq 20\%$ ; and iii) patient signed informed consent. Exclusion criteria were: i)

pregnancy, planned pregnancy, breast feeding or unwillingness to practice contraception, ii) participation in a controlled trial of an investigational medicinal product within the previous 12 weeks, iii) patients who had received a successful bone marrow transplant and iv) patients with a known history of human immunodeficiency virus, hepatitis B infection, or an active hepatitis C infection.

*Study design and objectives.* This study was conducted in an open label manner with participants receiving EE-TP in accordance with the provisions of Schedule 1 of The Medicines for Human Use (Marketing Authorisations Etc.) Regulations SI 1994/3144, where Schedule 1 provides an exemption from the need for a marketing authorisation for a relevant medicinal product, which is supplied on an individual patient basis to fulfil a "special need". The study objectives were to assess the safety, tolerability and efficacy of EE-TP when administered to patients with MNGIE at dose-escalating intravenous infusions.

After a 28 day screening and baseline assessment period, participants received their first intravenous infusion of EE-TP. For safety reasons the study was initiated with the recruitment of a single patient (Patient 1), who was administered a low dose of EE-TP (dose level 1, 4 U/kg). This dose was labelled with sodium [<sup>51</sup>Cr] chromate for the purpose of evaluating the *in vivo* survival characteristics of the thymidine phosphorylase-loaded erythrocytes. In subsequent treatment cycles, the administered dose of EE-TP was cautiously escalated with the aim of achieving metabolic correction, provided the patient successfully completed safety reviews between administrations. For Patient 1 the dose was escalated in treatment cycle 2 to level 2 ( $9 \pm 2$  U/kg) and this was followed by dose escalations in treatment cycles 4 (dose level 3,  $18 \pm 2$  U/kg) and 9 (dose level 4,  $29 \pm 2$  U/kg), Figure 1. This dose escalation strategy was applied to Patients 2 and 3, but without sodium [<sup>51</sup>Cr] chromate-labelling, and employing higher starting doses, based on the safety and tolerability assessments made on Patient 1. Patient 2 initiated treatment with dose level 3 and this was escalated to dose level 4 in treatment cycle 3 and then to dose level 5 ( $47 \pm 2$  U/kg) in treatment cycle 24. Patient 3 initiated treatment with dose level

5, and this was escalated to dose level 6 ( $108 \pm 5$  U/kg) in treatment cycle 3. See Figure 1 for dose escalation scheme employed for Patients 1, 2 and 3.




**Figure 1:** Dose escalation scheme for Patients 1, 2 and 3.

Following a screening period of 28 days an intravenous dose of EE-TP was administered on day 0. Patients were escalated to the next dose following a safety review. Blue numbers refer to dose levels, where level 1 = 4 U/kg, level 2 = 9 U/kg, level 3 = 18 U/kg, level 4 = 29 U/kg, level 5 = 47 U/kg and level 6 = 108 U/kg. Note that data pertaining to dose level 5 for Patient 2 is only covered in this report.

*Preparation and administration of EE-TP.* The study investigational product, EE-TP was prepared by the investigator team, using sterile, single-use materials and reagents throughout, under a Specials licence held by St. George's Healthcare Trust Pharmacy according to the Rules and Guidance for Pharmaceutical Manufacturers 2007 (MHRA), within Class A isolators contained within a Pharmacy Clean room. EE-TP was formulated by the encapsulation of thymidine phosphorylase within the patient's own autologous erythrocytes, *ex-vivo*, using a reversible hypo-osmotic dialysis process[21]. Briefly, a predetermined volume of the patient's blood was collected (55 to 400 mL; the volume collected being dependent on the activity of thymidine phosphorylase to be encapsulated and the patient's haematocrit) using aseptic techniques into tubes containing the low molecular weight heparin, dalteparin sodium (10 units/mL

blood). Within cleanroom facilities, the blood components were separated by centrifugation at 1,100 *g* for 10 minutes, followed by removal of the plasma and buffy coat (both retained for later use). The erythrocytes were then washed twice in cold (4°C) phosphate buffered saline (PBS; 136.89 mmol/L NaCl, 2.68 mmol/L KCl, 8.10 mmol/L Na<sub>2</sub>HPO<sub>4</sub>, 1.47 mmol/L KH<sub>2</sub>PO<sub>4</sub>, pH 7.4) with centrifugation. Seven volumes of washed and packed erythrocytes were mixed with three volumes of cold PBS containing 6,000 to 38,000 Units recombinant *E.coli* thymidine phosphorylase (Diatheva, Italy for treatment cycles 68-79 in patient 2, otherwise Sigma, Israel) and the suspension placed into dialysis bags with a molecular weight cut-off of 12,000 Daltons. The cells were dialysed against hypo-osmotic buffer (5 mmol/L KH<sub>2</sub>PO<sub>4</sub>, 5 mmol/L K<sub>2</sub>HPO<sub>4</sub>, pH 7.4) at 4°C in a refrigerated incubator, with rotation at 6 rpm, for 120 minutes. These conditions of hypo-osmolarity induce the erythrocytes to swell due to an influx of water, until at a critical size, pores form in the erythrocyte membrane; whilst permeable, thymidine phosphorylase enters the erythrocytes by diffusion. Erythrocyte resealing was achieved by restoration of iso-osmotic conditions by transferring the dialysis bags to pre-warmed iso-osmotic PBS supplemented with 5 mmol/L adenosine, 5 mmol/L glucose and 5 mmol/L MgCl<sub>2</sub>, pH 7.4, and rotated at 6 rpm for 60 minutes in an incubator set at 37°C. The resealed erythrocytes were washed three times in three volumes of supplemented PBS with centrifugation at 100 *g* for 20 min. The washed and packed EE-TP was mixed with the retained buffy coat, and re-suspended in an equal volume of plasma. EE-TP was administered to the patient by slow intravenous infusion within 45 minutes of manufacture. The manufacturing steps for EE-TP preparation are detailed in Figure 2.



**Figure 2:** Manufacture of EE-TP. Following venesection, blood was transferred to a Class A isolator for the manufacture of EE-TP under a Specials according to the Rules and Guidance for Pharmaceutical Manufacturers 2007 (MHRA). Blood was centrifuged to separate components and then the erythrocytes washed with PBS. Erythrocytes were then mixed with an appropriate activity of thymidine phosphorylase (TP) to a haematocrit of 70% and then dialysed against hypo-osmotic buffer for 90 minutes to create pores in the cell membrane. The lysed erythrocytes were then resealed by dialysis against iso-osmotic buffer for 60 minutes to encapsulate TP that had entered the cells. The resulting EE-TP was then washed to remove unencapsulated TP, mixed with the retained buffy coat and an equal volume of autologous plasma, and then infused into the patient.

*In vivo survival studies of EE-TP.* Thymidine phosphorylase-loaded erythrocytes were labelled using a standard  $^{51}\text{Cr}$  erythrocyte-labelling technique [22]. The washed and packed EE-TP prepared as described above was gently mixed with 0.75 MBq of sodium [ $^{51}\text{Cr}$ ] chromate (Amersham International, Buckinghamshire, UK), and allowed to stand at room temperature for 30 minutes. Unbound chromium was removed by the addition of 100 mg of ascorbic acid (100 mg/mL, Evans Medical, Leatherhead, UK) to the cell suspension, followed by a single wash in supplemented PBS. After resuspension in an equal volume of autologous plasma, EE-TP was slowly injected into patient 1 over a period of 10 minutes.

*In vivo survival* was assessed by monitoring the disappearance of  $^{51}\text{Cr}$  label from the circulation; 10 mL blood samples were taken from a vein in the contralateral arm 3, 6, 8, 14 days and then weekly until activity was not noticeably above background. To

check for intravascular haemolysis,  $^{51}\text{Cr}$  activity was measured in plasma. Percentage raw cell survival was calculated by expressing CPM/mL packed cells as a percentage of the calculated zero time value, after correction for natural decay. The data were then plotted against time on a semi-logarithmic scale and a best-fit line fitted.  $T_{1/2}$  thymidine phosphorylase-loaded erythrocyte survival was taken as the time for the concentration of  $^{51}\text{Cr}$  in the circulating blood to fall to 50% of its initial value. For the determination of mean cell life, CPM/mL packed cells (corrected for both natural decay and chromium elution from the erythrocytes) was expressed as a percentage of the corrected zero time value. Cell survival data were plotted against time on a linear scale and the mean cell life span derived from the intercept obtained by extrapolation of the line to zero activity.

Urinary excretion of label was assessed by making three 24-hour urine collections for the first 72 hours after injection;  $^{51}\text{Cr}$  activity in the urine was expressed as a percentage of the total  $^{51}\text{Cr}$  activity injected.

*Patient assessments:* Safety assessments included documentation of all adverse events, vital signs and standard clinical laboratory evaluations, including haematology and blood chemistry. Vital signs were monitored for 2 hours following administration of EE-TP, for the first three treatment cycles and after each dose escalation. Clinical laboratory evaluations were performed following the end of each treatment cycle, prior to the administration of EE-TP in the following treatment cycle.

To assess the metabolic efficacy of EE-TP in reducing or eliminating plasma thymidine and deoxyuridine, the disease associated metabolites, 5 mL of whole blood was collected into EDTA vacutainers at pre-therapy baseline (day 0) and then 5 days following each administration of EE-TP. The assessment of serum anti-thymidine phosphorylase antibodies for immunogenicity assessment was conducted by collecting 4 mL of whole blood into SST serum separation vacutainers at pre-therapy baseline (day 0) and then prior to each administration of EE-TP.

The clinical efficacy of EE TP was assessed by following longitudinal changes from baseline in body weight measurements, MRI scan (patient 2 only), disease scoring scales (MRC sum score, Overall neuropathy limitations scale (ONLS), Newcastle mitochondrial disease scale (NMDS) and Sensory sum score) and the Short Form (36) Health Survey (SF36).

*Analytical methods.* Plasma and urine thymidine and deoxyuridine levels were measured using Ultra-Performance Liquid Chromatography (UPLC). Briefly 200 $\mu$ L of plasma were mixed with 200  $\mu$ L 10% TCA, vortexed and then centrifuged for 2 minutes at 12,000 rpm. The supernatant was then washed four times with water saturated diethyl-ether, after which the samples were left to stand for 5 minutes to evaporate off any residual ether. Urine samples were prepared by diluting 1:31 with deionised water. Two microliters of processed plasma or urine sample were injected onto a UPLC column. Chromatographic separation of thymidine and deoxyuridine was achieved using reversed phase chromatography with gradient elution using a Waters Alliance UPLC system with UV detection. A pre-packed C18 column (Vanguard Pre Column BEH C18 1.7  $\mu$ m, 2.1x 5 mm) was used as the stationary stage. Analytes were eluted using a gradient of Buffer A (40 mM ammonium acetate, pH 5.0) and Buffer B (100% v/v Methanol), which was set at 100% Buffer A for 3.5 minutes followed by a linear gradient to 20% Buffer B at 12 minutes, at a flow rate of 1.0 mL/minute. Total elution time was 15 minutes at 20 $^{\circ}$  C with UV detection at 270 nm and 0.1 absorbance units full scale. Metabolites were identified by comparing spectra with pure standards. The retention times of deoxyuridine and thymidine were 9.29 and 11.76 minutes, respectively.

Thymidine phosphorylase activity was measured in an aliquot of each batch of EE-TP manufactured to ascertain the dose administered using a validated high performance liquid chromatography (HPLC) assay [23].

Anti-thymidine phosphorylase antibodies were measured using a validated thymidine phosphorylase assay [24]. Samples were screened for a positive or negative signal; for samples that

screened positive, the specificity of thymidine phosphorylase was determined in a confirmatory assay with a specificity cut-point of 93% inhibition.

*Study approval.* The study was approved by the National Research Ethics Service Committee. Participants provided written informed consent before participating in any study procedures.

## Results

### Patient Characteristics

Three patients between the ages 25 and 28 years were recruited into the study. The patients fulfilled the recruitment criteria through harbouring pathogenic mutations in the thymidine phosphorylase gene, having a deficiency of thymidine phosphorylase activity in the buffy coat, and having plasma thymidine and deoxyuridine concentrations greater than 3  $\mu\text{mol/L}$  and 5  $\mu\text{mol/L}$ , respectively. Table 1 shows demographic and baseline disease data. Clinical manifestations on recruitment for Patients 1 and 3 were peripheral polyneuropathy, external ophthalmoplegia, intestinal dysmotility, anorexia and cachexia, and for Patient 2 peripheral polyneuropathy and external ophthalmoplegia with minimal intestinal involvement.


**Table 1:** Subject demographics and baseline disease characteristics.

Patient	Sex	Age of diagnosis (years)	Age at start of EE-TP	TYMP mutation	Buffy coat TP activity (nmol/mg protein/hour) <sup>1</sup>	Pre-treatment plasma metabolites (µmol/L)	
						Thymidine	Deoxyuridine
1	Female	23	25	Homozygous for c.1431_1432insT	40	9	19
2	Male	26	28	Heterozygous c.866A>C c.1231_1243 del	0	21	31
3	Male	15	26	g.4009_4010insG g.4101G>A	24	12	19

<sup>1</sup>Normal range: 336 -1341 nmol/h/mg Protein

## In Vivo Survival of EE-TP

To ascertain that the encapsulation of thymidine phosphorylase had no effect on the *in vivo* survival of erythrocytes, the circulatory lifespan of the first EE-TP dose administered to patient 1 was estimated by <sup>51</sup>Cr labelling. The mean cell life and mean cell half-life ( $t_{1/2}$ ) were calculated to be 108 and 32 days, respectively, both within the normal reference range (Figure 3) and the mean daily urinary excretion of label was 0.9% (1.1% at 0-24 hours, 0.8% at 24-48 hours and 0.7% at 48-72 hours), also within the normal <sup>51</sup>Cr elution limits of 1.0 -3.2% per day (Table 2). No label was detected in the plasma demonstrating minimal intra-vascular haemolysis of the enzyme loaded cells.



**Figure 3:** *In vivo* survival of EE-TP in Patient 1. The first administered dose of EE-TP (4U/kg body weight) was labelled with <sup>51</sup>Cr and the survival characteristics of the loaded erythrocytes assessed by following the circulation.

**Table 2:** Urinary excretion of <sup>51</sup>Cr after infusion of EE-TP in Patient 1.

Urine collection period (hours)	Urinary excretion of <sup>51</sup> Cr (%)	Reference range (%)
0 - 24	1.1	1.0 – 3.2
24 - 48	0.8	
48 - 72	0.7	
Mean	0.9	


## Efficacy Outcomes

### Patient 1


Thirty one treatment cycles of EE-TP (including the chromium labelling study) were administered over a period of 28 months. EE-TP was administered at ascending dose levels of 4 , 9, 18 and 29 U/kg body weight/ 4 weeks up to treatment cycle 24; this was followed the administration of 14 U /kg body weight/ 2 weeks between treatment cycles 25 to 31 (Figure 4a). Pre-treatment plasma concentrations of thymidine and deoxyuridine were 10  $\mu\text{mol/L}$  and 20.0  $\mu\text{mol/L}$ , respectively; between 270 and 620 days from the start of therapy, these were reduced to intra cycle concentrations of 2-6  $\mu\text{mol/L}$  for thymidine and 3-13 $\mu\text{mol/L}$  for deoxyuridine (Figure 4b). From day 320 onwards, the urinary excretion of metabolites decreased from pre-therapy levels of 73 and 118  $\mu\text{mol}/24$  hours for thymidine and deoxyuridine, respectively, to 0-41 $\mu\text{mol}/24$  hours for thymidine and 0-49 $\mu\text{mol}/24$  hours for deoxyuridine (Figure 4c). By 200 days of therapy, the patient had gained 4 kg in weight (Figure 4d). This coincided with a decrease in nausea and vomiting, an increased walking distance and an increase in the physical and mental components of the SF36 health and well-being survey from 47 and 43, respectively, at pre-therapy, to 52 and 45 at day 189 (7 months). No changes were noted in the disease scoring scales at 7 months compared to pre-therapy scores (Table 3).

At day 220, the patient developed a flu-like illness and this was followed by a diagnosis of small intestinal bacterial overgrowth which lead to a subsequent 4 kg weight loss (Figure 4d). At day 252 transparenteral nutrition (TPN) was commenced with the administration of 2000 mL of lipid-based formula three times per week (500 Kcal medium-chain triglycerides and 1200 Kcal glucose) and 2000 mL of lipid-free formulation (1400 Kcal glucose) four times per week. Although this resulted in an initial 6 kg increase in weight up to day 686 (Figure 4d), the activities of alanine minotransferase (ALT) and gamma-glutamyl transpeptidase (GGT) became elevated, in a pattern indicative of hepatocellular damage (Figure 5). From day 620, the plasma metabolite levels increased to levels equal or greater than those determined pre-therapy (Figure 4b) and this correlated with the

development of lipaemia and lipiduria in association with the continuation of TPN. There was a progression of the peripheral polyneuropathy and a deterioration in the NMDS general neurological functioning and clinical assessment components, the MRC score, the sensory sum score and the SF36 health and well-being survey compared to assessments recorded pre-treatment and 7 months of treatment with EE-TP (Table 3). In an attempt to lower the metabolite levels, the frequency of treatment cycles was increased to fortnightly (14 U/kg/ two weeks) from treatment Cycle 26 onwards. This had no effect on the metabolite levels and the patient died from general debilitation at 20 days after the last administration of EE-TP.



**Figure 4:** Dose of EE-TP administered, plasma concentrations of deoxyribonucleosides, urinary excretion of deoxyribonucleosides and body weight in Patient 1 during 28 months of therapy with EE-TP. a) The dose of encapsulated thymidine phosphorylase infused was determined using HPLC. b) Mid cycle plasma thymidine (blue data points) and deoxyuridine (red data points) concentrations were measured by UPLC. c) Mid cycle urinary excretions of thymidine (blue data points) and deoxyuridine (red data points) were determined by measuring the concentration of metabolites in 24 hour urine collections. d) Patient body weight. The blue shaded area represents the timeframe in which TPN was administered.



**Figure 5:** Serum ALT and GGT activities in patient 1 during 31 treatment cycles of EE-TP administered over 28 months. Blue symbols represent GGT and red symbols ALT.


**Table 3:** Quality of Life and Disease scores for Patient 1, pre-therapy and during EE-TP therapy.

Scale	Months of therapy		
	Pre-therapy	7	24
SF36 (health and well-being):			
Physical component (population mean 50 ± 10)	47	52	42
Mental component (population mean 50 ± 10)	43	45	42
Newcastle mitochondrial disease scale (normal = 0):			
I (general neurological functioning, 0 to 50)	4	4	16
II (system specific functioning, 0 to 45)	6	6	6
III (clinical assessment, 0 to 50)	8	8	19
MRC sum motor score (normal = 80)	73	73	53
Overall neuropathy limitation scale: (0 = no disability, 12 = maximum disability)	4	4	4
Sensory sum score: (normal = 0, maximum score = 64)	26	26	34

## Patient 2

A total of 79 treatment cycles of EE-TP were administered over of period of 76 months. Data obtained for the first 27 months of treatment (cycles 1 to 30) is reported elsewhere [19]. Between

months 27 and 76, EE-TP doses of 47 U/kg per 4 weeks (dose level 5) were administered. Pre-treatment plasma concentrations of thymidine and deoxyuridine were 20.5  $\mu\text{mol/L}$  and 30.6  $\mu\text{mol/L}$ , respectively, and these were reduced to intra cycle values of 0-9  $\mu\text{mol/L}$  for thymidine and 0-15  $\mu\text{mol/L}$  for deoxyuridine (Figure 6a). These were lower than the levels previously reported for the first 27 months of therapy [19]. The intra-cycle urinary excretion rates of thymidine and deoxyuridine ranged between 0 - 354  $\mu\text{mol/24 hours}$  for thymidine and 4 - 218  $\mu\text{mol/24 hours}$  for deoxyuridine (Figure 6b); these were generally lower than reported for the first 27 months of therapy [19]. Pre-therapy excretion rates were 421  $\mu\text{mol/24 hours}$  for thymidine and 324  $\mu\text{mol/24 hours}$  for deoxyuridine, respectively. The previously reported 5.8 kg gain in body weight was sustained for a total of 23 months until day 1162, after which there was a 6 kg weight loss following a flu-like illness. Body weight then increased to 59 kg but then declined to 55.5 kg after 11 months, following a further flu-like illness (Figure 6c). The formerly reported decline in plasma creatine kinase activity, from a pre-therapy activity of 1200 U/L to activities very near or within the normal reference range of 40 - 320 U/L, were sustained throughout this current reporting period (Figure 6d).




**Figure 6:** Plasma concentrations of deoxyribonucleosides, urinary excretion of deoxyribonucleosides, body weight and plasma creatine kinase in Patient 2 between 27 and 76 months of therapy with EE-TP. a) Intra-cycle plasma thymidine and deoxyuridine concentrations were measured by UPLC. b) Intra-cycle urinary excretions of thymidine and deoxyuridine were determined by measuring the concentration of metabolites in 24 hour urine collections. Red dotted lines represent pre-therapy deoxyuridine plasma concentration and urinary excretion, and blue dotted lines represent pre-therapy thymidine plasma concentrations and urinary excretion. c) Patient body weight. d) Plasma creatine kinase activity. Pink dotted line represents pre-therapy activity.

Scores for the physical and mental components of the SF36 health and well-being survey increased from 52 and 59, respectively, at 23 months, to 55 and 60 at 50 months of therapy. The scores remained at these levels at 64 and 73 months (Table 4). Clinical assessments 64 months after initiating treatment with EE-TP demonstrated further improvements in sensory ataxia, balance and gait and fine finger functioning compared with assessments recorded at 23 months of therapy. The first improvement in the general neurological functioning component of the NMDS was recorded at 73 months of therapy, along with an improvement in the clinical assessment component at 64 months and a further improvement at 73 months of therapy. A decline in the system specific functioning component of the NMDS was noted at 64 months of therapy, compared with previous recorded scores. However, this was followed by a small improvement at 73 months of therapy. The MRC sum score for power remained unchanged since 23 months, with the patient demonstrating normal power in his right little finger and thumb. The score for the ONLS scale declined at 64 months of therapy, compared to scores recorded at pre-therapy and up to 50 months of therapy. There were improvements of 7 and 16 points on the Sensory sum score, respectively at 64 and 73 of months of therapy, compared to the score recorded at 23 months of therapy (Table 4). Patient reported outcomes correlated with neurological assessments and included improvements in the distal sensation in the upper and lower limbs and being able to hold a focused conversation whilst standing, whereas previously his lack of balance made visual engagement difficult.

**Table 4:** Quality of Life and Disease scores for Patient 2, pre-therapy and during therapy with EE-TP.

Scale	Months of therapy				
	Pre-therapy	23	50	64	73
SF36 (health and well-being):					
Physical component (population mean $50 \pm 10$ )	48	52	55	55	55
Mental component (population mean $50 \pm 10$ )	60	59	60	60	60
Newcastle mitochondrial disease scale (normal = 0):					
I (general neurological functioning, 0 to 50)	4	4	4	4	2
II (system specific functioning, 0 to 45)	2	2	2	6	5
III (clinical assessment, 0 to 50)	11	11	11	9	8
MRC sum motor score (normal = 80)	56	74*	74	74	74
Overall neuropathy limitation scale: (0 = no disability, 12 = maximum disability)	3	3	3	4	4
Sensory sum score: (normal = 0, maximum score = 64)	21	19	20	13	4

Brain MRI at month 28 of therapy showed almost symmetric, patchy areas of FLAIR and T2 hyperintensity in the cerebral periventricular, deep and subcortical white matter (Figure 7, Panel a). A follow-up brain MRI at 63 months showed a progression of T2/FLAIR hyperintense cerebral white matter changes (Figure 7, Panel b). Relative sparing of the U fibres and the corpus callosum was noted (Figure 7, Panel C left image). Infratentorially there was minor symmetrical FLAIR hyperintense signal change in the cerebellar peridentate white matter (Figure 7, Panel c right image).




**Figure 7:** MRI brain of Patient 2. MRI (axial FLAIR) at a) 28 months and b) 63 months shows progression of T2/FLAIR hyperintense cerebral white matter changes (leukoencephalopathy). The scan at 28 months shows patchy involvement of cerebral periventricular, deep and subcortical white matter (a) which 35 months later is more confluent and extensive (b). There is relative sparing of the U-fibers and the corpus callosum (c) left image) and infratentorially there is minor symmetrical FLAIR hyperintensity in the cerebellar peridentate white matter (c) right image).

### Patient 3

Four treatment cycles were administered over a period of 83 days. The initial EE-TP dose administered was 43 U/kg and this was escalated to 108 U/kg every 4 weeks in treatment cycles 3

and 5, respectively (Figure 8a). Pre-treatment plasma concentrations of thymidine and deoxyuridine were 12  $\mu\text{mol/L}$  and 19  $\mu\text{mol/L}$ , respectively; these declined to intra-cycle values of less than 4  $\mu\text{mol/L}$  and 2  $\mu\text{mol/L}$  for thymidine and deoxyuridine, respectively from 60 days onwards (Figure 8b). By day 3 of treatment, the intra-cycle urinary excretion rates of thymidine and deoxyuridine rapidly declined from 308 and 448  $\mu\text{mol}/24$  hour pre-therapy to levels less than 21  $\mu\text{mol}/24$  hours for deoxyuridine and 30 to 89  $\mu\text{mol}/24$  hours for thymidine (Figure 8c). Longitudinal assessments of disease scoring and the SF36 health and well-being survey were not conducted due to the premature termination of therapy due to the patient's inability to sustain long distance travel to the UK once per month to receive treatment. The patient's weight remained unchanged from the pre-therapy measure of 31.7 kg.



**Figure 8:** Dose of EE-TP administered, plasma concentrations of deoxyribonucleosides and urinary excretion of deoxyribonucleosides in Patient 3 during 6 months of therapy with EE-TP. a) The dose of encapsulated thymidine phosphorylase infused was determined using HPLC. b) Intra-cycle plasma thymidine (blue data points) and deoxyuridine (red data points) concentrations were measured by UPLC. c) Intra-cycle urinary excretions of thymidine (blue data points) and deoxyuridine (red data points) were determined by measuring the concentration of metabolites in 24 hour urine collections.

## Safety

Adverse reactions were observed in two of the three patients, Patients 1 and 2 in treatment cycles 1-17 and 1-11, respectively. These were short-lived, occurring within 5 minutes of infusion, and are described in Tables 5 and 6 for Patients 1 and 2, respectively. The reactions were managed by the administration of antihistamine and anti-inflammatory therapy, prior to EE-TP administration, with the addition of anti-emetics in Patient 1. It is noteworthy that the introduction of the highly purified GMP enzyme preparation produced by Diatheva in treatment Cycle 68 (Patient 2), permitted a gradual withdrawal of the pre-medication. Intravenous hydrocortisone (100 mg) was, however, re-introduced in treatment cycles 77- 79 as a precaution after the patient experienced chills three hours after infusion of EE-TP in the previous treatment cycle.

**Table 5:** Adverse reactions to EE-TP administration in Patient 1.

Infusion #	Observations	Pre-medication prior to infusion
2	nausea felt hot and faint erythema of face, neck and in arm proximal to infusion site tickle sensation in throat subsided spontaneously after 5 minutes	None
3	coughing spasm sensation in throat described as a tickle dry mouth felt hot subsided spontaneously after 5 minutes	Oral chlorphenamine 4mg 3 hrs prior
4	coughing spasm	Oral chlorphenamine 8 mg 3 hrs prior
5	generalized paresthesia nausea coughing spasm generalized erythema transient headache, visual disturbance subsided spontaneously after 5 minutes	Oral chlorphenamine 8 mg 3 hrs prior IV hydrocortisone 50 mg at time of reaction
6	no reactions	Oral dexamethazone 4 mg 6 hrs prior Oral chlorphenamine 8 mg 3 hrs prior IV maxalon 10 mg immediately prior to infusion
9	coughing spasm nausea felt hot	As infusion 6
10	nausea	Oral dexamethazone 4 mg 6 hrs prior Oral chlorphenamine 8 mg 3 hrs prior Oral ondansetron 4 mg 3 hrs prior
15	coughing spasm nausea felt cold and shaky	As infusion 10
17	coughing spasm nausea erythema of neck	Oral dexamethazone 8 mg 6 hrs prior Oral chlorphenamine 12 mg in two divided doses: 6 hrs (8 mg) and 2 hrs (25%) prior Oral ondansetron 8 mg 3 hrs prior
18-31	no reactions	Oral dexamethazone 10 mg 6 hours prior Oral chlorphenamine 12 mg two divided doses: 6 hours (8 mg) and 2 hrs (4 mg) prior Oral ondansetron 16 mg total in two equal doses, 6 and 3 hrs prior

**Table 6:** Adverse reactions to EE-TP administration in Patient 2.

<b>Infusion #</b>	<b>Observations</b>	<b>Pre-medication prior to infusion</b>
3	erythema of face felt hot	none
4	erythema of face felt hot	oral dexamethasone 4 mg 6 hrs prior oral chlorphenamine 12 mg , 4 mg and 8 mg given respectively 4 hrs and 1 hr prior oral ondansetron 8 mg given 1 hr prior
5	erythema of face felt hot	oral dexamethasone 10 mg 6 hrs prior oral chlorphenamine 12 mg, 4 mg and 8 mg given respectively 4 hr and 1 hr prior to infusion oral ondansetron 8 mg given 1 hr prior
6	erythema of face felt hot tightness in chest	oral dexamethasone 12 mg 6 hrs prior oral chlorphenamine 12 mg, 4 mg and 8 mg given respectively 4 hrs and 2 hrs prior to infusion + IV chlorphenamine 8 mg at start of infusion oral ondansetron 8 mg given 1 hr prior
7-8	felt hot erythema of face reduced compared to previous cycles	oral dexamethasone 10 mg 6 hrs prior IV hydrocortisone 100 mg at start of infusion oral chlorphenamine 12 mg, 2 hrs prior to infusion and IV 8 mg at start of infusion oral ondansetron 8 mg given 1 hr prior
9	slight erythema of face felt hot	oral monteleukast 20 mg daily for 3 days prior oral dexamethasone 10 mg 6 hrs prior IV hydrocortisone 100 mg at start of infusion oral chlorphenamine 12 mg, 2 hrs prior to infusion and IV 8 mg at start of infusion oral ondansetron 6 mg 1 hr prior
10	slight erythema of face felt hot	oral monteleukast 20 mg daily for 3 days prior to infusion oral ketotifen 2 mg daily in 2 equally divided doses for 4 days prior to infusion oral dexamethasone 10 mg 6 hrs prior oral chlorphenamine 12 mg, 2 hrs prior oral ondansetron 8 mg given 1 hr prior IV hydrocortisone 100 mg at start of infusion. IV 8 mg at start of infusion
11	very slight erythema of face	as infusion 10
12-13	No reactions	oral dexamethasone 10 mg 6 hrs prior oral chlorphenamine 12 mg, 2 hrs prior oral ondansetron 4 mg given 1 hr prior IV hydrocortisone 100 mg at start of infusion IV 8 mg at start of infusion
14-22	No reactions	phased withdrawal of pre-medication starting with ondansetron, then oral dexamethasone and chlorphenamine
23-26	No reactions	IV chlorphenamine 4 mg at start of infusion IV hydrocortisone 100 mg at start of infusion
27	very slight erythema of face	IV chlorphenamine 8 mg at start of infusion

28-39	No reactions until infusion 39 when during last 5 minutes of infusion: erythema of face, tightness in chest	IV chlorphenamine 8 mg at start of infusion IV hydrocortisone 50 mg at start of infusion
40-68	No reactions	oral dexamethasone 10 mg 5 hrs prior to infusion oral chlorphenamine 4 mg 4 hrs prior to infusion IV chlorphenamine 10 mg at start of infusion IV hydrocortisone 100 mg at start of infusion
69-	No reactions	oral dexamethasone 10 mg 5 hrs prior to infusion oral chlorphenamine 4 mg 4 hrs prior to infusion IV chlorphenamine 10 mg at start of infusion IV hydrocortisone 50 mg at start of infusion
70	No reactions	oral dexamethasone 10 mg 5 hrs prior to infusion oral chlorphenamine 4 mg 4 hrs prior to infusion IV chlorphenamine 10 mg at start of infusion
71	No reactions	IV chlorphenamine 5 mg at start of infusion IV hydrocortisone 50 mg at start of infusion
72	No reactions	oral dexamethasone 10 mg 4 hrs prior to infusion
73	No reactions	oral dexamethasone 6 mg 4 hrs prior to infusion
74-75	No reactions	oral dexamethasone 4 mg 4 hrs prior to infusion
76	Chill 3 hour post infusion	none
77-79	No reactions	IV hydrocortisone 100 mg at start of infusion

Immunogenicity, i.e. the presence of anti-thymidine phosphorylase antibodies was positive in the serum samples from 2 patients in the screening assay (treatment cycle 8 for Patient 1, and treatment cycles 9 onwards for Patient 2). The confirmatory assay revealed that only the positive samples from Patient 2 were specific for thymidine phosphorylase antibodies. There were no correlations between the plasma levels of deoxyribonucleosides and the appearance of anti-thymidine phosphorylase antibodies.

No clinically significant alterations in vital signs were reported. No abnormalities of haematological or clinical chemistry parameters were observed (other than those reported above).

## Discussion

MNGIE is a relentlessly progressive, degenerative disease with a very poor prognosis, and causes a great deal of morbidity for affected individuals. Gastrointestinal dysmotility caused by degeneration of the alimentary peripheral nervous system occurs in nearly all patients. The resulting symptoms problems include early satiety, dysphagia, nausea and vomiting after eating, episodic abdominal distention and pain, diarrhoea and severe weight loss. Peripheral polyneuropathy leads to distal weakness and sensory symptoms and may be disabling. Ptosis, external ophthalmoplegia and hearing loss are also common [1]. At present there is no regulatory approved treatment to prevent or reverse the inexorable clinical deterioration.

A number of experimental therapies have been explored which share the therapeutic strategy of reducing or eliminating the systemic concentrations of thymidine and deoxyuridine with the ultimate aim of ameliorating intracellular deoxyribonucleotide imbalances and preventing further mtDNA damage [6,11–19]. The permanent restoration of thymidine phosphorylase activity has been provided through AHSCT and more recently orthotopic liver transplantation [14–17,20]. A retrospective analysis of patients who received AHSCT showed that 9 of 24 patients (37.5%) were alive at follow-up, with survival being strongly associated with human a leukocyte antigen match of 10/10 and an absence of liver disease and gastro-intestinal pseudo-

obstruction [17]. Death was generally caused by transplantation complications or disease progression, and consequently a published consensus conference proposal recommends that only patients with an optimally matched donor and in good clinical condition should be considered for treatment using AHSCT [14]. Patients with MNGIE frequently experience delays in attaining the correct diagnosis, during which time their clinical condition often deteriorates; a majority of patients are therefore ineligible for AHSCT. A clinical trial, sponsored by Columbia University, USA is currently recruiting patients with MNGIE to evaluate the safety of AHSCT (NCT02363881). Orthotopic liver transplantation for MNGIE has been reported in two patients without complications and with sustained normalisation of thymidine and deoxyuridine (follow-up 90 days to 18 months) [20]. However, a longitudinal evaluation of data collected from additionally transplanted patients is necessary to confirm the efficacy of this treatment approach.

We report here the application of an enzyme replacement therapy for MNGIE which uses a genetically engineered recombinant *E.coli* thymidine phosphorylase sharing a 40% sequence homology with the human sequence [25]. The enzyme is encapsulated into the patient's autologous erythrocytes *ex vivo* to produce EE-TP which is then administered to the patient. In these studies the encapsulation process was conducted using a centralised manual manufacturing process taking 7 to 8 hours from the time of venesection. Consequently, patients were required to attend the study site on the day of treatment for venesection in the morning for EE-TP manufacture and then infusion later in the day. Two of the patients travelled from abroad every 4 weeks to receive treatment. For clinical trials, a decentralised manufacturing process for EE-TP will be employed using an automated device, thereby assuring consistency in the EE-TP drug product and permitting treatment to be expanded to the patient's locality. Patient 2 was transitioned to the automated EE-TP manufacturing process in treatment cycle 80, along with a fourth patient who was recruited into this compassionate programme.

Erythrocyte-mediated enzyme replacement therapy is applicable to disorders where the pathological plasma metabolite is able to

permeate the erythrocyte membrane. EE-TP aims to clear the associated MNGIE disease metabolites, thymidine and deoxyuridine, from the extracellular and cellular compartments, leading to an amelioration of the intracellular nucleotide imbalances and ultimately arrest/reverse the progression of the clinical disease by reversal of the mitochondrial dysfunction in MNGIE. The encapsulation within erythrocytes has the benefit of prolonging the circulatory half-life of thymidine phosphorylase, thereby reducing the frequency of EE-TP required, and potentially minimising immunogenic reactions against the *E.coli* enzyme. Indeed, the viability of the erythrocyte as a vehicle for sustaining therapeutic blood levels of thymidine phosphorylase was demonstrated in Patient 1, where the first dose of EE-TP was labelled with <sup>51</sup>Cr. The circulating mean cell life, and mean cell half- life were 108 days and 32 days, respectively, and well within the normal mean cell life range of 89 to 131 days and mean cell half-life range of 19 to 29 days [22]. The absence of <sup>51</sup>Cr in the plasma and the finding that the urinary excretion of <sup>51</sup>Cr was within the normal Cr elution limit of 1.0 to 3.2% per day indicate that the label was cell associated and there was minimal intravascular haemolysis of EE-TP on infusion .

We investigated the safety, tolerability and efficacy of EE-TP in three adult patients with MNGIE. Findings from the initial 27 months of treating Patient 2 with EE-TP were published previously and so we report here an update on the subsequent 49 months of treatment of this patient with EE-TP [19]. Six dose levels of EE-TP were investigated, ranging from 4U/kg to 108 U/kg every 4 weeks. A cautious escalation strategy was employed by staggering recruitment, starting with Patient 1 who was administered dose levels 1 to 4 (4-29 U/kg/4 weeks), followed by Patient 2 who was administered dose levels 3 to 5 (18-47 U/kg/4 weeks) and then Patient 3 who was administered dose levels 5 and 6 (47 and 108 U/kg/4 weeks). The decision to escalate to the next dose was based on the patient successfully completing a safety review between EE-TP administrations and the objective of further reduction of the disease plasma metabolites, thymidine and deoxyuridine. Although dose levels of 4 and lower provided some lowering of plasma metabolite

concentrations, dose levels 5 and 6 provided superior reductions in Patients 2 and 3, with intra-cycles plasma thymidine and deoxyuridine often decreasing to below the diagnosis concentrations for MNGIE. Dose levels of 4 and above were able to significantly reduce the urinary excretion of thymidine and deoxyuridine in all three patients.

Minor clinical improvements were observed in Patient 1 and more substantial improvements in Patient 2. Both patients demonstrated several periods of weight gain but also periods of weight loss associated with intercurrent illnesses. TPN was commenced in Patient 1 and although this resulted in an initial increase in weight, it was subsequently followed by weight loss, coinciding with a sharp increase in plasma metabolites, an elevation of the liver enzymes ALT and GGT and the development of hyperlipaemia and hyperlipiduria. A change of EE-TP dosing from 29 U/kg / 4 weeks to 14 U/kg/ 2 weeks failed to reduce the metabolites which increased to levels higher than those recorded pre-therapy. Studies have shown that liver disease can induce pathological alterations in the structure and function of the erythrocyte membrane and TPN administration can generate changes in the lipid composition and hence fluidity of the erythrocyte membrane [26–28]. Correct membrane function plays a pivotal role in modulating the activity of membrane channels and transporters and we therefore hypothesize that the lipid component of TPN interfered with the efficacy of EE-TP in Patient 1 at the level of the erythrocyte nucleoside transporter, ENT1. Hepatic steatosis and elevated transaminases in association with TPN administration has previously been reported in patients with MNGIE, and also disease-related hepatopathies, including hepatic steatosis, hepatomegaly, increased transaminases, abnormal liver function, triglyceride hyperlipidemia and cirrhosis [29,30]. The administration of TPN should therefore be carefully considered in such patients as the lipid component is metabolised by the mitochondrion, the site of the primary pathology in MNGIE. Thymidine phosphorylase is highly expressed in the liver of healthy individuals; for patients with MNGIE who express residual enzyme activity, the administration of TPN should be cautiously appraised.

Clinical assessments of Patient 2 between 27 and 76 months showed further improvements since the previous reporting period (0 to 27 months). Improvements in sensory ataxia, balance and gait and fine finger movements were recorded and patient reported outcomes included improvements in the distal sensation in hands and feet. Plasma creatine kinase, a marker of muscle damage, continued to decrease and remain within the normal range, indicating that EE-TP was able to relieve the skeletal muscle from the toxic effects of the accumulated thymidine and deoxyuridine. These results, and the clinical improvements noted in patients who received AHSCT, indicate that systemic reductions in thymidine and deoxyuridine concentrations are able to reverse the somatic mtDNA mutations that accumulate in post-mitotic cells [17]. An early normalization of nucleosides is essential to avoid the cumulative effects of metabolite imbalances on mtDNA and to prevent as much mitochondrial damage as possible. We, therefore, propose an expeditious treatment with EE-TP. This therapeutic approach would also increase patients eligibility for permanent treatments, for example with AHSCT or OLT, once a match is identified, or for future promising gene therapies [31,32]. Pivotal to this is an increased clinical awareness of MNGIE to enable an early diagnosis before disease progresses to an irreversible state.

Leukoencephalopathy is a hallmark of MNGIE and in the majority of affected individuals, it is initially patchy but progressively becomes more diffuse, appearing as hypointense on T1- and hyperintense on T2- weighted images and in FLAIR and fast spin echo (FSE) T2 sequences (1). A follow-up brain MRI scan of Patient 2 showed a progression of T2/FLAIR hyperintense cerebral white matter changes. EE-TP therapy was therefore not able to halt the progression of the leukoencephalopathy. However, it is not known whether therapy was able slow down the rate of progression. An additional consideration is the extent and distribution of leukoencephalopathy are not distinctly correlated with age and the clinical phenotype [33]. More extensive MRI studies of untreated and treated patients with MNGIE are required to establish whether the leukoencephalopathy can be reversed. *Post*

*mortem* histopathological studies of brains have shown the presence of albumin in the cytoplasm of reactive astrocytes in patients with MNGIE, compared to age-matched healthy individuals, suggesting functional blood brain barrier alterations and consequent vasogenic oedema as a cause of leukoencephalopathy in MNGIE [34]. Furthermore, a mild perivascular gliosis was also observed in immunohistochemical analyses [35]. Thymidine phosphorylase is also referred to as gliostatin, and has been shown to exert strong inhibitory effects on glial cells and neurotrophic effects on cortical neurons [36–38]. The leukoencephalopathy in MNGIE may therefore be a consequence of the absence of the inhibitory activity of thymidine phosphorylase, rather than its catalytic activity. If this is the case, the white matter aspect of MNGIE will not be responsive to treatment strategies which reduce or remove the pathological deoxyribonucleosides.

Adverse events for EE-TP observed in this compassionate study were nausea, coughing spasms and erythema of the face and neck in two of the three patients. These events were not classified as serious and were transient in nature, occurring within the first five minutes of EE-TP infusion. The events did not lead to patient discontinuation of EE-TP therapy, and were managed by subsequent pre-medication with antihistamine, corticosteroid anti-inflammatory and anti-emetic drugs. The introduction of a highly purified enzyme preparation allowed a careful and gradual withdrawal of this premedication.

Recombinant biologicals are increasingly used as a treatment modality for diseases due to the deficiency of an enzyme or protein, and the development of antibodies against administered recombinant proteins is well documented [39]. Recombinant thymidine phosphorylase is encapsulated within autologous erythrocytes and is less likely to elicit the strong antibody responses and neutralising antibodies that have been reported in the literature for other recombinant proteins. Antibodies specific to thymidine phosphorylase were detected in one patient, Patient 2 from treatment cycle 10 onwards. This observation does not raise any specific concerns with regard to neutralising antibodies as the efficacy of encapsulated thymidine phosphorylase in

metabolising the plasma metabolites improved over the 5.5 years of administration and positive clinical responses to treatment were recorded. The development of specific antibodies is not surprising considering that senescent erythrocytes are naturally sequestered from the vascular compartment by macrophages of the monocyte-macrophage system, where macrophages are able to present peptides to T lymphocytes. Pre-clinical studies have demonstrated that the administration of erythrocyte encapsulated antigens to BALB/c mice is able to elicit humoral responses [40].

## Conclusions

In conclusion, we have demonstrated the safety, tolerability and efficacy of EE-TP in three patients with MNGIE under a compassionate use programme. Regulatory and ethical approval for a Phase 2 clinical trial of EE-TP has been granted to further investigate this treatment approach in patients with MNGIE.

## Patents

Bax, B.E and Bain, M.D. (2018) *Treatment for mitochondrial neurogastrointestinal encephalomyopathy (MNGIE)*. EP2760459B1.

## References

1. Pacitti D, Levene M, Garone C, Nirmalanathan N, Bax BE. Mitochondrial Neurogastrointestinal Encephalomyopathy: Into the Fourth Decade, What We Have Learned So Far. *Front. Genet.* 2018; 9: 1–22.
2. Nishino I, Spinazzola A, Papadimitriou A, Hammans S, Steiner I, et al. Mitochondrial Neurogastrointestinal Encephalomyopathy: An Autosomal Recessive Disorder due to Thymidine Phosphorylase Mutations. *Ann. Neurol.* 2000; 47: 792–800.
3. Nishino I, Spinazzola A, Hirano M. Thymidine Phosphorylase Gene Mutations in MNGIE, a Human Mitochondrial Disorder. *Science* (80-). 1999; 283: 689–692.

4. Okamura K, Santa T, Nagae K, Omae T. Congenital Oculoskeletal Myopathy with Abnormal Muscle and Liver Mitochondria. *J. Neurol. Sci.* 1976; 27: 79–91.
5. Hirano M, Silvestri G, Blake DM, Lombes A, Minetti C, et al. Mitochondrial Neurogastrointestinal Encephalomyopathy (MNGIE): Clinical, Biochemical, and Genetic Features of an Autosomal Recessive Mitochondrial Disorder. *Neurology.* 1994; 44: 721–727.
6. Spinazzola A, Marti R, Nishino I, Andreu AL, Naini A, et al. Altered Thymidine Metabolism due to Defects of Thymidine Phosphorylase. *J. Biol. Chem.* 2002; 277: 4128–4133.
7. Martí R, Nishigaki Y, Hirano M. Elevated Plasma Deoxyuridine in Patients with Thymidine Phosphorylase Deficiency. *Biochem. Biophys. Res. Commun.* 2003; 303: 14–18.
8. Nishigaki Y, Martí R, Copeland WC, Hirano M. Site-Specific Somatic Mitochondrial DNA Point Mutations in Patients with Thymidine Phosphorylase Deficiency. *J. Clin. Invest.* 2003; 111: 1913–1921.
9. Nishigaki Y, Martí R, Hirano M. ND5 Is a Hot-Spot for Multiple Atypical Mitochondrial DNA Deletions in Mitochondrial Neurogastrointestinal Encephalomyopathy. *Hum. Mol. Genet.* 2004; 13: 91–101.
10. Hirano M, Nishigaki Y, Martí R. Mitochondrial Neurogastrointestinal Encephalomyopathy (MNGIE): A Disease of Two Genomes. *Neurologist.* 2004; 10: 8–17.
11. Yavuz H, Özel A, Christensen M, Christensen E, Schwartz M, et al. Treatment of Mitochondrial Neurogastrointestinal Encephalomyopathy with Dialysis. *Arch. Neurol.* 2007; 64: 435–438.
12. La Marca G, Malvagia S, Casetta B, Pasquini E, Pela I, et al. Pre- and Post-Dialysis Quantitative Dosage of Thymidine in Urine and Plasma of a MNGIE Patient by Using HPLC-ESI-MS/MS. *J. Mass Spectrom.* 2006; 41: 586–592.
13. Lara MC, Weiss B, Illa I, Madoz P, Massuet L, et al. Infusion of Platelets Transiently Reduces Nucleoside Overload in MNGIE. *Neurology.* 2006; 67: 1461–1463.
14. Halter J, Schüpbach WMM, Casali C, Elhasid R, Fay K, et al. Allogeneic Hematopoietic SCT as Treatment Option for Patients with Mitochondrial Neurogastrointestinal

- Encephalomyopathy (MNGIE): A Consensus Conference Proposal for a Standardized Approach. *Bone Marrow Transplant*. 2011; 46: 330–337.
15. Filosto M, Scarpelli M, Tonin P, Lucchini G, Pavan F, et al. Course and Management of Allogeneic Stem Cell Transplantation in Patients with Mitochondrial Neurogastrointestinal Encephalomyopathy. *J. Neurol*. 2012; 259: 2699–2706.
  16. Hirano M, Martí R, Casali C, Tadesse S, Uldrick T, et al. Allogeneic Stem Cell Transplantation Corrects Biochemical Derangements in MNGIE. *Neurology*. 2006; 67: 1458–1460.
  17. Halter JP, Michael W, Schüpbach M, Mandel H, Casali C, et al. Allogeneic Haematopoietic Stem Cell Transplantation for Mitochondrial Neurogastrointestinal Encephalomyopathy. *Brain*. 2015; 138: 2847–2858.
  18. Moran NF, Bain MD, Muqit MMK, Bax BE. Carrier Erythrocyte Entrapped Thymidine Phosphorylase Therapy for Mngie. *Neurology*. 2008; 71: 686–688.
  19. Bax BE, Bain MD, Scarpelli M, Filosto M, Tonin P, et al. Clinical and Biochemical Improvements in a Patient with MNGIE Following Enzyme Replacement. *Neurology*. 2013; 81: 1269–1271.
  20. De Giorgio R, Pironi L, Rinaldi R, Boschetti E, Caporali L, et al. Liver Transplantation for Mitochondrial Neurogastrointestinal Encephalomyopathy. *Ann. Neurol*. 2016; 80: 448–455.
  21. Godfrin Y, Horand F, Franco R, Dufour E, Kosenko E, et al. International Seminar on the Red Blood Cells as Vehicles for Drugs. *Expert Opin. Biol. Ther*. 2012; 12: 127–133.
  22. Bax BE, Bain MD, Talbot PJ, Parker-williams EJ, Chalmers RA. Survival of Human Carrier Erythrocytes in Vivo. *Clin. Sci*. 1999; 96: 171–178.
  23. Fairbanks LD, Levene M, Bax BE. Validation of a HPLC Method for the Measurement of Erythrocyte Encapsulated Thymidine Phosphorylase (EE-TP) Activity. *J. Pharm. Biomed. Anal*. 2013; 76: 8–12.
  24. Levene M, Pacitti D, Gasson C, Hall J, Sellos-Moura M, et al. Validation of an Immunoassay for Anti-Thymidine Phosphorylase Antibodies in Patients with MNGIE Treated

- with Enzyme Replacement Therapy. *Mol. Ther. - Methods Clin. Dev.* 2018; 11: 1–8.
25. Barton GJ, Ponting CP, Spraggon G, Finnis C, Sleep D. Human Platelet-derived Endothelial Cell Growth Factor Is Homologous to Escherichia Coli Thymidine Phosphorylase. *Protein Sci.* 1992; 1: 688–690.
  26. Maksane S, Guhasarkar S, Banerjee R, Dandekar S. Evaluation of Surface Properties of Erythrocyte Membranes in Liver Diseases. *Int. J. Res. Med. Sci.* 2015; 3: 593.
  27. Dahlan W, Richelle M, Kulapongse S. Modification of Erythrocyte Membrane Lipid Composition Induced by a Single Intravenous Infusion of. 1958; 6.
  28. Innis SM. Alteration of Erythrocyte Lipid Composition Following Total Parenteral Nutrition in the Rat. *J. Parenter. Enter. Nutr.* 1989; 13: 47–50.
  29. Garone C, Tadesse S, Hirano M. Clinical and Genetic Spectrum of Mitochondrial Neurogastrointestinal Encephalomyopathy. *Brain.* 2011; 134: 3326–3332.
  30. Wang J, Chen W, Wang F, Wu D, Qian J, et al. Nutrition Therapy for Mitochondrial Neurogastrointestinal Encephalopathy with Homozygous Mutation of the TYMP Gene. *Clin. Nutr. Res.* 2015; 4: 132.
  31. Yadak R, Cabrera-Pérez R, Torres-Torronteras J, Bugiani M, Haeck JC, et al. Preclinical Efficacy and Safety Evaluation of Hematopoietic Stem Cell Gene Therapy in a Mouse Model of MNGIE. *Mol. Ther. - Methods Clin. Dev.* 2018; 8: 152–165.
  32. Torres-Torronteras J, Cabrera-Pérez R, Barba I, Costa C, de Luna N, et al. Long-Term Restoration of Thymidine Phosphorylase Function and Nucleoside Homeostasis Using Hematopoietic Gene Therapy in a Murine Model of Mitochondrial Neurogastrointestinal Encephalomyopathy. *Hum. Gene Ther.* 2016; 27: 656–667.
  33. Scarpelli M, Ricciardi GK, Beltramello A, Zocca I, Calabria F, et al. The Role of Brain MRI in Mitochondrial Neurogastrointestinal Encephalomyopathy. *Neuroradiol. J.* 2013; 26: 520–530.
  34. Szigeti K, Sule N, Adesina AM, Armstrong DL, Saifi GM, et al. Increased Blood-Brain Barrier Permeability with

- Thymidine Phosphorylase Deficiency. *Ann. Neurol.* 2004; 56: 881–886.
35. Gramegna LL, Pisano A, Testa C, Manners DN, D'Angelo R, et al. Cerebral Mitochondrial Microangiopathy Leads to Leukoencephalopathy in Mitochondrial Neurogastrointestinal Encephalopathy. *Am. J. Neuroradiol.* 2018; 39: 427–434.
  36. Asai K, Hirano T, Kaneko S, Moriyama A, Nakanishi K, et al. A Novel Glial Growth Inhibitory Factor, Gliostatin, Derived from Neurofibroma. *J. Neurochem.* 1992; 59: 307–317.
  37. Asai K, Nakanishi K, Isobe I, Eksioglu YZ, Hirano A, et al. Neurotrophic Action of Gliostatin on Cortical Neurons. Identity of Gliostatin and Platelet-Derived Endothelial Cell Growth Factor. *J. Biol. Chem.* 1992; 267: 20311–20316.
  38. Ueki T, Nakanishi K, Asai K, Okouchi Y, Isobe I, et al. Neurotrophic Action of Gliostatin on Cocultured Neurons with Glial Cells. *Brain Res.* 1993; 622: 299–302.
  39. Faraji F, Karjoo Z, Moghaddam MV, Heidari S, Enameh RZ, et al. Challenges Related to the Immunogenicity of Parenteral Recombinant Proteins: Underlying Mechanisms and New Approaches to Overcome It. *Int. Rev. Immunol.* 2018; 1–15.
  40. Murray AM, Pearson IFS, Fairbanks LD, Chalmers RA, Bain MD, et al. The Mouse Immune Response to Carrier Erythrocyte Entrapped Antigens. *Vaccine.* 2006; 24: 6129–6139.