Book Chapter

The Role of Iron in the P-Acquisition Mechanisms of the Unicellular N₂-Fixing Cyanobacteria *Halothece* sp., Found in Association with the Mediterranean Seagrass *Posidonia oceanica*

Víctor Fernández-Juárez^{1*}, Antoni Bennasar-Figueras², Antonio Tovar-Sanchez³ and Nona Sheila R Agawin¹

¹Marine Ecology and Systematics (MarEs), Department of Biology, Universitat de les Illes Balears (UIB), Spain

²Grup de Recerca en Microbiologia, Departament de Biologia, Universitat de les Illes Balears (UIB), Spain

³Department of Ecology and Coastal Management, Andalusian Institute for Marine Science, ICMAN (CSIC), Spain

***Corresponding Author:** Víctor Fernández-Juárez, Marine Ecology and Systematics (MarEs), Department of Biology, Universitat de les Illes Balears (UIB), Palma de Mallorca, Spain

Published **March 27, 2020**

This Book Chapter is a republication of an article published by Víctor Fernández-Juárez, et al. at Frontiers in Microbiology in August 2019. (Fernández-Juárez V, Bennasar-Figueras A, Tovar-Sanchez A and Agawin NSR (2019) The Role of Iron in the P-Acquisition Mechanisms of the Unicellular N₂-Fixing Cyanobacteria *Halothece* sp., Found in Association With the Mediterranean Seagrass *Posidonia oceanica*. Front. Microbiol. 10:1903. doi: 10.3389/fmicb.2019.01903)

How to cite this book chapter: Víctor Fernández-Juárez, Antoni Bennasar-Figueras, Antonio Tovar-Sanchez, Nona Sheila R Agawin. The Role of Iron in the P-Acquisition Mechanisms of the Unicellular N₂-Fixing Cyanobacteria *Halothece* sp., Found in Association with the Mediterranean Seagrass *Posidonia*

oceanica. In: Alexandre Morrot, editor. Prime Archives in Microbiology. Hyderabad, India: Vide Leaf. 2020.

© The Author(s) 2020. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions: VFJ and NSRA designed the experiments. VFJ conducted all experiments and led the writing of the paper. All authors contributed to the writing and review of the manuscript, and NSRA is the supervisor of the laboratory.

Funding: This work was supported by funding to NSRA through the Agencia Estatal de Investigación and the European Regional Development Funds project (CTM2016-75457-P).

Acknowledgments: We acknowledge the support and help of Scientific Technical Service (Dr. Guillem Ramis Munar and Maria Trinidad Garcia Barceló) of the University of the Balearic Islands for cytometry and gas-chromatography analyses, respectively. We also thank Pere Ferriol Buñola and Alba Coma Ninot for the help in the acquisition of the cultures.

Abstract

Posidonia oceanica, an endemic seagrass of the Mediterranean Sea harbours a high diversity of N₂-fixing prokaryotes. One of these is *Halothece* sp., a unicellular N₂-fixing cyanobacteria detected through *nifH* analysis from the epiphytes of *P. oceanica*. The most related strain in culture is *Halothece* sp. PCC 7418 and this was used as the test organism in this study. In the Mediterranean Sea, phosphorus (P) and iron (Fe) can be the major limiting nutrients for N₂ fixation. However, information

about the mechanisms of P-acquisition and the role of metals (*i.e.* Fe) in these processes for N₂-fixing bacteria is scarce. From our genomic analyses of the test organism and other phylogenetically related N₂-fixing strains, *Halothece* sp. PCC 7418 is one of the strains with the greatest number of gene copies (eight copies) of Alkaline Phosphatases (APases). Our structural analysis of PhoD (alkaline phosphatase type D) and PhoU (phosphate acquisition regulator) of *Halothece* sp. PCC 7418 showed the connection among metals (Ca²⁺ and Fe³⁺), and the P-acquisition mechanisms. Here, we measured the rates of Alkaline Phosphatase Activity (APA) through MUF-P hydrolysis under different combinations of concentrations of inorganic P (PO₄³⁻) and Fe in experiments under N₂-fixing (low NO₃⁻ availability) and non-N₂ fixing (high NO₃⁻ availability) conditions. Our results showed that APA rates were enhanced by the increase in Fe availability under low levels of PO₄³⁻, especially under N₂-fixing conditions. Moreover, the increased PO₄³⁻-uptake was reflected in the increased of the P-cellular content of the cells under N₂ fixation conditions. We also found a positive significant relationship between cellular P and cellular Fe content of the cells ($r^2 = 0.71$, $p < 0.05$). Our results also indicated that Fe-uptake in *Halothece* sp. PCC 7418 was P and Fe-dependent. This study gives first insights of P-acquisition mechanisms in the N₂-fixing cyanobacteria (*Halothece* sp.) found in *P. oceanica* and highlights the role of Fe in these processes.

Keywords

Halothece sp. PCC 7418; *Posidonia Oceanica*; Alkaline Phosphatase (APase); N₂ Fixation; PO₄³⁻-Fe Uptake; Iron; PhoD and PhoU

Introduction

Posidonia oceanica is an endemic seagrass in the Mediterranean Sea, forming extensive meadows with valuable established key ecological services: high primary productivity, as a carbon sink, as a habitat and nursery for a variety of micro- and macro-

organisms, as sediment stabilizers, as buffers for ocean acidification, and as an important site for biogeochemical processes (*e.g.* nitrogen cycles) [1-3]. Atmospheric nitrogen (N_2) fixation associated with *P. oceanica* meadows are similar in rates or even higher than tropical seagrasses and may play a key role in maintaining the high productivity of the *P. oceanica* in oligotrophic waters [1,4]. N_2 fixation in *P. oceanica* is carried out by microorganisms called diazotrophs that can be found on the surface of the leaves, roots and rhizomes (epiphytic population) or even inside of the roots (endophytic population) [5,6]. Among the diazotrophic prokaryotes, a huge variety of diazotrophic cyanobacteria have been detected based on the sequence analysis of *nifH* gene (gene coding for the nitrogenase enzyme responsible for the N_2 fixation) on the leaves of *P. oceanica* [1,4].

In general, cyanobacteria are key components in the marine food web, contributing significantly to primary production in oligotrophic oceans [7,8]. Compared with other phytoplankton taxa, cyanobacteria have elevated ratio of nitrogen (N):phosphorus (P) (a molar ratio above 25 compared with the general Redfield ratio of 16 in marine phytoplankton) and can be a consequence of having two light-harvesting complexes [9-11]. Changes affecting the N:P ratios in their environment by limiting concentration of N or P, could change their N:P tissue composition and may have consequences in their adaptation and survival and possibly the N_2 fixation activities of diazotrophic cyanobacteria [6,12]. Nonetheless, these versatile microorganisms may have several adaptive mechanisms to changes in their dynamic marine environment (*e.g.* nutrient availability) [8,13,14].

Phosphorus, (*i.e.* inorganic phosphorus, PO_4^{3-}), together with iron (Fe) are hypothesized to be the major limiting nutrients for N_2 fixation [15-17]. Phosphorus is vital for the storage and retrieval system of genetic information (DNA/RNA), for the energy metabolism through ATP dependence [18-20] and in most bacteria, it is important for the structure of the cell membrane. During P-starvation, microorganisms produce enzymes that are capable to hydrolyse P-esters contained in

dissolved organic phosphorus (DOP) releasing dissolved inorganic phosphorus (DIP), that the cells can utilise. These enzymes are called Alkaline Phosphatases (APases) and in marine bacteria they are included in three main families: PhoA, PhoX and PhoD. APases are metalloenzymes that require metal co-factors. PhoA forms a coordinate with two zinc (Zn^{2+}) and one magnesium (Mg^{2+}) ions; PhoX forms a coordinate with three calcium (Ca^{2+}) and one/two Fe^{3+} ions [21]; and PhoD coordinates with an unknown number of Ca^{2+} ions. In *Bacillus subtilis* model, PhoD has an active site formed with one Fe^{3+} and two Ca^{2+} ions [22]. This information suggests the possible interaction between metals (*e.g.* Fe^{3+} , Ca^{2+} , Mg^{2+} , Zn^{2+}) in the mechanisms of P-acquisition involving APases. In *Halothece* sp. PCC 7418, two types of APases have been reported: PhoA (two copies) and PhoD (one copy). Calcium dependence was proven in PhoD in *Halothece* sp. PCC 7418 (Kageyama et al., 2011). However, Fe dependence of PhoD and the relative importance between these two types of APases (PhoA and PhoD) have not been demonstrated in *Halothece* sp. PCC 7418.

APases are included in what is known as the Pho regulon. It is a huge regulatory group of genes that control P-acquisition. Pho regulon is composed of elements related with (1) high-affinity phosphate transport (PstS, PstC, PstA and PstB) and low-affinity phosphate transport, (2) extracellular enzymes capable of obtaining PO_4^{3-} from organic phosphates (APases), and (3) polyphosphate metabolism (PpK, PpX and PpA) as P reservoir or elements with unknown functions (PhoU) [19,23,24]. PhoU coordinates with metal cluster (Zn^{2+} or Fe^{3+}), and may have a role in the control of autokinase activity of the PhoR and Pst systems [25]. The Pho regulon is mainly controlled by PhoR-PhoB, a two-component regulatory system [19]. PhoR is an inner-membrane histidine kinase, while PhoB is a transcriptional factor that recognises and binds to consensus sequence named PHO box. In cyanobacteria, PHO box is formed by three tandem repeats of 8 bp separated by 5 bp, unlike PHO Box from *Escherichia coli*, formed by two direct repeats of 7 bp separated by 5 bp [20,24,26].

The P-acquisition mechanisms in bacteria are well studied in the Atlantic ocean, where Fe is shown to enhance the P-acquisition mechanisms in N₂-fixing cyanobacterial species, *Trichodesmium* spp. and *Crocospaera watsonii* [27-29]. However, there is scarcely any information about the relation between metals (e.g. Fe) and P-acquisition mechanisms in N₂-fixing cyanobacteria found in association with the Mediterranean seagrass, *P. oceanica*, taking into account the multiple ecological benefits of this seagrass in the region. The Mediterranean Sea is oligotrophic, characterized by low water column PO₄³⁻ concentrations and a decreasing gradient of PO₄³⁻ concentrations from west to east basins [30]. Knowledge on the P-acquisition mechanisms of N₂-fixing organisms in an environment with limiting levels of PO₄³⁻ is particularly important. Moreover, the Mediterranean Sea is subject to Saharan atmospheric dust deposition containing Fe [31] which can play a role in the P-acquisition mechanisms of the organisms.

To study, for the first time, the P-acquisition mechanisms in N₂-fixing cyanobacteria associated with the dominant coastal ecosystem in the region (*P. oceanica* seagrass beds), we selected a diazotrophic unicellular cyanobacteria, *Halothece* sp. found on the leaves of *P. oceanica* [4] as our test species. The most related culturable strain is *Halothece* sp. PCC 7418 and this was used as the test organism in this study. The halotolerant *Halothece* sp. PCC 7418 (originally called *Synechococcus* PCC 7418), also known as *Aphanothece halophytica*, was originally isolated from Solar Lake on the eastern shore of the Sinai Peninsula in 1972 (UniProt source). First, we made a genomic analyses of the Pho regulon to check the regulatory group of genes that control the P-acquisition mechanisms and then a structural analysis of PhoD (alkaline phosphatase type D) and PhoU (phosphate acquisition regulator) of *Halothece* sp. PCC 7418 to investigate the connection among metals (e.g. Ca²⁺ and Fe³⁺) and the P-acquisition mechanisms of this species. Second, we investigated experimentally how the availability of Fe affects the Alkaline Phosphatase Activity (APA) and their PO₄³⁻-uptake rates and the magnitude of the effect under different levels of PO₄³⁻ and NO₃⁻ availability, and how the availability of PO₄³⁻ and Fe affect Fe-uptake rates of the cells.

Materials and Methods

Genome Analysis

With the goal of comparing *Halotheca* sp. PCC 7418 Pho regulon with its closest genomes [32], the distribution of the number of copies of Pho regulon components in selected strains was analysed. The genome from *Halotheca* sp. PCC 7418 (GenkBank: NC_019779.1) and genomes from other closely related microorganisms were compared using the dedicated bacterial information system Pathosystems Resource Integration Center (PATRIC). This database, and the analysis tools included, offers an easy interface in which annotated genes that are included in different subsystems can be searched [33].

Three-Dimensional Predicted Structures

Sequences of PhoD and PhoU in FASTA format were sent to the I-Tasser server for protein 3D-structure prediction [34], with their domains previously checked in Pfam 32.0 [35]. The predicted structures for PhoD and PhoU of *Halotheca* sp. PCC 7418 were sent to POSA [36] for a structural alignment against PhoD of *B. subtilis* (PDB: 2YEQ) and PhoU of *Pseudomonas aeruginosa* (PDB: 4Q25) respectively; the two more evolutionarily related homologous proteins available to date in databases (*i.e.*, that have both similar sequences and 3D models). To describe the Fe coordination positions of these proteins, residues from both (*i.e.*, PhoD and PhoU of *Halotheca* sp. PCC 7418 against 2YEQ and 4Q25, respectively) were mapped through alignment with Uniprot Clustal Omega [37]. The predicted structures and the corresponding structural alignments were visualised with Pymol [38].

Strain and Culture Conditions

Halotheca sp. PCC 7418, was obtained from the Pasteur Culture Collection of Cyanobacteria (PCC) and maintained in 250 ml acid-cleaned Quartz Erlenmeyer flasks containing 150 ml of ASNIII + Tu4X medium (initial pH 7.5) [39]. The medium was supplemented with 0.1-0.3 % (w/v) of glucose and grown in a rotary shaker (120 r.p.m) with a photoperiod of 12 h light: 12 h

dark under low intensity fluorescent light ($30 \mu\text{E m}^{-2}\text{s}^{-1}$) at 25°C . Three conditions were established for inorganic phosphorus (PO_4^{3-}) concentrations: [Low PO_4^{3-}] ($0.1 \mu\text{M}$), [Medium PO_4^{3-}] ($1 \mu\text{M}$) and [High PO_4^{3-}] ($45 \mu\text{M}$). Furthermore, three conditions for Fe were established: [Low Fe] (2 nM), [Medium Fe] (20 nM) and [High Fe] ($7.5 \mu\text{M}$). These PO_4^{3-} and Fe concentration were combined in nine conditions ([Low PO_4^{3-} - Low Fe], [Low PO_4^{3-} - Medium Fe], [Low PO_4^{3-} - High Fe], [Medium PO_4^{3-} - Low Fe], [Medium PO_4^{3-} - Medium Fe], [Medium PO_4^{3-} - High Fe], [High PO_4^{3-} - Low Fe], [High PO_4^{3-} - Medium Fe] and [High PO_4^{3-} - High Fe]), and these treatments were tested in two sets of experiments: growth under 4.4 mM of NO_3^- (optimal concentration) and 0.15 mM of NO_3^- (low concentration, and referred from now on as [Low NO_3^-]) (Table 1). The solutions of PO_4^{3-} , Fe and NO_3^- were prepared from K_2HPO_4 , ferric citrate and NaNO_3 , respectively. The batch cultures were maintained for over 10 days for each experiment and the initial inoculum of cells was added at exponential phase ($\text{O.D}_{750 \text{ nm}} \cong 0.2$) from their original ASNIII + Tu4X medium.

Selected treatments ([Low PO_4^{3-} - Low Fe], [Low PO_4^{3-} - High Fe], [High PO_4^{3-} - Low Fe] and [High PO_4^{3-} - High Fe]) were also used to compare the results under NO_3^- starvation (6.66 nM) and optimal NO_3^- conditions. Cultures were maintained at the same conditions as described above for over 12 days. During the last day, PO_4^{3-} , Fe and/or NO_3^- were added to the different treatments to achieve optimal concentrations of PO_4^{3-} ($45 \mu\text{M}$), Fe ($7.5 \mu\text{M}$) and NO_3^- (4.4 mM) to evaluate the changes in the APA rates, and the new conditions were maintained for over four days. The different conditions of the experiments are shown in Table 1.

Table 1: List of all experimental treatments conducted in this study. In the recovery experiments, PO_4^{3-} , Fe and/or NO_3^- were added to the different initial treatments to achieve optimal conditions to evaluate the changes in APA rates.

Experiments			
Condition (optimal and Low NO_3^-)			Description
[Low PO_4^{3-} - Low Fe]	[Medium PO_4^{3-} - Low Fe]	[High PO_4^{3-} - - Low Fe]	1 st experiment-optimal NO_3^- (4.4 mM)
[Low PO_4^{3-} - Medium Fe]	[Medium PO_4^{3-} - Medium Fe]	[High PO_4^{3-} - - Medium Fe]	2 nd experiment- low NO_3^- (0.15 mM)
[Low PO_4^{3-} - High Fe]	[Medium PO_4^{3-} - High Fe]	[High PO_4^{3-} - - High Fe]	3 rd experiment- NO_3^- starvation (6 nM), comparing with optimal NO_3^- in selected treatments
Recovery experiments			
Initial Treatment	Condition of NO_3^-	Nutrient added (at day 12)	Resulting treatment (maintained for 4 days)
[Low PO_4^{3-} - Low Fe]	optimal NO_3^- (4.4 mM)	PO_4^{3-} and Fe	[High PO_4^{3-} - High Fe] in optimal NO_3^- treatment.
[Low PO_4^{3-} - Low Fe]	NO_3^- starvation (6.66 nM)	PO_4^{3-} , Fe and NO_3^-	[High PO_4^{3-} - High Fe] in optimal NO_3^- treatment.

The importance of PhoD in *Halothece* sp. PCC 7418 was investigated by changing the availability of the metal co-factors for PhoA (Zn^{2+} and Mg^{2+}). The method used was as described above in the initial main experiments except that the medium was depleted with Mg and Zn and the condition of PO_4^{3-} and Fe was: [Medium PO_4^{3-} - High Fe] under optimal NO_3^- .

All cultures were performed in duplicate, and the study parameters (APA, N_2 fixation, uptake rates of PO_4^{3-} and Fe, TDP and/or P/Fe/Mn cellular content) were evaluated during the different phases of the culture ($\text{O.D}_{750 \text{ nm}} \cong 0.01\text{-}0.2$). A subsample of the cells (1.5 ml) was taken from the culture flasks during the experiment and were counted through flow cytometric analysis (as described below) to normalize the results

per cell. All samples were manipulated in a class-100 clean hood, to avoid Fe contamination.

Flow Cytometry Analysis

Cells were fixed with glutaraldehyde 25% (v/v) in H₂O (Sigma-Aldrich) [final concentration 0.05 % (v/v)] and were counted with a Becton Dickinson FACS-Verse cytometer (Beckton & Dickinson, Franklin Lakes, New Jersey, USA). Fluorescent beads, BD FACSuite™ CS&T research beads (Beckton & Dickinson and Company BD Biosciences, San Jose, USA), were used as internal standard to calibrate the instrument. The cytometer shows fluorescence patterns for FITC, PE, PerCP-CyTM5.5 and APC. To count the *Halothece* sp. PCC 7418 cells, we selected FITC (488 nm excitation, 530/30 nm emission) and PE (488 nm excitation, 576/26 nm emission) combination fluorescence signals which show clearly the population of the cells. A total of 10.000 cells were counted in each sample and the counted cells were expressed as cells μl^{-1} .

Alkaline Phosphatase Activity (APA)

APA was evaluated through a fluorometric assay, in which the hydrolysis of the fluorogenic substrate (S) 4-methylumbelliferyl phosphate (MUF-P, Sigma-Aldrich) to 4-methylumbelliferyl (MUF) was measured. Generally, an end point enzymatic assay was conducted with a concentration of 2 μM MUF-P during the exponential phase of the culture ($\text{O.D}_{750\text{ nm}} \cong 0.1$). After one hour incubation in darkness at room temperature, APA was measured in a microtiter plate that contained borate buffer at pH 10 (3:1 of sample:buffer). The MUF production (fmole MUF $\text{cell}^{-1} \text{ h}^{-1}$) was measured with a Cary Eclipse spectrofluorometer (FL0902M009, Agilent Technologies) at 359 nm (excitation) and 449 nm (emission) and using a calibration standard curve with commercial MUF (Sigma-Aldrich).

Saturation curves of velocity (V , fmole MUF $\text{cell}^{-1} \text{ h}^{-1}$) vs substrate (S , μM) were made under [Low NO_3^-] condition during the final exponential phase of the culture ($\text{O.D}_{750\text{ nm}} \cong 0.2$), using different concentrations of MUF-P: 0 μM , 0.05 μM , 0.1 μM , 0.5

μM , $2 \mu\text{M}$ and $5 \mu\text{M}$. The maximum velocity (V_{max}) at saturating substrate concentrations was obtained from each plot of V vs S . The Michaelis-Menten constant, K_m (μM) which represents the substrate concentration at half V_{max} was calculated using de Hill plot equation [40]. The evolution of MUF-P hydrolysis rates ($\text{fmole MUF cell}^{-1}$) with time (h) was recorded over one hour in the treatments under [Low PO_4^{3-} -Low Fe], [Low PO_4^{3-} -High Fe] and [High PO_4^{3-} -High Fe] at the last day of the experiment with $5 \mu\text{M}$ of MUF-P under NO_3^- starvation and NO_3^- optimal conditions and the APA rate ($\text{fmole MUF cell}^{-1} \text{ h}^{-1}$) was calculated as the slope of the fitted line.

PO_4^{3-} uptake Rates, Nutrient Concentrations in the Culture Medium and in the Cells

Samples for the determination of PO_4^{3-} and total dissolved P (TDP) were centrifuged for 15 minutes at $16,000 \times g$ under 4°C . The supernatant was collected from the centrifuged tubes and used for PO_4^{3-} determinations following standard spectrophotometric methods [41]. TDP concentrations were also analysed using the latter method after persulfate digestion. Samples for Fe analyses of culture media were filtered through sterile $0.2 \mu\text{m}$ filters (MFV5-025, FilterLab) at different times (initial and final) during the experiments. The metal (Fe) concentrations of culture medium were measured by inductively coupled plasma mass spectrometry (ICP-MS; iCap, Thermo Scientific), following the trace-metal clean techniques described in Tovar-Sanchez et al. [42] and Tovar-Sanchez & Sañudo-Wilhelmy [43].

The PO_4^{3-} concentrations in the culture medium were determined at different times: 0, 1, 4 and 10 days in the experimental treatment of [High PO_4^{3-}]: [Low Fe], [Medium Fe] and [High Fe]), and under [Low NO_3^-] and optimal NO_3^- conditions. Specific PO_4^{3-} uptake rates ($\text{pmole PO}_4^{3-} \text{ cell}^{-1} \text{ d}^{-1}$) were calculated as described in [44]. Briefly, specific PO_4^{3-} uptake rates were calculated as the mass balance of PO_4^{3-} over

the multiple days by taking the differences of PO_4^{3-} concentrations at two different times (T_0 - T_1 , T_0 - T_4 and T_0 - T_{10}) and normalized by the number of cells counted at different time points (0, 1, 4 and 10) through the following equation:

$$\text{PO}_4^{3-}\text{-uptake (pmole PO}_4^{3-} \text{ cell}^{-1} \text{ d}^{-1}) = \frac{C-D}{T_i-T_f} \quad \text{eq.} \quad (1)$$

C is $\mu\text{mole PO}_4^{3-} \text{ cell}^{-1}$ at the initial time (T_i) and D is the $\mu\text{mole PO}_4^{3-} \text{ cell}^{-1}$ at the final time (T_f).

TDP concentrations were also measured at different times: 0, 4, 8 and 12 days in the experiments under NO_3^- starvation at [Low PO_4^{3-}] and [High PO_4^{3-}] conditions. Fe-uptake rates were measured under N_2 -fixing conditions (*i.e.* [Low NO_3^-] conditions). Initial and final Fe concentrations of the culture media were measured, and the difference between time=0 and time=10 (T_0 - T_{10}) was used to determine the Fe-uptake during the 10 days of the experiment. Specific Fe-uptake ($\text{fmole Fe cell}^{-1} \text{ d}^{-1}$) was calculated the same way as the specific PO_4^{3-} -uptake rates described above.

Cellular contents of phosphorus (P), Fe and other metals (*i.e.* Mn, V, Co, Ni or Zn) were also determined by collecting the cells under [Low NO_3^-] treatment conditions through filtration of a known volume of culture (20 ml) with 0.2- μm acid-cleaned polycarbonate filters (Merck-Millipore). Elemental concentrations of P and Fe in the cyanobacterial samples were determined by inductively coupled plasma mass spectrometry (ICP-MS; iCap, Thermo Scientific), after microwave acid digestion (CEM, Mars 5) using nitric acid (high purity Suprapur®, Merck) [42,43].

Acetylene Reduction Assay (ARA)

N_2 -fixing activities were measured with the acetylene reduction assay (ARA) method under known N_2 -fixing conditions for unicellular cyanobacteria (*i.e.* low NO_3^- concentrations, anaerobic environment, dark phase of the photoperiod, Reddy et

al., [45]), and under low-medium levels of Fe and in low-medium-high levels of PO_4^{3-} . A volume of 50 ml from treatments with [Low NO_3^-] condition at day 8 of the experiment was transferred to anaerobic tubes for cultivation for 2 days, and after which, ARA measurements were done following the method described in Agawin et al. [46]. Duplicate 10 ml samples of culture from each experimental tube, were filtered through 0.45 μm GF/F filters (MFV5-025, FilterLab). The filters were deposited in hermetic vials containing 1 ml of the corresponding culture medium. Acetylene (C_2H_2) was added at 20 % (v/v) final concentration in each vial using gas-tight Hamilton syringes. The filters were incubated in the vials for three hours at room temperature in the dark. After the 3 h incubation time, 10 ml of headspace gas were removed with a gas-tight Hamilton syringe from the incubation vials or tubes, transferred and stored in Hungate tubes and sealed with hot melt adhesive glue (SALKI, ref. 0430308) to minimize gas losses [46]. Ethylene and acetylene were determined using a GC (model HP-5890, Agilent Technologies) equipped with a flame ionization detector (FID). The column was a Varian wide-bore column (ref. CP7584) packed with CP-PoraPLOT U (27.5 m length, 0.53 mm inside diameter, 0.70 mm outside diameter, 20 μm film thickness). Helium was used as carrier gas at a flow rate of 30 ml min^{-1} . Hydrogen and airflow rates were set at 30 ml min^{-1} and 365 ml min^{-1} , respectively. The split flow was used so that the carrier gas flow through the column was 4 ml min^{-1} at a pressure of 5 psi. Oven, injection and detector temperatures were set at 52 °C, 120 °C and 170 °C, respectively. Ethylene produced was calculated using the equations in Stal [47]. The acetylene reduction rates were converted to N_2 fixation rates ($\text{pmole N}_2 \text{ ml}^{-1} \text{ h}^{-1}$) using a factor of 4:1 [48].

Statistical Analyses

Univariate Analysis of variance (ANOVA) factor analyses and post-hoc (Bonferroni) was used to study the effect of the nutrient treatment conditions to APA rates, P-cellular content and specific PO_4^{3-} and Fe uptake rates. In other cases, where we

want to highlight a specific point, we use individual t-tests. Regression analyses were used to determine the relationships between P-cellular content vs N₂ rates fixation, P-cellular content vs Fe-cellular content and P/Fe-cellular content vs other metals (*i.e.* Mn). The statistical analyses were performed using the SPSS program version 21 (IBM Corp year 2012).

Results

Pho Regulon of *Halotheca* sp. PCC 7418

The distribution of the number of copies of Pho regulon components of *Halotheca* sp. PCC 7418 and its closest genomes [32] are shown in Figure 1 and Supplementary Table 1. The *Gloeocapsa* sp. PCC 7428 genome had the highest number of copies detected (up to 45), suggesting that this species is one of the better adapted species to P-limitation. On the other hand, *Nostoc punctiforme* PCC 73102 and *Chroococcidiopsis thermalis* PCC 7203 genomes had the lowest number of copies of the Pho regulon components. Our test microorganism *Halotheca* sp. PCC 7418 genome was the fourth cyanobacterium containing more copies of the Pho regulon (26): 1 for *phoU*, 4 for *pstS*, 3 for *pstC*, 2 for *pstA*, 3 for *pstB*, 1 for *phoR-phoB*, 8 for APases, 1 for *ppK*, 1 for *ppX* and 1 for *ppA*. With 8 copies of APases, it was the second cyanobacterium containing more APases (8), only surpassed by *Gloeocapsa* sp. PCC 7428 (19), suggesting a key role of the APases in *Halotheca* sp. PCC 7418. Annotation in PATRIC did not annotate any specific APase, except for a PhoD. No low-affinity phosphate transporters were detected.

Figure 1: Distribution of the Pho regulon in *Halothece* sp. PCC 7418 (*) and genomes of its closest cyanobacterial relatives. Annotated genes were identified in PATRIC.

Structural Analysis of PhoD and PhoU of *Halothece* sp. PCC 7418

Three-dimensional structure of PhoD and its implication in APA

The annotated PhoD of *Halothece* sp. PCC 7418 displayed 511 amino acids (aa) with two domains *i.e.*; PhoD-like phosphatase N-terminal domain and PhoD-like phosphatase domain. Predicted structure of PhoD (C-score = 0.00, estimated TM-score = 0.71 ± 0.11 , estimated RMSD = $7.4 \pm 4.3\text{\AA}$) had 10 α -helix and 21 β -chains. PhoD of *Halothece* sp. PCC 7418 was homologue to the crystal structure of PhoD of *B. subtilis* (2YEQ) of 522 aa, with an identity of 47.5% and coverage of 91.6%. Figure 2A shows the structural alignment between PhoD of *Halothece* sp. PCC 7418 and 2YEQ. The sequence alignment displayed up to 40.11% similarity and was used with the intention to describe the hypothetical catalytic center. The catalytic center for PhoD of *Halothece* sp. PCC 7418, using the catalytic center of 2YEQ (in parenthesis) as a template, consisted of Cys 160 (Cys 124), Asp 187 (Asp 151), Tyr 188 (Tyr 152), Asp 242 (Asp 209), Asp 243 (Asp 210), Asn 248 (Asn 215), Asp 249 (Asn 216), Asp 420 (Asp 380) and His 422 (His 382) (Figure 2B). All these amino acids are described in 2YEQ as the active site and coordinate with two Ca^{2+} and one

Fe^{3+} ions [22]. Only one substitution was detected in Asp 249, where in 2YEQ is Asn 216.

Figure 2: Structural analysis of PhoD and PhoU of *Halothece* sp. PCC 7418. **A)** Predicted structure of PhoD of *Halothece* sp. PCC 7418 is represented in red and aligned with the crystal structure 2YEQ that is displayed in blue cyan. **B)** Active center of PhoD of the test bacterium showing all the aminoacids involved in coordination with two Ca^{2+} and Fe^{3+} ions. **C)** Cluster 1 of PhoU, trinuclear metal site with Fe, between $\alpha 5$ and $\alpha 6$. **D)** Cluster 2 of PhoU, tetranuclear metal site with three Fe and one Ni, between $\alpha 2$ and $\alpha 3$. Black aminoacids are from predicted PhoU and red aminoacids are from 4Q25. All the structures were represented with Pymol.

The *in-silico* results described above of PhoD and how it coordinates with Ca^{2+} and Fe^{3+} ions in its active site in *Halothece* sp. PCC 7418 corroborates with the results of the experiment testing the relative importance of PhoD and PhoA in *Halothece* sp. PCC 7418, showing that the APA rates, with the depletion of Mg^{2+} and Zn^{2+} which are the metal co-factors of

PhoA, did not differ considerably with sufficient availability of Mg^{2+} and Zn^{2+} (Figure 3). This suggests that PhoD (and not PhoA) is the more active APase in *Halothece* sp. PCC 7418.

Figure 3: End point assay measurement of Alkaline Phosphatase Activity (APA) under Mg/Zn omission compared with optimal condition. Values are the mean and the error bar is the spanning range between the two duplicate measurements.

Three-Dimensional Structure of PhoU

Annotated PhoU had 224 amino acids (aa) and presented two PhoU domains. Predicted structure of PhoU (C-score = 0.55, estimated TM-score = 0.79 ± 0.09 , estimated RMSD = 4.5 ± 2.9 Å) had 7 α -helix without β -chains. The protein with more

structure homology was PhoU of *P. aeruginosa* (4Q25) of 250 aa with an identity of 32.5% and coverage of 93.3%. Sequence alignment with 4Q25 showed 27.45% of identity and we used this alignment to describe its metal clusters (Figures 2C and 2D). Results showed that *Halotheca* sp. PCC 7418 using 4Q25 as a template displayed at least one metal cluster, and possibly a second one, forming a trinuclear metal site with three Fe and tetranuclear metal site with three Fe and one nickel (Ni). The first cluster was complete and had the same aa as *P. aeruginosa* (in parenthesis) and was formed by Asp 164 (Asp 163), Asp 168 (Asp 167), Glu 202 (Glu 202) and Asp 206 (Asp 206), between α -helixes 5-6 (Figure 2C). The second cluster was incomplete and did not have all the aa that are present in *P. aeruginosa*. Only three aa of seven aa in *P. aeruginosa* (in parenthesis) coincide with *Halotheca* sp. PCC 7418, and this cluster consisted of Asp 61 (Asp 59), Arg 62 (Asp 60), Asp 65 (Asn 63), Tyr 68 (Glu 66), Arg 100 (Ile 98), Arg 104 (Arg 102) and Asp 107 (Asp 105) between α -helix 2-3 (Figure 2D) [49].

Alkaline Phosphatase Activity (APA) in *Halotheca* sp. PCC 7418

Generally, APA rates were significant higher ($p < 0.05$) in [Low NO_3^-] conditions compared with optimal NO_3^- conditions (Figure 4A). Under [Low NO_3^-] APA rates were ≈ 7 times higher in [Low- Medium PO_4^{3-}] and ≈ 77 times higher in [High PO_4^{3-}] compared with their rates under optimal NO_3^- conditions. Moreover, under optimal NO_3^- conditions, APA rates did not have significant differences among the treatments (Figure 4A). Under [Low NO_3^-], treatment combinations of PO_4^{3-} and Fe levels had a significant effect on APA rates (ANOVA, $p < 0.05$), where the rates were significant higher ($p < 0.05$) at the highest Fe levels and at low to medium PO_4^{3-} levels, compared with other treatment combinations (Figure 4A). Figure 4B shows the differences in the kinetics of APA for treatments under [Low NO_3^-] at low and medium PO_4^{3-} levels and low and high Fe levels. At high Fe levels with low to medium PO_4^{3-} levels, the V vs S curve did not reach saturation levels with the maximum S added (5 μM MUF-P). The Vmax and Km, calculated using the

available data for these treatments, were: V_{max} , 4.92 ± 0.56 fmole cell⁻¹ h⁻¹; K_m of 3.47 ± 0.94 μ M at [Low PO₄³⁻- High Fe] and V_{max} , 4.26 ± 0.43 fmole cell⁻¹ h⁻¹; K_m of 7.24 ± 0.57 μ M at [Medium PO₄³⁻- High Fe]. On the contrary to high Fe levels, APase kinetics reached saturation levels with the maximum S added (Figure 4B) at low Fe levels with V_{max} , 1.55 ± 0.19 fmole cell⁻¹ h⁻¹; K_m of 1.53 ± 0.31 μ M at [Low PO₄³⁻- Low Fe] and V_{max} , 1.88 ± 0.06 fmole cell⁻¹ h⁻¹; K_m of 2.02 ± 0.94 μ M at [Medium PO₄³⁻- Low Fe].

The APA rates calculated were considerable higher (up to 6-fold) under NO₃⁻ starvation compared with under NO₃⁻ optimal conditions (Figure 5A). Treatments with [High Fe] in [Low PO₄³⁻] reached the maximum rates at 0.21 ± 0.07 fmole cell⁻¹ h⁻¹ under NO₃⁻ starvation condition, and at 0.03 ± 0.01 fmole cell⁻¹ h⁻¹ under NO₃⁻ optimal condition (Figure 5A). Additions of PO₄³⁻, Fe and/or NO₃⁻ to obtain optimum concentration of PO₄³⁻, Fe and/or NO₃⁻ resulted in reduced APA rates particularly under the initial NO₃⁻ starvation conditions (Figure 5B).

Figure 4: Rates of Alkaline Phosphatase Activity (APA) in *Halothece* sp. PCC 7418 under optimal NO₃⁻ and [Low NO₃⁻] conditions. **A**) End point assay of APA under optimal and [Low NO₃⁻] conditions with 2 μM of MUF-P. **B**) Saturation curve, velocity (V, fmole MUF cell⁻¹ h⁻¹) vs substrate (S, μM) under [Low NO₃⁻] in [Low PO₄³⁻- Low Fe], [Low PO₄³⁻- High Fe], [Medium PO₄³⁻- Medium Fe] and [Medium PO₄³⁻- High Fe], up to 5 μM of MUF-P. PO₄³⁻ is represented as Pi. In A) and B), values are the mean and the error bar is the spanning range between the duplicate measurements. Asterisks (*) indicate significant differences (p < 0.05) between NO₃⁻ optimal and [Low NO₃⁻] conditions of the same PO₄³⁻ and Fe combination treatments, by individual T-student test. Different letters indicate pairwise significant differences (p < 0.05) among treatments in [Low NO₃⁻] using a posthoc test (Bonferroni) after ANOVA over the whole dataset was done.

Figure 5: Rates of Alkaline Phosphatase Activity (APA, fmoles MUF cell⁻¹ h⁻¹) in *Halothece* sp. PCC 7418 under NO₃⁻ starvation conditions compared with the optimal NO₃⁻ treatments under **A**) treatment conditions: [Low PO₄³⁻- Low Fe], [Low PO₄³⁻- High Fe] and [High PO₄³⁻- High Fe] and **B**) under re-inoculum of nutrients in treatment conditions: [Low PO₄³⁻- Low Fe, + PO₄³⁻ + Fe + NO₃⁻] and [Low PO₄³⁻- Low Fe + PO₄³⁻ +Fe]. PO₄³⁻ is represented as Pi. Values are the mean and the error bar is the spanning range between the duplicate measurements.

Phosphorus (P)-Uptake at different Levels of Fe Availability and Fe-Uptake at different Levels of PO_4^{3-} Availability

Generally, P-cellular content varied significantly (ANOVA, $p < 0.05$) under N_2 -fixing conditions [Low NO_3^-] among treatment combinations with significantly higher values at [High PO_4^{3-} -High Fe] treatment compared with other treatment combinations (Figure 6A). On the other hand, specific PO_4^{3-} -uptake rates under N_2 -fixing conditions [Low NO_3^-] and optimal NO_3^- conditions generally did not vary significantly (ANOVA, $p > 0.05$) among treatment combinations (Figure 6B). However, specific t-tests conducted under [Low NO_3^-] conditions, showed PO_4^{3-} -uptake rates to be on average 200 times significantly higher ($p < 0.05$) than the rates under optimal conditions of NO_3^- in T_0 - T_4 and T_0 - T_{10} in low to medium Fe levels (Figure 6B). Different concentrations of Fe in [High PO_4^{3-}] did not show significant differences in PO_4^{3-} -uptake rates ($p > 0.05$) (Figure 6B).

The time course of depletion of total dissolved phosphate (TDP) in the culture media showed that under optimal NO_3^- conditions, the media were depleted with TDP while under NO_3^- starvation conditions, the cells were not capable in depleting TDP from the media (Figure 7A). Fe did not have a significant effect in TDP depletion ($p > 0.05$). The time course of depletion of TDP in the re-inoculum conditions at [Low PO_4^{3-} - Low Fe] (under NO_3^- starvation and NO_3^- optimal conditions), showed the same tendency, in which under NO_3^- starvation conditions, TDP was not depleted (Figure 7B).

Figure 6: A) P-cellular content (fmole P cell⁻¹) under N₂-fixing conditions in *Halotheca* sp. PCC 7418, at different PO₄³⁻ and Fe levels. **B)** Specific net PO₄³⁻-uptake rates (pmole PO₄³⁻-uptake cell⁻¹ d⁻¹) in *Halotheca* sp. PCC 7418 under [Low NO₃⁻] and optimal NO₃⁻ conditions at three different periods (T₀-T₁, T₀-T₄ and T₄-T₁₀). PO₄³⁻ is represented as Pi. Values are the mean and the error bar is the spanning range between the duplicate measurements. Different letters indicate pairwise significant differences (p < 0.05) among treatments using a posthoc test (Bonferroni) after ANOVA over the whole dataset was done, and asterisks (*) indicate significant differences (p < 0.05) between [Low NO₃⁻] and NO₃⁻ optimal conditions by individual t-student tests for each treatment combination of PO₄³⁻ and Fe.

Figure 8 shows the specific Fe-uptake rates at different levels of PO₄³⁻ and Fe under N₂-fixing conditions. Results showed that generally, specific Fe-uptake rates varied significantly at different treatment combinations of PO₄³⁻ and Fe (ANOVA, p < 0.05). Fe-uptake rates were significantly higher (p < 0.05) at [High PO₄³⁻] conditions compared to [Low PO₄³⁻] and [Medium

PO_4^{3-}] conditions. There were also significant differences ($p < 0.05$) of increased Fe-uptake with increasing availability of Fe.

Phosphorus (P)-Cellular Content and its Relationship with N_2 Fixation and Fe-Cellular Content

Phosphorus cellular content of *Halothece* sp. PCC 7418 showed significant positive linear correlation with N_2 fixation rates ($p < 0.05$, $r^2 = 0.86$, $n = 12$) (Figure 9A). Moreover, the P-cellular content of the cells showed significant positive linear correlation with their Fe contents ($p < 0.05$, $r^2 = 0.71$, $n = 18$) (Figure 9B). The P and Fe-cellular contents of the cells did not show significant correlations with other metals (*i.e.* Mn).

Discussion

Pho Regulon and the Three-Dimensional Structure of PhoD and PhoU of *Halothece* sp. PCC 7418: Elucidating the role of Fe as Co-Factor

The Pho regulon of *Halothece* sp. PCC 7418 is composed of genes whose protein products are involved in different functions: autokinase activity of PhoR and phosphate transport (PhoU); high-affinity phosphate transport (PstS, PstC, PstA and PstB), in a two-component regulatory system (PhoR-PhoB); extracellular enzymes capable of obtaining PO_4^{3-} from organic phosphates (Alkaline Phosphatases, APases); and polyphosphate metabolism (PpK, PpX and PpA) [19]. However, no low-affinity transporters were annotated while some studies demonstrated that this strain exhibit low-affinities transporters [50]. *Halothece* sp. PCC 7418 contains a Pho regulon with 11 distinct genes in single or multiple copies altogether accounting 26 distinct loci in the whole genome, suggesting that *Halothece* sp. PCC 7418 is well adapted to survive to P-limiting conditions. In model strains whose P-acquisition mechanisms are well studied such as *Trichodesmium* spp. and *Crocospaera watsonii* they only have 15 copies and 19 copies respectively in their Pho regulon [28,29].

Genome analysis indicated that *Halotheca* sp. PCC 7418 and *Gloeocapsa* sp. PCC 7428 were the strains with more copies of Alkaline Phosphatases (APase), 8 and 19, respectively (Figure 1). These two cyanobacteria are halotolerant species, and there are studies that suggest that salt stress enhance APA in halophytic strains [51]. In a previous study [51], *Halotheca* sp. PCC 7418 only showed three APases: two PhoA and one PhoD. Of the 8 APases found in our study for the same species, one of them is also annotated as PhoD and the rest are not annotated to a specific type of APase. PhoD, together with PhoX, is one of the most abundant APases in marine habitats and its activity may be controlled by availability of its metal co-factor(s) (e.g. Fe^{3+} , Ca^{2+} , Mg^{2+} , Zn^{2+}) [32,52].

Three-dimensional analyses with PhoD of *Halotheca* sp. PCC 7418 revealed its active site as a homologue to the crystal structure of PhoD of *B. subtilis* with two Ca^{2+} and one Fe^{3+} ions as co-factors [22] (Figure 2B). Previous studies on APase activity in *Halotheca* sp. PCC 7418 indicated Ca^{2+} dependence of PhoD [51] but the Fe^{3+} dependence was not investigated. The experiment conducted here wherein the omission of Mg^{2+} and Zn^{2+} (but not Fe^{3+} in the culture medium) did not result in any significant changes in APase activity (Figure 3), suggesting that the APases of *Halotheca* sp. PCC 7418 (i.e. PhoD) do not require these metals (Mg^{2+} and Zn^{2+}) as co-factors as in the case of PhoA [51], and the most active APase could be PhoD.

Iron is not only important as a co-factor for the activities of APase but can be essential in other components of Pho regulon like PhoU in which the results of the 3D-dimensional analyses in this study showed PhoU of *Halotheca* sp. PCC 7418 forming at least one Fe-containing metal cluster, and possible a second cluster (Figures 2C and 2D), using as a model, the PhoU of *P. aeruginosa* (4Q25). PhoU can participate in the PO_4^{3-} transport across the cell membranes of bacteria in the regulation of the phosphate-specific transport systems [19] and in controlling cellular phosphate metabolism [53]. The specific role of PhoU in *Halotheca* sp. PCC 7418, however, remains to be investigated.

Alkaline Phosphatase Activity (APA) in *Halothece* sp. PCC 7418: Experimental Analysis of Regulation by Fe, NO₃⁻, and P Availabilities

Experimental measurements of APA in *Halothece* sp. PCC 7418 under different levels in Fe availability revealed higher activities with higher levels of Fe (Figures 4A, 4B and Figure 5A), confirming the regulatory role of Fe in the APase (*i.e.* PhoD) in this species as we predicted in our 3D-structural analyses of its PhoD (Figure 2B). However, the effect of Fe availability on the rates of APA in *Halothece* sp. PCC 7418 depends on the availability of inorganic sources of nitrogen (*i.e.* NO₃⁻) wherein at low NO₃⁻ concentrations, increasing Fe availability enhanced the APA rates (Figures 4A and 4B). We showed that under [Low NO₃⁻] and at high Fe levels, APA was not saturated (Figure 4B). We hypothesized that under these conditions, the V_{max} of APases from *Halothece* sp. PCC 7418 is so high that increasing MUF-P concentrations, up to 10 μM (in the other assays that were additionally conducted), was not high enough to saturate the enzyme because of the enhancement of APA by high levels of the Fe co-factor.

At high or optimal NO₃⁻ concentrations, APA rates in general are lower than in NO₃⁻ starvation conditions (Figure 5A) and even lower than in [Low NO₃⁻] treatments (Figure 4A). These results can be due to peculiar characteristics of the N₂ fixation process. High concentrations of readily assimilable forms of dissolved inorganic nitrogen (DIN, *i.e.* NH₄, NO₃⁻) are known to inhibit N₂ fixation as evidenced by DIN inhibition studies [54]. The N₂ fixation process ($\text{N}_2 + 8\text{e}^- + 16\text{ATP} + 8\text{H}^+ \rightarrow 2\text{NH}_3 + \text{H}_2 + 16\text{ADP} + 16\text{PO}_4^{3-}$) is an energetically costly processes requiring 16 ATPs and 25% more energy is needed to reduce N₂ than to reduce NO₃⁻ to NH₄. A N₂-fixing cell such as *Halothece* sp. PCC 7418 would rather reduce first the available NO₃⁻ than to fix N₂. Conversely, the N₂-fixing process is stimulated with low NO₃⁻ availability [55,56]. Since the energy (ATP) to fuel N₂ fixation is dependent on PO₄³⁻, the demand for PO₄³⁻ is theoretically enhanced when the cells are doing N₂ fixation (in conditions under low NO₃⁻ availability). Thus, APase activities are expected to be stimulated under low NO₃⁻ conditions, and

consequently depend on the availability of Fe because APases such as PhoD may have Fe as co-factor. Moreover, Fe is an important structural component of the nitrogenase enzyme catalyzing the N_2 fixation process. Nitrogenase contains 38 Fe atoms per holoenzyme since nitrogenase is characterized by slow reaction rates the N_2 -fixers need a large cellular pool of this enzyme, and thus more Fe is needed [57]. The enhanced rates of APase under N_2 -fixing conditions (low NO_3^- availability) and high Fe availability with low PO_4^{3-} levels is expected as APases activities are induced with low PO_4^{3-} levels in the medium [58]. The control of NO_3^- and PO_4^{3-} availabilities in APase activities for N_2 -fixing cells such as *Halothece* sp. PCC 7418 is further supported here with the results of decreased APA rates when NO_3^- , PO_4^{3-} and Fe were added to cells growing previously with low PO_4^{3-} , low Fe and/or low NO_3^- levels (Figure 5B).

Phosphorus (P) and Fe-Uptake and Cellular Contents in *Halothece* sp. PCC 7418

The PO_4^{3-} -uptake measurements in *Halothece* sp. PCC 7418 were done in the experimental units with high PO_4^{3-} levels because (1) the method used for PO_4^{3-} analyses was not sensitive enough to measure very low levels of PO_4^{3-} ($\leq 0.1 \mu M$) and (2) APase activities are not induced at high PO_4^{3-} levels allowing us to evaluate if Fe is also important in PO_4^{3-} transport mechanisms and not only in APase activities. PO_4^{3-} -uptake rates of *Halothece* sp. PCC 7418 was significantly higher under N_2 -fixing conditions ([Low NO_3^-]) than in non- N_2 fixing conditions due to the high demand of P for the energy costly N_2 fixation (Figure 6B). The dependence of N_2 fixation on P in *Halothece* sp. PCC 7418 is evidenced here with the significant linear correlation between cellular P content of the cells and their rates of N_2 fixation (Figure 9A), consistent with studies carried out in *Trichodesmium* spp. in the Atlantic [12]. In addition, not only N_2 -fixing conditions can enhance the P-requirements of cyanobacteria. It is also reported that under nitrogen limitation, phytoplankton can accumulate carbohydrates and phospholipids, increasing their P-cellular content [59]. Different concentrations

of Fe, however, did not show significant differences in PO_4^{3-} -uptake at high levels of PO_4^{3-} availability. This suggests that PO_4^{3-} -uptake mechanisms in this case are not dependent on Fe levels or the Fe present in all treatments (from low to high Fe concentrations) are sufficient for the cells (Figure 6B). The latter case may be most likely since we found significant correlation between P-cellular and Fe-cellular content of the cells (Figure 9B). These results are also consistent with our data that highest P-cellular content was found at high Fe levels (Figure 6A), suggesting the narrow connection between P and Fe. The relation between P and Fe cellular contents is also supported by evidences that high concentrations of elemental P are found associated (or co-localized spatially) with Fe within the cells of phytoplankton (*Chlorella* sp. and *Chlamydomonas* sp. [60]. The Fe-uptake measurements in *Halotheca* sp. PCC 7418 in N_2 -fixing conditions revealed that Fe-uptake was correlated with P with high Fe-uptake rates at higher PO_4^{3-} levels (Figure 8). This may be due to the P-dependence (ATP) of Fe transporters [61,62]. Results also show the tendency of higher Fe-uptake rates in higher concentrations of Fe in the media, suggesting a passive transport of this metal in *Halotheca* sp. PCC 7418. However, this needs to be further investigated.

The time course of depletion of total dissolved phosphate (TDP) in the media (Figure 7A), showed that under NO_3^- starvation conditions, cells did not deplete TDP, and even increased at the final stage of the experiment suggesting a liberation of cellular TDP of dying cells. Extreme NO_3^- starvation conditions are suggested here to be detrimental to the growth of *Halotheca* sp. PCC 7418 and may have consequences on their P-uptake mechanisms, explaining why APA rates were lower than in [Low NO_3^-] conditions. Even when the nutrients (PO_4^{3-} , Fe and/or NO_3^-) were re-inoculated in the cultures that were previously starved with NO_3^- , the cells did not acclimate and were not capable to deplete TDP from the media (Figure 7B). Whereas many of the researches have focused on the inhibition or sensitivity of N_2 fixation to increased availability of dissolved inorganic nitrogen (e.g. NO_3^- , NH_4^+) [54], investigations on the physiological conditions for growth of N_2 -fixers are few. Spiller

and Shanmugan [63], gave some evidences that a unicellular species of marine N_2 -fixer *Synechococcus* sp. strain SF1 (isolated from macroalgae, *Sargassum fluitans*) is dependent on the presence and type of carbon (C) source to support its growth with N_2 as the sole nitrogen source. Their results showed for example that without the addition of C source (*e.g.* HCO_3^-), there was no growth of the species tested with N_2 as the sole source. Moreover, some studies have reported less cell yield of unicellular N_2 fixers when grown with N_2 as sole N source compared with addition of NO_3^- since N_2 fixation is an energetically costly process [63,64]. Our result that extreme NO_3^- starvation condition (at nM levels close to N_2 as sole source) is suggested to be detrimental to the growth of *Halotheca* sp. PCC 7418 may be due to the type of C source (glucose and citrate) in our treatments which may not be the optimum for growth of this species with N_2 as sole N source. This hypothesis however needs more investigations.

Figure 7: TDP (μM) under NO_3^- starvation and optimal NO_3^- conditions. **A)** TDP consumption (μM) under NO_3^- starvation and NO_3^- optimal conditions. **B)** TDP consumption in the re-inoculated conditions in [Low PO_4^{3-} - Low Fe] (under NO_3^- starvation and NO_3^- optimal conditions). PO_4^{3-} is represented as Pi. Values are the mean and the error bar is the spanning range between the duplicate measurements. In A) letters indicate significant differences ($p < 0.05$) by T-student test.

Figure 8: Fe-uptake in *Halothece* sp. PCC 7418 (fmole Fe cell⁻¹ d⁻¹) measured as the difference of the initial Fe concentration and the final Fe concentration at day 10 day of the experiment under [Low NO₃⁻]. PO₄³⁻ is represented as Pi. Values are the mean and the error bar is the spanning range between the duplicate measurements, and letters indicate significant differences (p < 0.05) between treatments using a posthoc test (Bonferroni) after ANOVA over the whole dataset was done.

Figure 9: Linear regression analyses **A)** between P-cellular content and N₂ fixation and **B)** between P and Fe-cellular content. Cellular content was measured in [Low NO₃⁻] using pooled data from all PO₄³⁻ and Fe combination treatments.

In summary, this is the first study investigating the interaction between PO_4^{3-} , Fe and NO_3^- availabilities in the P-acquisition mechanisms of a unicellular N_2 -fixing bacteria found in association with the Mediterranean seagrass *P. oceanica*. The results suggest that APase activities under inorganic P-limited conditions are enhanced by increased Fe availabilities. The PO_4^{3-} and Fe dependence of *Halotheca* sp. PCC 7418 depends whether they are grown in N_2 -fixing conditions (*i.e.* low NO_3^- levels) or not. Genomic and structural analyses have also shown the tight association between P-acquisition mechanisms and Fe in *Halotheca* sp. PCC 7418. Studies combining genomic and protein structural analyses and experimental approaches are important to investigate in detail the control of environmental factors (*e.g.* availability of metals and nutrients) to the functioning of N_2 -fixing organisms found in important species of seagrasses.

References

1. Agawin NSR, Ferriol P, Cryer C, Alcon E, Busquets A, et al. Significant nitrogen fixation activity associated with the phyllosphere of Mediterranean seagrass *Posidonia oceanica*: First report. *Marine Ecology Progress Series*. 2016; 551: 53-62.
2. Campagne CS, Salles JM, Boissery P, Deter J. The seagrass *Posidonia oceanica*: Ecosystem services identification and economic evaluation of goods and benefits. *Marine Pollution Bulletin*. 2015; 97: 391-400.
3. Gutiérrez JL, Jones CG, Byers JE, Arkema KK, Berkenbusch K, et al. Physical ecosystem engineers and the functioning of estuaries and coasts. *Treatise on Estuarine and Coastal Science*. 2012; 7: 58-81.
4. Agawin NSR, Ferriol P, Sintes E, Moyà G. Temporal and spatial variability of in situ nitrogen fixation activities associated with the Mediterranean seagrass *Posidonia oceanica* meadows. *Limnology and Oceanography*. 2017; 62: 2575-2592.
5. Agawin NSR, Ferriol P, Sintes E. Simultaneous measurements of nitrogen fixation activities associated with

- different plant tissues of the seagrass *Posidonia oceanica*. *Marine Ecology Progress Series*. 2019; 611: 111-127.
6. Sohm JA, Webb EA, Capone DG. Emerging patterns of marine nitrogen fixation. *Nature Reviews Microbiology*. 2011; 9: 499-508.
 7. Agawin NSR, Duarte CM, Agusti S. Nutrient and temperature control of the contribution of picoplankton to phytoplankton biomass and production. *American Society of Limnology and Oceanography*. 2000; 45: 591–600.
 8. Herrero A, Flores E. *The cyanobacteria: molecular biology, genomics, and evolution*. Norfolk: Caister Academic Press. 2008.
 9. Geider RJ, La Roche J. Redfield revisited: variability of C:N:P in marine microalgae and its biochemical basis. *European Journal of Phycology*. 2002; 37: 1-17.
 10. Quigg A, Irwin AJ, Finkel ZV. Evolutionary inheritance of elemental stoichiometry in phytoplankton. *Proceedings of the Royal Society B: Biological Sciences*. 2011; 278: 526–534.
 11. Redfield AC. On the proportions of organic derivatives in sea water and their relation to the composition of plankton. *James Johnstone Memorial Volume*, University Press of Liverpool. 1934; 176-192.
 12. Sañudo-Wilhelmy SA, Kustka AB, Gobler CJ, Hutchins DA, Yang M, et al. Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central Atlantic Ocean. *Nature*. 2001; 411: 66-69.
 13. Schwarz R, Forchhammer K. Acclimation of unicellular cyanobacteria to macronutrient deficiency: emergence of a complex network of cellular responses. *Microbiology*. 2005; 151: 2503-2514.
 14. Tandeau de Marsac N, Houmard J. Adaptation of cyanobacteria to environmental stimuli: new steps towards molecular mechanisms. *FEMS Microbiology Letters*. 1993; 104: 119-189.
 15. Mills MM, Ridame C, Davey M, La Roche J, Geider RJ. Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic. *Nature*. 2004; 429: 232-294.
 16. Moore C, Mills MM, Achterberg EP, Geider RJ, Laroche J, et al. Large-scale distribution of Atlantic nitrogen fixation

- controlled by iron availability. *Nature Geoscience*. 2009; 2: 867-871.
17. Moore C, Mills MM, Arrigo KR, Berman-Frank I, Bopp L, et al. Processes and patterns of oceanic nutrient limitation. *Nature Geoscience*. 2013; 6: 701-710.
 18. Kornberg A. Inorganic polyphosphate: toward making a forgotten polymer unforgettable. *Journal of Bacteriology*. 1995; 177: 491-496.
 19. Santos-Beneit F. The Pho regulon: a huge regulatory network in bacteria. *Frontiers in Microbiology*. 2015; 6.
 20. Tiwari B, Singh S, Kaushik MS, Mishra AK. Regulation of organophosphate metabolism in cyanobacteria. A review. *Microbiology*. 2015; 84: 2913-2915.
 21. Yong SC, Roversi P, Lillington J, Rodriguez F, Krehenbrink M, et al. A complex iron-calcium cofactor catalyzing phosphotransfer chemistry. *Science*. 2014; 345: 1170-1173.
 22. Rodriguez F, Lillington J, Johnson S, Timmel CR, Lea SM, et al. Crystal structure of the *Bacillus subtilis* phosphodiesterase PhoD reveals an iron and calcium-containing active site. *Journal of Biological Chemistry*. 2014; 289: 308893-99.
 23. Blanco AG, Sola M, Gomis-Rüth FX, Coll M. Tandem DNA Recognition by PhoB, a two-component signal transduction transcriptional activator. *Structure*. 2002; 10: 701-713.
 24. Yuan ZC, Zaheer R, Morton R, Finan TM. Genome prediction of PhoB regulated promoters in *Sinorhizobium meliloti* and twelve proteobacteria. *Nucleic Acids Research*. 2006; 34: 2686-2697.
 25. Gardner SG, Johns KD, Tanner R, McCleary WR. The PhoU protein from *Escherichia coli* interacts with PhoR, PstB, and metals to form a phosphate-signaling complex at the membrane. *Journal of Bacteriology*. 2014; 196: 1741-1752.
 26. Su Z, Olman V, Xu Y. Computational prediction of Pho regulons in cyanobacteria. *BMC Genomics*. 2007; 8: 1-12.
 27. Browning TJ, Achterberg EP, Yong JC, Rapp I, Utermann C, et al. Iron limitation of microbial phosphorus acquisition in the tropical North Atlantic. *Nature Communications*. 2017; 8: 1-7.
 28. Dyhrman ST, Haley ST. Phosphorus scavenging in the unicellular marine diazotroph *Crocospaera watsonii*.

- Applied and Environmental Microbiology. 2006; 72: 1452-1458.
29. Fu FX, Zhang Y, Bell PRF, Hutchins DA. Phosphate uptake and growth kinetics of *Trichodesmium* (Cyanobacteria) isolates from the North Atlantic Ocean and the Great Barrier Reef, Australia. *Journal of Phycology*. 2005; 41: 62-73.
 30. Tanhua T, Hainbucher D, Schroeder K, Cardin V, Álvarez M, et al. The Mediterranean Sea system: a review and an introduction to the special issue. *Ocean Science*. 2013; 9: 789-803.
 31. Statham PJ, Hart V. Dissolved iron in the Cretan Sea (eastern Mediterranean). *Limnology and Oceanography*. 2005; 50: 1142-1148.
 32. Luo H, Benner R, Long RA, Hu J. Subcellular localization of marine bacterial alkaline phosphatases. *Proceedings of the National Academy of Sciences*. 2009; 106: 21219-21223.
 33. Wattam AR, Davis JJ, Assaf R, Boisvert S, Brettin T, et al. Improvements to PATRIC, the all-bacterial bioinformatics database and analysis resource center. *Nucleic Acids Research*. 2017; 45: D535-D542.
 34. Zhang Y. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics*. 2008; 9: 40.
 35. Finn RD, Coghill P, Eberhardt RY, Eddy SR, Mistry J, et al. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Research*. 2016; 44: D279-D285.
 36. Li Z, Natarajan P, Ye Y, Hrabec T, Godzik A. POSA: A user-driven, interactive multiple protein structure alignment server. *Nucleic Acids Research*. 2014; 42: W240–W245.
 37. The UniProt Consortium. UniProt: a hub for protein information. *Nucleic Acids Research*, 43(Database issue). 2014; D204-212.
 38. DeLano W. The PyMOL Molecular Graphics System. Schrödinger, LLC. 2002. Available Online at: www.pymol.org
 39. Stanier RY, Deruelles J, Rippka R, Herdman M, Waterbury JB. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *Microbiology*. 1979; 111: 1-61.

40. Ascenzi P, Amiconi G. Logarithmic plots in enzymology. Representation of the Michaelis-Menten equation. *Biochemical Education*. 1987; 15: 83-84.
41. Hansen HP, Koroleff F. Determination of nutrients. In: K Grasshoff, K Kremling, M Ehrhardt, editors. *Methods of Seawater Analysis*. 2007.
42. Tovar-sanchez A, Sañudo-Wilhelmy SA, Kustka AB, Agustí S, Dachs J, et al. Effects of dust deposition and river discharges on trace metal composition of *Trichodesmium* spp. in the tropical and subtropical North Atlantic Ocean. *Limnology and Oceanography*. 2006; 51: 1755–1761.
43. Tovar-Sanchez A, Sañudo-Wilhelmy SA. Influence of the Amazon River on dissolved and intra-cellular metal concentrations in *Trichodesmium* colonies along the western boundary of the sub-tropical North Atlantic Ocean. *Biogeosciences*. 2011; 8: 217–225.
44. Ghaffar S, Stevenson RJ, Khan Z. Effect of phosphorus stress on *Microcystis aeruginosa* growth and phosphorus uptake. *PLoS ONE*. 2017; 12: e0174349.
45. Reddy KJ, Haskell JB, Sherman DM, Sherman LA. Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus *Cyanothece*. *Journal of Bacteriology*. 1993; 175: 1284-1292.
46. Agawin NSR, Benavides M, Busquets A, Ferriol P, Stal LJ, et al. Dominance of unicellular cyanobacteria in the diazotrophic community in the Atlantic Ocean. *Limnology and Oceanography*. 2014; 59: 623-637.
47. Stal LJ. Acetylene reduction technique for assay of nitrogenase. *Methods Enzymol*. 1988; 167: 474-484.
48. Jensen BB, Cox RP. Direct measurements of steady-state kinetics of cyanobacterial N₂ uptake by membrane-leak mass spectrometry and comparisons between nitrogen fixation and acetylene reduction. *Applied and Environmental Microbiology*. 1983; 45: 1331-1337.
49. Lee SJ, Park YS, Kim SJ, Lee BJ, Suh SW. Crystal structure of PhoU from *Pseudomonas aeruginosa*, a negative regulator of the Pho regulon. *Journal of Structural Biology*. 2014; 188: 22-29.
50. Tripathi K, Sharma NK, Kageyama H, Takabe T, Rai AK. Physiological, biochemical and molecular responses of the

- halophilic cyanobacterium *Aphanothece halophytica* to Pi-deficiency. *European Journal of Phycology*. 2013; 48: 461-473.
51. Kageyama H, Tripathi K, Rai AK, Cha-Um S, Waditee-Sirisattha R, et al. An alkaline phosphatase/phosphodiesterase, PhoD, induced by salt stress and secreted out of the cells of *Aphanothece halophytica*, a halotolerant cyanobacterium. *Applied and Environmental Microbiology*. 2011; 77: 5178-5183.
 52. Zeng YF, Ko TP, Lai HL, Cheng YS, Wu TH, et al. Crystal structures of *Bacillus* alkaline phytase in complex with divalent metal ions and inositol hexasulfate. *Journal of Molecular Biology*. 2011; 409: 214-224.
 53. Lubin EA, Fiebig A, Laub MT, Henry JT, Crosson S. Identification of the PhoB regulon and role of PhoU in the phosphate starvation response of *Caulobacter crescentus*. *Journal of Bacteriology*. 2015; 198: 187-200.
 54. Knapp AN. The sensitivity of marine N₂ fixation to dissolved inorganic nitrogen. *Frontiers in Microbiology*. 2012; 3: 1-14.
 55. Manhart JR, Wong PP. Nitrate Effect on Nitrogen Fixation (Acetylene Reduction): Activities of legume root nodules induced by rhizobia with varied nitrate reductase activities. *Plant physiology*. 1980; 65: 502-505.
 56. Nelson DC, Waterbury JB, Jannasch HW Nitrogen fixation and nitrate utilization by marine and freshwater *Beggiatoa*. *Arch. Microbiol*. 1982; 133: 172.
 57. Hoffman BM, Lukoyanov D, Yang ZY, Dean DR, Seefeldt LC. Mechanism of nitrogen fixation by nitrogenase: The next stage. *Chemical Reviews*. 2014; 114: 4041-4062.
 58. Romano S, Bondarev V, Kölling M, Dittmar T, Schulz-Vogt HN. Phosphate limitation triggers the dissolution of precipitated iron by the marine bacterium *Pseudovibrio* sp. FO-BEG1. *Frontiers in Microbiology*. 2017; 8: 1-11.
 59. Liefer JD, Garg A, Fyfe MH, Irwin AJ, Benner I, et al. The Macromolecular Basis of Phytoplankton C:N:P Under Nitrogen Starvation. *Frontiers in Microbiology*. 2019; 10: 1-16.
 60. Diaz J, Ingall E, Vogt S, de Jonge MD, Paterson D, et al. Characterization of phosphorus, calcium, iron, and other

- elements in organisms at sub-micron resolution using X-ray fluorescence spectromicroscopy. *Limnology and Oceanography: Methods*. 2009; 7: 42-51.
61. Kranzler C, Rudolf M, Keren N, Schleiff E. Iron in Cyanobacteria. *Advances in Botanical Research*. 2013; 65: 57-105.
 62. Noinaj N, Guillier M, Barnard TJ, Buchanan SK. TonB-dependent transporters: regulation, structure, and function. *Annual Review of Microbiology*. 2010; 64: 43-60.
 63. Spiller H, Shanmugam KT. Physiological conditions for nitrogen fixation in a unicellular marine cyanobacterium, *Synechococcus* sp. strain SF1. *Journal of bacteriology*. 1987; 169: 5379–5384.
 64. Agawin NSR, Rabouille S, Veldhuis MJW, Servatius L, Hol S, et al. Competition and facilitation between unicellular nitrogen-fixing cyanobacteria and non – nitrogen-fixing phytoplankton species. *Limnology And Oceanography*. 2007; 52: 2233–2248.