Book Chapter

Generation of Hematopoietic-Like Stem Cells from Adult Human Peripheral Blood Following the Treatment with Platelet-Derived Mitochondria

Haibo Yu, Wei Hu, Xiang Song, Dante Descalzi-Montoya, Zheng Yang, Robert Korngold and Yong Zhao*

Center for Discovery and Innovation, Hackensack Meridian Health, USA

***Corresponding Author:** Yong Zhao, Center for Discovery and Innovation, Hackensack Meridian Health, Nutley, NJ 07110, USA

Published **November 16, 2020**

This Book Chapter is a republication of an article published by Yong Zhao, et al. at International Journal of Molecular Sciences in June 2020. (Yu, H.; Hu, W.; Song, X.; Descalzi-Montoya, D.; Yang, Z.; Korngold, R.; Zhao, Y. Generation of Hematopoietic-Like Stem Cells from Adult Human Peripheral Blood Following Treatment with Platelet-Derived Mitochondria. *Int. J. Mol. Sci.* 2020, 21, 4249.)

How to cite this book chapter: Haibo Yu, Wei Hu, Xiang Song, Dante Descalzi-Montoya, Zheng Yang, Robert Korngold, Yong Zhao. Generation of Hematopoietic-Like Stem Cells from Adult Human Peripheral Blood Following the Treatment with Platelet-Derived Mitochondria. In: Sławomir Lach, editor. Prime Archives in Molecular Sciences. Hyderabad, India: Vide Leaf. 2020.

© The Author(s) 2020. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which

permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Funding: This research received no external funding.

Acknowledgments: We are grateful to Mr. Poddar and Mr. Ludwig for generous funding support via Hackensack UMC Foundation. We thank Drs. Hal E. Broxmeyer and Jason Butler for helpful discussions and critical comments on the manuscript. We thank Drs. Bryan DeBusk and H. Nikki March and Laura Zhao for editing this manuscript.

Authors contribution: Y.Z.: initiated the study, supervised experiments, and contributed to concepts, experimental design, data analysis and interpretation, manuscript writing, and final approval of manuscript; H.Y.: performed most experiments and data analysis; W.H., and X.S.: performed stem cell culture and flow cytometry. H.Y., D.D., Z.Y. performed animal studies. R.K. helped in data analysis and revising manuscript.

Conflicts of Interest: This work has been submitted for a provisional patent application (Application No. 62/976 830). Y.Z, H.Y, W.H. and X.S. were listed as inventors. All other authors have no financial interests that may be relevant to the submitted work.

Abstract

Adult stem cells represent a potential source for cellular therapy to treat serious human diseases. We characterized the insulin-producing cells from adult peripheral blood (designated PB-IPC), displaying a unique phenotype. Mitochondria are normally located in cellular cytoplasm where they generate ATP to power the cell's functions. *Ex vivo* and *in vivo* functional studies established that treatment with platelet-derived mitochondria can reprogram the transformation of adult PB-IPC into functional CD34⁺ hematopoietic stem cells (HSC-like cells), leading to the production of blood cells such as T cells, B cells, monocytes/macrophages, granulocytes, red blood cells, and megakaryocytes (MK)/platelets. These findings reveal a novel

function of mitochondria directly contributing to cellular reprogramming, thus overcoming the limitations and safety concerns by using conventional technologies to reprogram embryonic stem (ES) and induced pluripotent stem (iPS) cells in regenerative medicine.

Keywords

Mitochondria; PB-IPC; Hematopoietic Stem Cells; Differentiation; Blood

Introduction

Stem cell research has the potential to revolutionize treatments for certain life-changing injuries and devastating human diseases such as diabetes and Alzheimer's disease. To date, researchers have characterized multiple types of human stem cells with varying potentials for regeneration. Animal studies and human trials have demonstrated their translational capability to treat human diseases. For over three decades now, the most common stem cell therapy approved by the FDA has been hematopoietic cell transplantation (HCT) (also termed hematopoietic stem cell transplantation, HSCT) for treatment of bone marrow failure, malignant blood disorders, post chemotherapy and/or radiation cell regeneration, genetic-based blood disorders, and autoimmune diseases [1-3]. However, several major limitations have restricted the broad clinical application of allogeneic HCT, including the difficulty in identifying a human leukocyte antigen (HLA) fully matched or haploidentical donor, the scarcity of CD34⁺ hematopoietic stem cells (HSC) amongst all sources of harvested cells ($\leq 1\%$) [4] but particularly limited by the incidence of graft-versus-host disease (GVHD), opportunistic infections, relapse of primary disease and toxicities associated with immunosuppressive drugs and radiation. An autologous source of HSC would address the problems of matching and GVHD, but engraftment could still be hampered by the limited number of CD34⁺ HSC. Since the success rate of engraftment for clinical HCT is correlated with the number of functional CD34⁺ hematopoietic progenitor cells (HPC) and HSC in the transplant [5,6], researchers have evaluated whether embryonic stem (ES)

cells and induced pluripotent stem (iPS) cells can be manipulated to produce HSC through reprogramming by small molecules or by viral transduction of transcription factors [7-12]. Thus far, these approaches have been limited by an inability to generate true functional HSC in sufficient numbers for therapeutic use, as well as safety and ethical concerns and potential immune rejection issues to ES- or iPS derivatives [5,13]. Alternative approaches are needed to circumvent these limitations.

We previously characterized a new type of multipotent stem cell from human cord blood (designated cord blood-derived multipotent stem cells, CB-SC) [14] that is distinguished from other known stem cell types including HSC and mesenchymal stem cells (MSC). Based on the comprehensive immune modulation characteristics of CB-SC [15], we developed the Stem Cell Educator (SCE) therapy, which consists of isolating the patient's blood cells through a blood cell separator, co-culturing the patient's immune cells with adherent CB-SCs *in vitro*, and returning the "educated" immune cells to the patient's circulation [15-17]. Our clinical trials indicate that SCE therapy reverses multiple immune dysfunctions and corrects autoimmune memory [17], promotes regeneration of islet β cells, improves metabolic control for the treatment of type 1 diabetes (T1D) [15,17,18] and type 2 diabetes (T2D) [16,18], and treats other autoimmune diseases such as hair regrowth in alopecia areata (AA) [19]. Using a similar previously utilized approach for an isolation of CB-SC [15-19], we characterized adult peripheral blood insulin-producing cells (PB-IPC) [20] from adult peripheral blood by virtue of their ability adhering to the hydrophobic surface with positive charge. While investigating the mechanisms underlying the effects of SCE therapy in *ex vivo* studies [18], we found the migration of platelet-released mitochondria to pancreatic islets, where they can enhance islet β -cell function and C-peptide (an insulin byproduct) production, supporting clinical observations of the long-lasting improved health status in subjects with T1D and T2D post receiving SCE therapy [18]. Subsequently, we sought to determine whether platelets-derived mitochondria could directly influence the functions or reprogram the differentiation of adult PB-IPC. Notably, our recent *ex vivo* studies established that

mitochondrion-induced PB-IPC (miPB-IPC) can give rise to retinal pigment epithelium (RPE) cells and neuronal cells in the presence of different inducers [21]. Here, we found that platelets-derived mitochondria may be used to transform PB-IPC into CD34⁺ hematopoietic stem cell-like cells, giving rise to multiple blood cell lineages.

Results

Ex vivo Differentiation of Mitochondrion-induced PB-IPC (miPB-IPC) into the Mitochondrion-induced CD34⁺ HSC-like Cells (miCD34⁺ HSC) after the Treatment with Platelet-Derived Mitochondria

Platelets are enucleate cells without human genomic DNA. We obtained high purity apheresis platelets (> 99% of CD41⁺CD42⁺ platelets [18]) from New York Blood Center for our studies. To determine the purity of isolated mitochondria from platelets, different markers were applied by flow cytometry including MitoTrack Deep Red staining, anti-cytochrome C and anti-heat shock protein (HSP) 60 Abs for mitochondrial markers, calnexin for endoplasmic reticulum (ER), and GM130 for Golgi apparatus. Flow cytometry demonstrated that 99% of isolated mitochondria were positive for MitoTrack Deep Red, HSP 60 and cytochrome C; there were about 5% of cytochrome C⁺ calnexin⁺ and 4% of cytochrome C⁺ GM130⁺ (**Figure 1A**). These double-positive staining might be caused by the interaction and conjugation of mitochondria with ER or Golgi apparatus respectively. Flow cytometry demonstrated the purity of isolated mitochondria was $\geq 90\%$. (**Figure 1A**). We prepared the purified platelet-derived mitochondria from autologous or allogeneic peripheral blood, as described previously [18] and treated PB-IPC that were isolated and expanded from blood samples of adult donors at the New York Blood Center (n = 51; mean age of 48.76 ± 14.97 ; age range from 18 to 72 years old; 24 males and 27 females). Notably, the expression of HSC marker CD34 was up-regulated on PB-IPC after the treatment with mitochondria. A phenotypic analysis of miPB-IPC after two weeks of mitochondria treatment was striking in that the expression of CD34 on miPB-IPC increased from $0.71 \pm 0.25\%$ to $14.8 \pm 3.1\%$ ($P = 7.88\text{E-}06$, n = 5) (**Figure 1B**). Using an optimized panel of

cell markers [22], we found that mitochondrion-induced CD34⁺ (miCD34⁺) cells displayed a phenotype of CD34⁺CD38⁻/_{low}CD45RA⁻CD49f⁺CD90⁺Flt3^{-/low}CD7⁺CD10⁺CD71⁺BAH1^{-/low} (14.8 ± 3.1%, n = 5) (**Figure 1C**). In comparison to regular blood CD34⁺CD45RA⁻CD90⁺ Flt3^{-/low}CD7⁺CD71⁺ HSC (0.49 ± 0.19%, n = 4) from non-mobilized healthy donors, the miCD34⁺ cells expressed similar surface markers as CD34⁺CD45RA⁻CD90⁺ Flt3^{-/low}CD7⁺CD71⁺ (15.3 ± 2.9%, n = 5, *P* < 0.01), but higher levels of CD10 (a marker defining human lymphoid progenitors [23]) (99.4 ± 0.36% versus 20.6 ± 3.1%, *P* < 0.01), CD49f (a common biomarker for most populations of stem cells [24]) (98.8 ± 1.3% versus 15.4 ± 2.9%, *P* < 0.01), and lower level of BAH-1 (a marker for human megakaryocyte-erythroid progenitor [22]) (0.51 ± 0.2% versus 32.5 ± 3.9%, *P* < 0.01) (**Figure 1 C and D**). Due to coexpressions of CD7 and CD10 (the surface markers for common lymphoid progenitor (CLP) cells [25]) on miCD34⁺ HSC, the data suggested that miCD34⁺ HSC might have a high potential to give rise to lymphocytes.

Figure 1: Differentiation of PB-IPC into CD34⁺ HSC-like cells after the treatment with platelet-derived mitochondria. **(A)** The purity analysis of isolated mitochondria. The different markers have been applied by flow cytometry including MitoTrack Deep Red staining, anti-cytochrome C and anti-heat shock protein (HSP) 60 Abs for mitochondrial markers, calnexin for endoplasmic reticulum (ER), and GM130 for Golgi apparatus. Isotype-matched IgGs (grey histogram) served as negative controls (n = 3). **(B)** CD34 expression upregulation after treatment with mitochondria in miPB-IPC. Data represent mean ± SD of five experiments. **(C)** Phenotypic characterization of gated miCD34⁺ HSC (red) in total miPB-IPC. Isotype-matched IgGs served as controls. Data were representative from five preparations. **(D)** Phenotypic characterization of gated CD34⁺CD45RA⁻ HSC (bottom, red) CD34⁺CD45RA⁺ cell population (top, blue) in total PBMC (n = 4). Isotype-matched IgGs served as controls. Data were representative from one of four preparations.

Differentiation of miCD34⁺ HSC into T Cells

To determine whether miCD34⁺ cells were functional as stem cells, they were purified from miPB-IPC and treated with

different inducers (**Figure 2A**). We first examined their potential to differentiate into T cells by treating purified miCD34⁺ cells with recombinant FMS-like tyrosine kinase (FLT)-3 ligand, interleukin (IL)-2, and IL-7 for 3 days. Phase-contrast microscopy revealed marked morphological changes and the differentiated T cells had numbers of cell clusters in this cytokine-treated group, with some cells released into the supernatant (**Figure 2B, right**). Cells in the control groups exhibited a smooth surface and failed to show any morphological changes (**Figure 2B, left, and 2C, left**). Confocal microscopy demonstrated the differentiated cells strongly expressed human T cell marker CD4, with weak expression of CD8 (**Figure 2D**). Flow cytometry further confirmed the differentiation of miCD34⁺ HSC into CD3⁺CD4⁺CD8⁻CD38⁺ T cells at a percentage of $76.93 \pm 3.21\%$ (**Figure 2E**, $n = 4$), which were CD3⁺CD4⁺TCR $\alpha\beta$ ⁺ T cells ($82.65 \pm 5.2\%$, $n = 3$) (**Figure 2F**). Intracellular staining with T cell functional markers indicated that these T cells produced Th1 cytokine IL-12 ($65.3 \pm 20.1\%$, $n = 3$) and Th2 cytokines IL-4 ($28.5 \pm 9.99\%$, $n = 3$) and IL-5 ($53.9 \pm 11.2\%$, $n = 3$), with a very low level of interferon (INF)- γ ($0.61 \pm 0.3\%$, $n = 3$) (**Figure 2G**). Additional functional tests established the significantly up-regulated expressions of cytokines such as IL-4 ($P = 0.0025$, $n = 3$), IL-5 ($P = 0.0049$, $n = 3$), and IL-12 ($P = 0.037$, $n = 3$) after the treatment with PMA and ionomycin. The level of INF- γ failed to show a marked change ($P = 0.085$, $n = 3$). The data confirmed that the differentiated T cells were responding to the stimulation of PMC/ionomycin (**Figure 2H**). The rapid differentiation of T cells with high efficiency clearly demonstrated that miCD34⁺ HSC were converted into functional and definitive hematopoietic progenitors.

Figure 2: In vitro differentiation of purified miCD34⁺ HSC to T cells. (A) Outline of the whole protocol from the generation of miPB-IPC to multipotent cellular differentiation of miPB-IPC. The miCD34⁺ HSC were purified for *in vitro* and *in vivo* differentiations respectively. (B) Phase-contrast images showing T cell differentiation of purified miCD34⁺ HSC in the presence of FLT-3 ligand, IL-2, and IL-7 for 3 days (n = 4). Untreated cells served as a control (left). The treated-CD34⁺ HSC displayed substantially morphological changes with cell clusters (orange stars), and some floating cells (red arrows) released from “cell clusters” (orange stars). Magnification ×200. (C) Phase-contrast images show the morphology of control PB-IPC (left) and treated PB-IPC in the presence of Mitochondria (right) (n = 4). Original magnification: ×200. (D) Z-stacked confocal images demonstrated strong expression of human T cell marker CD4 with a low expression of CD8 (n = 4). Untreated miCD34⁺ HSC served as control for immunostaining (left panel). (E) Flow cytometry revealed that differentiated T cells were CD3⁺CD4⁺CD8⁺CD38⁺ (n = 4). Untreated miCD34⁺ HSC served as controls (top panel). (F) Expression of T cell receptors α/β (TCRαβ) in the gated CD3⁺ and CD4⁺ T cells (n = 4). (G) Intracellular

staining of differentiated T cells with Th1/Th2 cell cytokine markers ($n = 3$). Isotype-matched IgG served as a control. Data are presented as mean $\pm$ SD from three experiments. **(H)** Fold changes of cytokine expression levels in the in vitro differentiated T cells post the stimulation with PMA and ionomycin ($n = 3$). Data represent mean $\pm$ SD.

Ex vivo Differentiation of miCD34⁺ HSC into other Hematopoietic Lineages

To further examine their potential for differentiation, we treated miCD34⁺ HSC with macrophage colony-stimulating factor (M-CSF) for 3 days, whereupon they became adherent, well-spaced cells. Functional analysis established that M-CSF-treated miCD34⁺ HSC exhibited strong phagocytosis of fluorescent latex beads (**Figure 3A, middle**), while untreated cells were mostly negative for this effect (**Figure 3A, left**). Flow cytometry confirmed that $34.3 \pm 4.3\%$ of the cells were CD11b⁺CD209⁺ macrophages (MΦ) and about $53.66 \pm 3.8\%$ were CD11b⁺CD209⁻ macrophages (MΦ) (**Figure 3A, right**). In contrast, there were only $15.29 \pm 1.5\%$ of CD11b⁺CD209⁻ macrophages and $0.43 \pm 0.12\%$ of CD11b⁺CD209⁺ macrophages in untreated miCD34⁺ HSC.

Figure 3: Ex vivo multiple differentiation of miCD34⁺ HSC. **(A)** Differentiation of miCD34⁺ HSC into macrophages after treatment with 50 ng/mL M-CSF for 3 days. The differentiated M ϕ displayed macrophage markers such as CD11b and CD209, and exhibited phagocytosis of fluorescence beads (green) (n = 4). Untreated miCD34⁺ HSC served as controls. **(B)** Differentiation of miCD34⁺ HSC into granulocytes after treatment with 100 ng/mL G-CSF + 25 ng/mL FLT-3L for 3 days, followed by Wright-Giemsa staining (left), and flow cytometry for granulocyte marker CD66b (n = 4). Untreated miCD34⁺ HSC served as controls. **(C)** Differentiation of miCD34⁺ HSC into erythrocytes shown by phase contrast imaging of mature RBC (indicated by red arrow), and by Wright-Giemsa staining with typical morphology of mature RBC (pointed by red arrows) (n = 4). **(D)** Analysis of the percentage of the matured CD45⁺ hemoglobin⁺ RBC in the gated CD235a⁺ cells by flow cytometry (n = 4). **(E)** Differentiation of miCD34⁺ HSC into megakaryocytes (MK)/platelets after the treatment with FLT-3L + TPO for 7 days, exhibiting CD42⁺ and a polynuclear appearance (n = 4). **(F)** Analysis of polyploidy MK post-treatment with FLT-3L + TPO for 7 days, shown by histogram (blue, middle) and dot plot with

MK/platelets' marker CD42a (top). Normal platelets (dark blue) and T cells (grey) from healthy donors served as controls for cells with no nucleus (0N) and one nucleus (1N) respectively (bottom) ($n = 4$). (G) Wright-Giemsa staining showed the differentiated miCD34⁺ HSC with multiple nuclei (pointed by red arrows) after the treatment with FLT-3L + TPO for 7 days ($n = 4$).

Next, we induced miCD34⁺ HSC with granulocyte colony-stimulating factor (G-CSF). After 3 days, $81.14 \pm 3.7\%$ of treated cells displayed the granulocyte-specific marker CD66b, with a reduced nuclear-cytoplasmic ratio and multi-lobed nuclei shown by Wright-Giemsa staining (**Figure 3B**). However, the untreated miCD34⁺ HSC failed to express CD66b and displayed a big size of nuclei, with a large of nuclear-cytoplasmic ratio (**Figure 3B**, $n = 4$). Moreover, after being treated with erythropoietin (EPO) for 5 days, miCD34⁺ HSC turned into nucleated cells strongly positive for the erythroid (Er) lineage marker CD235a and facilitated RBC maturation via expulsion of their nuclei with additional EPO treatment, exhibiting a distinctive biconcave shape and enucleated RBC (**Figure 3C, right**). In total, $41.4 \pm 11.46\%$ of cells were terminally-differentiated into enucleated CD235a⁺CD45⁻hemoglobin⁺ RBCs (**Figure 3D**, $n = 4$). However, untreated cells failed to show these differentiations or only expressed background levels of these markers. Further flow cytometry analysis demonstrated that the level of hemoglobin expression was increased in the matured RBCs, with the mean fluorescence intensity of hemoglobin⁺CD45⁻ mature RBCs at $13.61 \pm 4.29\%$, while hemoglobin⁺CD45⁺ immature RBCs was $8.29 \pm 1.61\%$ ($P = 0.044$, $n = 4$).

Additionally, we examined the commitment of miCD34⁺ HSC to MKs and platelets, which are critical for blood clotting. After treatment with FLT-3 ligand and thrombopoietin (TPO) for 7 days, production of CD42⁺ MKs was achieved with typical polyploidization (mostly from 2N to 7N) (**Figure 3 E-G**) and the formation of non-nucleated CD42⁺ platelets ($21.3 \pm 4.1\%$, $n = 4$) (**Figure 3E, F**), yielding of 95 ± 17 platelets per MK. Approximately 54% of mature CD42⁺ platelets were released into the supernatant (**Figure 3F**, $n = 4$).

In vivo Differentiation of miCD34⁺ HSC into other Hematopoietic Lineages after Transplant into NSG Mice

To further demonstrate their multipotent features, the purified miCD34⁺ HSC were transplanted into irradiated NOD/Lt-*scid*/IL2R γ ^{null} (NSG) mice (**Figure S1**). We examined the chimaerism of human CD45⁺ cells in peripheral blood, spleen and bone marrow of miCD34 HSC-engrafted mice at 12 weeks, by using flow cytometry analysis with blood cell lineage-specific markers [11,26] including T cells (CD3⁺CD4⁺), B cells (CD19⁺), monocytes (CD14⁺), granulocytes (CD66b⁺), erythroid cells (CD235a⁺), and megakaryocytes/platelets (CD41b⁺). We found that the engraftment levels of human CD45⁺ cells in blood ($9.93 \pm 9.62\%$, $P = 0.035$) and spleen ($25.37 \pm 21.89\%$, $P = 0.018$) were much higher than that in bone marrow ($0.33 \pm 0.15\%$) post transplantation at 12 weeks (**Figure 4A**, $n = 6$ mice). The miCD34⁺ HSC-derived CD3⁺CD4⁺ T cells remained as a predominant population at $57.92 \pm 8.49\%$ of human CD45⁺ blood cells after transplantation for 12 weeks, with different proportions of other engrafted cells in the blood. Similar data ($59 \pm 13.55\%$ of CD3⁺CD4⁺ T cells) were obtained from splenocytes of miCD34⁺ HSC-engrafted mice (**Figure 4 B and C**, $n = 5$ mice). By comparison, the percentage of miCD34⁺ HSC-derived CD3⁺CD4⁺ T cells in bone marrow were $49.45 \pm 14.01\%$ at 12 weeks, CD19⁺ B cells were $5.24 \pm 2.68\%$, CD41b⁺ megakaryocytes/platelets were $4.3 \pm 2.0\%$, with CD14⁺ monocytes at $1.71 \pm 2.36\%$, CD66b⁺ granulocytes at $0.51 \pm 0.46\%$, and CD235a⁺SYTO60⁺ nucleated erythroid cells at $0.22 \pm 0.13\%$ and CD235a⁺SYTO60⁻ enucleated erythroid cells at $0.08 \pm 0.05\%$ (**Figure 4D**, $n = 6$ mice). The percentage of CD14⁺ monocytes was $2.1 \pm 1.81\%$ in peripheral blood of miCD34⁺ HSC-transplanted mice. Additional flow cytometry failed to detect the primary phenotype of undifferentiated miCD34⁺HSC. There was little to no human CD34⁺ cells in peripheral blood ($0.07 \pm 0.04\%$), spleen ($0.04 \pm 0.03\%$) and bone marrow ($0.01 \pm 0.01\%$) of miCD34 HSC-engrafted mice at 12 weeks (**Figure 4E**, $n = 5$ mice).

Figure 4: *In vivo* multiple differentiation of miCD34⁺ HSC after transplant into the irradiated NSG mice. To determine the multilineage differentiation of miCD34⁺ HSC after transplantation into irradiated NSG mice, only the viable cells from different samples were gated for analysis after excluding the propidium iodide (PI)-positive dead cells. The gated human leukocyte common antigen CD45-positive and mouse CD45.1-negative viable cells were analyzed for characterization with different lineage-specific surface markers such as CD3 and CD4 for T cells, CD19 for B cells, CD41b for megakaryocytes/platelets, CD14, CD11b and CD11c for monocyte/macrophage, CD66b for granulocytes, and CD235a for erythroid cells. SYTO60 was utilized to stain the CD235a⁺ nucleated erythroid cells. Isotype-matched IgGs served as controls for flow cytometry. (A) Engrafted levels of hCD45⁺mCD45.1⁺ cells in peripheral blood, spleen, and bone marrow of miCD34⁺ HSC-transplanted NSG mice at 12 weeks (3×10^5 cells/mouse in 200 μ l physiological saline, i.v., n = 6). (B) Phenotypic characterization of miCD34⁺ HSC post engraftment in the irradiated NSG

mice (3×10^5 cells/mouse in 200 μ l physiological saline, i.v., $n = 5$). Tissue samples were collected post the transplantation for 12 weeks. Only the viable cells (green histogram) were gated for analysis after excluding the propidium iodide (PI)-positive dead cells. The gated human leukocyte common antigen CD45-positive and mouse CD45.1-negative viable cells were analyzed. Representative data are from one of three experiments with similar results. Isotype-matched IgGs served as controls for flow cytometry. **(C)** Multi-lineage differentiation of miCD34⁺ HSC after transplantation into irradiated NSG mice for 12 weeks. **(D)** Erythroid reconstitution of miCD34⁺-transplanted NSG mice at 12 weeks ($n = 6$). Only the viable cells (green histogram) were gated for analysis after excluding the propidium iodide (PI)-positive dead cells. SYTO60 was utilized to stain the CD235a⁺ nucleated erythroid cells [11]. **(E)** Characterization of miCD34⁺ HSC post engraftment for 12 weeks in the irradiated NSG mice ($n = 5$). Representative data are from one of three experiments with similar results. The gated human CD45-positive and mouse CD45.1-negative viable cells were analyzed. Isotype-matched IgGs served as controls for flow cytometry. **(F)** Myeloid differentiation of miCD34⁺ HSC after transplant in NSG mice at 12 weeks (top) and 16 weeks (bottom, $n = 3$).

To further confirm miCD34⁺HSC giving rise to monocytes/macrophages (myeloid lineage differentiation), we performed an additional animal study in miCD34⁺HSC-transplanted mice at 12 and 16 weeks respectively by using macrophage-associated markers anti-human CD11b and CD11c mAbs. Flow cytometry demonstrated that the percentage of mCD45⁺hCD45⁺hCD3⁺hCD11b⁺hCD11c⁺ macrophages was $75.22 \pm 18.33\%$ in the splenocytes of miCD34⁺ HSC-transplanted mice at 16 weeks (**Figure 4F**, $n = 3$). In contrast, the percentage of mCD45⁺hCD45⁺hCD3⁺hCD11b⁺ T cells was declined from $59 \pm 14.01\%$ at 12 weeks to $4.05 \pm 2.87\%$ at 16 weeks (**Figure 4F**). Thus, the data clearly demonstrated the monocyte/macrophage (myeloid) differentiation of miD34⁺ HSC after transplantation into the irradiated NSG mice. Considering other lineage differentiations (e.g., T cells, B cells, megakaryocytes and red blood cells), these data indicated the multi-lineage differentiations of miCD34⁺ HSC.

Notch Signaling Pathway Contributes to the miCD34⁺ HSC Differentiation after the Treatment with Platelet-Derived Mitochondria

Notch signaling has been well established as an essential regulator for HSC generation and differentiation [27]. Specifically, Notch signaling pathway plays a crucial role in the T-cell development and maturation at different stages [28]. Both *ex vivo* and *in vivo* data demonstrated the multiple differentiations of miCD34⁺ HSC. To dissect their molecular mechanisms underlying mitochondrial treatment, we explored the action of Notch signaling during the induction of differentiation of PB-IPC toward the miCD34⁺ HSC. Flow cytometry revealed that mitochondria expressed Notch ligands: Jagged 1 (JAG1) ($25.13 \pm 16.0\%$), Jagged 2 (JAG2) ($68.04 \pm 14.6\%$) and Delta-like 3 (DLL3) ($69.3 \pm 25.96\%$); DLL1 ($2.21 \pm 1.74\%$) and DLL4 ($0.23 \pm 0.09\%$) (**Figure 5A**) were not expressed. The expression of Notch receptors 1 – 4 on PB-IPC were markedly upregulated after the treatment with platelet-derived mitochondria (**Figure 5B**). To examine the role of Notch signaling in miCD34⁺ HSC differentiation of mitochondrion-induced PB-IPC, N-[N-(3,5-difluorophenacetyl)-L-alanyl]-S-phenylglycine t-butyl ester (DAPT) treatment was used to block γ -secretase (**Figure 5C**), an enzyme critical for the release of the Notch intracellular domain (NICD) into the nucleus to initiate gene transcription [29]. The percentage of CD34⁺ cells was significantly increased in the group treated with mitochondria + DAPT ($26.2 \pm 5.68\%$) (**Figure 5D**). Contrastingly, treatment with DAPT alone showed a very low ability to induce CD34⁺ cells ($2.59 \pm 0.13\%$), indicating mitochondria is required for miCD34⁺ HSC cell differentiation.

Figure 5: Notch pathway in the differentiation of PB-IPC toward miCD34⁺ HSC. **(A)** Analysis of Notch ligands on platelet-derived mitochondria by flow cytometry (n = 3). **(B)** Expression of Notch receptors on PB-IPC by flow cytometry. PB-IPC were treated with mitochondria for 7 days and collected for flow cytometry. Untreated PB-IPC served as control. n = 3. **(C)** Phase-contrast images show the morphology of PB-IPC (left) and treated PB-IPC in the presence of mitochondria (middle) and mitochondria + DAPT (right). Original magnification: $\times 200$. **(D)** CD34 expression upregulation after the treatment with mitochondria and/or DAPT. DAPT-treated PB-IPC served as control. n = 4. Data represent mean $\pm$ SD.

Discussion

The use of autologous stem cells for regenerative medicine is more ethically acceptable and likely more successful than the use of other stem cells, and such therapies would avoid many of the immune rejection and safety concerns associated with other stem cells (e.g., ES- or iPS-based therapies). PB-IPC can be easily isolated from peripheral blood and expanded in serum-free culture medium to avoid the painful and invasive procedures required to withdraw bone marrow. Using autologous PB-IPC from patients as a starting material, mitochondrial treatment can generate functional autologous miCD34⁺ HSC on a large scale, giving rise to different blood cell lineages such as T cells, M Φ , granulocytes, erythrocytes, and MKs/platelets. Current data indicated that miCD34⁺ HSC exhibited a rapid multiple potential for differentiation post-graftment in the irradiated NSG mice.

Thus, these cells offer great promise as a solution for the current bottlenecks associated with conventional stem cell transplants and have tremendous potential for patient benefit in the clinic. Generation of functional autologous miCD34⁺ HSC from PB-IPC may provide unmet medical need.

Both *ex vivo* and *in vivo* studies have demonstrated the T-cell differentiation of miCD34⁺ HSC at high efficiency. To exclude the possibility of the expansion of contaminated cells, we have carefully designed experiments and analyzed the differentiated cells through morphology and flow cytometry with multiple different T-cell specific markers such as CD3, CD4, CD8, TCR α/β , TCR γ/δ , Th1 and Th2 cytokines. There were a few CD8⁺ T cells adhered to PB-IPC after overnight (12 hrs) culture in non-tissue culture-treated Petri dishes, but no CD4⁺ T cells. After an expansion of PB-IPC for 7 days, PB-IPC were attached to the bottom of the Petri dish with high purity (> 97%) and remained negative for CD4, with a few CD8 T cells (~ 2.63%). Before the treatment with platelet-derived mitochondria, all floating cells and cellular debris were washed away with PBS twice. The purified miCD34⁺ HSC cells were utilized for transplantation. Thus, the possibility of transplant mature PBMC and/or lymphocytes was very limited in our protocol. Normally, NSG mice fail to support human T-cell growth and differentiation. A few CD8⁺ T cells might be temporarily expanded after transplant in mice, however, flow cytometry demonstrated that most of T cells were CD3⁺CD4⁺ T cells in the miCD34⁺ HSC-transplanted NSG mice. Therefore, the engrafted CD3⁺CD4⁺ T cells were differentiated from miCD34⁺ HSC. Additionally, our *ex vivo* studies demonstrated the high efficiency of CD3⁺CD4⁺CD8⁻CD38⁺ T-cell differentiation from miCD34⁺ HSC, with an expression of Th1- and Th2-associated cytokines (IL-4, IL-5, and IL-12). To confirm the CD4 expression on T cells, not on monocytes/macrophages, T cell-specific marker CD3 was utilized for cellular gating during the flow cytometry. Therefore, the CD4⁺ T cells were derived from the differentiation of miCD34⁺ HSC.

Notch receptors act as key regulators in both CD34⁺ HSC and T-cell development and maturation at different stages [27,28].

After binding to its specific ligands, the Notch intracellular domain (NICD) is cleaved off by the enzyme γ -secretase and translocated to the nucleus, where it binds to the transcription factor recombination signal sequence-binding protein Jkappa (RBP-J), leading to the gene regulation. To understand the mechanism underlying miCD34⁺ HSC- and T-cell differentiations, current study revealed the upregulation of Notch receptors 1 – 4 on miPB-IPC, while platelet-derived mitochondria expressed Notch ligands Jagged 1 (JAG1), JAG3 and Delta-like 3 (DLL3), but not DLL1 and DLL4, highlighting the Notch signaling pathway may contribute to the miCD34⁺ HSC and T-cell differentiations after the treatment with platelet-derived mitochondria. Notably, the percentage of miCD34⁺ HSC was significantly upregulated after the mitochondrial treatment with addition of the γ -secretase inhibitor DAPT. The data confirmed that the canonical Notch pathway contributed to the differentiation of miCD34⁺ HSC through the interaction of Notch receptors on PB-IPC and their ligands on platelet-derived mitochondria. Due to the action of Notch pathway in the maintenance of quiescent feature of hematopoietic stem and progenitor cells [30], blocking with DAPT promote the miCD34⁺ HSC differentiation of PB-IPC, which is consistent with previous report [30]. Additional detailed molecular mechanisms need to be explored to better understand the individual and synergistic effects during the interaction of mitochondrial Notch ligands (JAG1, JAG3, and DLL3) with Notch receptors 1-4 on PB-IPC.

The roadmap of cell differentiation and maturation is tightly modulated through the systematic integration of distinct activating or repressing signaling pathways located both in the cytoplasm and nucleus, specifically for the hematopoietic system underlying the hierarchical differentiation model [22,26,31]. Our previous work [18] demonstrated that the highly-purified adult peripheral blood (PB)-derived platelets (> 99% purity) strongly display the ES cell-associated pluripotent gene markers such as transcription factors OCT3/4 and SOX2, with little or no expression of NANOG. Real time PCR Array revealed the expressions of human stem cell-related transcription factors and human stem cell-associated markers in the mitochondria of both

human PB-platelets [18], highlighting that the stemness' markers are localized in platelets' mitochondria, which may contribute to the induction of multipotency of PB-IPC after the treatment with PB-derived mitochondria. Our recent mechanistic studies confirmed that mitochondria enter cells and directly penetrate the nucleus of PB-IPC after the treatment with platelet-derived mitochondria, where they can produce profound epigenetic changes [21]. Due to the essential role of mitochondria in the reprogramming of somatic cells to iPS cells [32], additional molecular mechanisms underlying the mitochondrial reprogramming of PB-IPC need to be determined. In conclusion, innovative reprogramming of adult PB-IPC by the treatment with platelet-derived mitochondria may overcome the limitations and safety concerns by using conventional transgenic technologies to reprogram ES and iPS cells in the clinic.

Materials and Methods

PB-IPC cell culture

Human buffy coat blood units (N = 51; mean age of 48.97 ± 14.11 ; age range from 18 to 72 years old; 24 males and 27 females) were purchased from the New York Blood Center (New York, NY, <http://nybloodcenter.org/>). Human buffy coats were initially added 40 ml chemical-defined serum-free culture X-VIVO 15TM medium (Lonza, Walkersville, MD) and mixed thoroughly with 10ml pipette, and then used for isolation of peripheral blood-derived mononuclear cells (PBMCs). PBMCs were harvested as previously described [33]. Briefly, mononuclear cells were isolated from buffy coats blood using Ficoll-PaqueTM PLUS ($\gamma=1.007$, GE Healthcare), followed by removing the red blood cells using Red Blood Cell Lysis buffer (eBioscience, San Diego, CA). After three washes with saline, the whole PBMC were seeded in $150 \times 15\text{mm}$ Petri dishes (BD Falcon, NC) at 1×10^6 cells/ml, 25ml/dish in chemical-defined serum-free culture X-VIVO 15TM medium (Lonza, Walkersville, MD) without adding any other growth factors and incubated at 37°C in 8% CO_2 [34]. Seven days later, PB-IPC were growing and expanded by adhering to the hydrophobic bottom of Petri dishes. Consequently, PB-IPC were washed three times with saline and all floating cells were removed. Next, the serum-free

NutriStem[®] hPSC XF culture medium (Corning) was added for continue cell culture and expansion at 37 °C in 8% CO₂. The expanded PB-IPC were usually applied for experiments during 7-14 days.

Isolation of Mitochondria From Platelets

The mitochondria were isolated from PB-platelets using the Mitochondria Isolation kit (Thermo scientific, Rockford, USA, Prod: 89874) according to the manufacturer's recommended protocol [18]. Adult human platelet units (N = 16; mean age of 30.81 ± 8.64; age range from 16 to 40 years old; 9 males and 7 females) were purchased from the New York Blood Center (New York, NY, <http://nybloodcenter.org/>). The concentration of mitochondria was determined by the measurement of protein concentration using a NanoDrop 2000 Spectrophotometer (ThermoFisher Scientific, Waltham, MA). The isolated mitochondria were aliquoted and kept in -80 °C freezer for experiments.

For mitochondrial staining with fluorescent dyes, mitochondria were labeled with MitoTracker Deep Red FM (100 nM) (Thermo Fisher Scientific, Waltham, MA) at 37°C for 15 mins according to the manufacturer's recommended protocol, followed by two washes with PBS at 3000 rpm × 15 mins [18].

In vitro differentiation of PB-IPC into miCD34⁺ HSC

PB-IPC were treated with 100 µg/ml platelet-derived mitochondria for 7 – 14 days in the non-treated 24-well plates or Petri dishes with the serum-free NutriStem[®] hPSC XF culture medium (Corning), at 37 °C and 8% CO₂. According to our current protocol, the miCD34⁺ HSC cells were purified from mitochondria-treated PB-IPC by immunomagnetic sorting with Miltenyi Biotech CD34 MicroBead Kit (Miltenyi Biotech, catalog #130-097-047) according to the manufacturer's instructions. The differentiation of miCD34⁺ HSC were characterized by flow cytometry.

Flow Cytometry

Flow cytometric analyses of surface and intra-cellular markers were performed as previously described [18]. PB-IPC were washed with PBS at 2000 rpm for 5 mins. Mitochondria were washed with PBS at 12,000g for 10 mins at 4 °C. Samples were pre-incubated with human BD Fc Block (BD Pharmingen) for 15 mins at room temperature, and then directly aliquoted for different antibody staining. Cells were incubated with different mouse anti-human monoclonal antibodies (mAb). For surface staining, cells were stained for 30 min at room temperature and then washed with PBS at 2000 rpm for 5 mins prior to flow analysis. Isotype-matched mouse anti-human IgG antibodies (Beckman Coulter) served as a negative control for all fluorescein-conjugated IgG mAb. SYTOTM60 (Thermo Fisher) was combined with CD235a (GLY-A) staining to determine the nucleated erythroid cells. Staining with propidium iodide (PI) (BD Biosciences) was used to exclude dead cells during the flow cytometry analysis. For intra-cellular staining, cells were fixed and permeabilized according to the PerFix-nc kit (Beckman Coulter) manufacturer's recommended protocol. After staining, cells were collected and analyzed using a Gallios Flow Cytometer (Beckman Coulter) equipped with three lasers (488 nm blue, 638 red, and 405 violet lasers) for the concurrent reading of up to 10 colors. The final data were analyzed using the Kaluza Flow Cytometry Analysis Software (Beckman Coulter).

Cells were incubated with different mouse anti-human monoclonal antibodies (mAb) from Beckman Coulter (Brea, CA) including FITC-conjugated anti-CD45RA, anti-IFN γ , anti-CD4, anti-CD235a, anti-CD8 and anti-CD42a; phycoerythrin (PE)-conjugated anti-CD34; PE-Texas Red-conjugated CD3; phycoerythrin-Cy5 (PE-Cy5)-conjugated anti-CD90; phycoerythrin-Cy5.5 (PE-Cy5.5)-conjugated anti-CD19; phycoerythrin-Cy7 (PE-Cy7)-conjugated anti-CD49f, anti-CD11b and anti-CD45; APC-conjugated anti-CD4; APC-Alexa Fluor 700-conjugated anti-CD71; APC-Alexa Fluor 750-conjugated anti-CD7, anti-CD66b and anti-CD8; pacific blue (PB)-conjugated anti-CD38; Krome Orange-conjugated anti-

CD14 and anti-135 (FLT3) -BV510. From BD Biosciences (San Jose, CA), the investigator purchased the AlexaFluor-488-conjugated anti-human Cytochrome C; FITC-conjugated anti-CD90 (THY1) and anti-CD11C; BV 510-conjugated anti-CD45, PE-conjugated anti-IL4, anti-IL5, anti-BAH1, anti-IL12; PE-CF594-conjugated anti-CD10, BV421-conjugated anti-CD209 and PE-conjugated anti-mouse CD45.1. Antibodies were purchased from Biolegend (San Diego, CA) including the FITC-conjugated anti-human Hsp60, anti-human TCR $\alpha\beta$, anti-human Notch 1 and anti-human Notch 2; PE-conjugated anti-human Notch 3, anti-human Dll1, anti-human Dll4 and anti-human Jagged2; APC conjugated anti-human Notch 4 ; phycoerythrin-Cy7 (PE-Cy7)-conjugated anti-TCR $\gamma\delta$ and pacific blue (PB)-conjugated anti-CD3. FITC-conjugated anti-human Jagged1 and PE-conjugated anti-human Dll3 mAbs were purchased from R&D Systems (Minneapolis, MN). Hemoglobin $\beta/\gamma/\delta$ (H-76) rabbit polyclonal antibody, AlexaFluor-546-conjugated Calnexin and AlexaFluor-647-conjugated GM130 were purchased from Santa Cruz Biotechnology (Dallas, TX). The SYTOTM60 was purchased from Thermo Fisher.

Multiple differentiations of miCD34⁺ HSC

Initially, miCD34⁺ HSC were purified from miPB-IPC by using CD34 MicroBead Kit human-lyophilized (Miltenyi Biotec), through the auto MACS Pro Separator (Miltenyi Biotec) according to manufacturer's recommended protocol. The purified miCD34⁺ HSC were treated with different inducers for cellular differentiations.

To test T cell differentiation, the purified miCD34⁺ HSC (1×10^5 cells/ml) were planted in 24-well non-treated plates in the presence of HSC-Brew GMP Basal Medium (Miltenyi Biotec, Gladbach, Germany) with addition of cytokines 25 ng/mL hFlt3L and 25 ng/mL rhIL -7 (R&D Systems), at 37 °C in 5% CO₂. After the treatment for 3-7 days, cells were photographed and analyzed by confocal microscopy and flow cytometry with different T cell markers such as CD3, CD4, CD8, TCR α/β , CD38, Th1 cytokines (IL-4 and IL-5) and Th2 cytokines (IFN- γ and IL-12). Untreated miCD34⁺ HSC served as negative control.

T cells from healthy donors served as positive controls. For combined immunocytochemistry, the differentiated cells were fixed in 24-well plates with 4% paraformaldehyde for 20 mins and permeabilized with 0.5% triton X-100 for 5 mins, blocking non-specific binding with 2.5% horse serum, and followed by immunostaining with FITC-conjugated mouse anti-human CD4 and CD8. After covering with Mounting Medium with DAPI (Vector Laboratories), cells were photographed with Nikon A1R confocal microscope on Nikon Eclipse Ti2 inverted base, using software NIS Elements Version 4.60.

For the differentiation of miCD34⁺ HSC to macrophages, the purified miCD34⁺ HSC (1×10^5 cells/ml) were treated with 50 ng/mL M-CSF (Sigma) in 24-well non-treated plates in the presence of HSC-Brew GMP Basal Medium, at 37 °C in 5% CO₂. After the treatment for 2 - 3 days, cells were analyzed with phagocytosis and by flow cytometry with macrophage marker CD11b and CD209. Untreated miCD34⁺ HSC served as negative control. To detect the function of differentiated macrophages, the fluorescence latex beads (Sigma) were added to M-CSF-treated and untreated miCD34⁺ HSC cultures. After 4 hours of incubation with latex beads, cells were washed 3 times with PBS. The phagocytosis was viewed and evaluated under microscopy. The positive cells had minimum 5 beads per cell.

To differentiate the miCD34⁺ HSC into granulocytes, the purified miCD34⁺ HSC (1×10^5 cells/ml) were treated with 25 ng/mL hFlt3L + 100 ng/mL G-CSF (R&D Systems) in the presence of HSC-Brew GMP Basal Medium, in the 24-well non-treated plates, at 37 °C in 5% CO₂. After the treatment for 3-5 days, cells were photographed and analyzed by flow cytometry with granulocyte marker CD66b and staining with Wright-GIEMSA (Sigma) according to the manufacturer's instruction. Untreated miCD34⁺ HSC served as negative control. PBMC from healthy donors served as positive controls.

To differentiate miCD34⁺ HSC into the RBC, the purified miCD34⁺ HSC (1×10^5 cells/ml) were initially treated with 25 ng/mL hFlt3L + 3 units/mL EPO (R&D Systems) in the presence of HSC-Brew GMP Basal Medium, in the 24-well non-treated

plates, at 37 °C in 5% CO₂. After this treatment for 5 days, cells were re-treated with 3 units/mL EPO for additional 3 – 7 days. Consequently, cells were photographed and analyzed by flow cytometry with erythrocyte markers CD235a and hemoglobin. Untreated miCD34⁺ HSC served as negative control. For intracellular flow cytometry, all floating cells were collected and centrifuged at 2,700g × 15 mins. First, post blocking non-specific binding with Fc Blocker (BD Biosciences), cells were fixed and permeabilized using PerFix-nc kit (Beckman Coulter) according to the manufacturer's recommended protocol. Second, cells were incubated with rabbit anti-human hemoglobinβ/γ/δ polyclonal antibody (Santa Cruze) at 1 : 100 dilution, room temperature for 30mins, and then washed with PBS at 2700g × 15 mins. Next, cells were labeled with Cy5-conjugated AffiniPure donkey anti-rabbit 2nd Ab (Jackson ImmunoResearch Laboratories, West Grove, PA), in combination with staining with mouse anti-human CD235a-FITC (Beckman Coulter) and CD45-PE-CY7 mAbs for 30 mins, and followed by flow cytometry analysis.

To differentiate miCD34⁺ HSC into megakaryocytes and platelets, the purified miCD34⁺ HSC (1 × 10⁵ cells/ml) were initially treated with 25 ng/mL hFlt3L + 100 ng/mL TPO (R&D Systems) in the presence of HSC-Brew GMP Basal Medium, in the 24-well non-treated plates, at 37 °C in 5% CO₂. After this treatment for 3 - 7 days, cells were photographed and collected for flow cytometry with MK/platelets' marker CD42a. Untreated miCD34⁺ HSC served as negative control. For the analysis of polyploidization: Firstly, the viable TPO-treated miCD34⁺ cells were stained with CD42a mAb and Hoechst 33342 (Sigma) and photographed under confocal microscope. Secondly, using healthy donor-derived matured T cells (1N) and platelets (0N) as control, the polyploidy of differentiated CD42a⁺ MKs were analyzed by flow cytometry after staining with Propidium Iodide (PI) (Abcam, Cambridge MA) according to the manufacturer's recommended protocol.

To morphological determine the differentiation of miCD34⁺ HSC to granulocytes, RBCs, and megakaryocytes/platelets, Wright-Giemsa staining was performed on the treated and

untreated cells respectively, and observed and photographed under the inverted Nikon ECLIPSE Ti2 microscope.

For DAPT blocking experiment, PB-IPC were treated with 100 $\mu\text{g/ml}$ mitochondria + 10 μM DAPT (Sigma, Catalog# D5942) for 7 - 14 days in the non-tissue culture-treated 24-well plates or Petri dishes with the serum-free NutriStem® hPSC XF culture medium (Corning), at 37 °C and 8% CO₂ conditions. Consequently, both treated and untreated PB-IPC were collected to examine the expression of CD34 by flow cytometry.

Animal Study and Engraftment of miCD34⁺ HSC in irradiated NSG Mice

All animal experiments were performed according to approval of the institutional animal care and use committee of Hackensack Meridian Health. To demonstrate their multipotent features of miCD34⁺ HSC, the purified miCD34⁺ HSC were transplanted into irradiated NOD/Lt-*scid*/*IL2R γ* ^{null} (NSG) mice [35]. NSG mice were purchased from Jackson Laboratories (Bar Harbor, ME), breeding and maintaining under pathogen-free conditions at the animal facility of Center for Discovery and Innovation. To determine the repopulating potential of miCD34⁺ HSC, NSG mice, aged 12 – 16 weeks, were irradiated with 200 cGy by using RS-2000 irradiator. The miCD34⁺ HSC were transplanted to irradiated NSG mice at 3×10^5 cells/mouse via tail vein (in 200 μl saline, i.v., n = 10 mice) 24 hours after irradiation, according to a protocol approved by the Animal Care Committee (ACC) of Center for Discovery and Innovation. Only physiological saline injectin (200 μl) served as a control (n = 6 mice). After transplantation, mice were monitored twice a week for 16 weeks. To examine the differentiation of miCD34⁺ HSC, mice were sacrificed at 12th and 16th week respectively post-transplantation to collect samples of peripheral blood, spleen, and bone marrow for flow cytometry analysis. To determine the multilineage differentiation of miCD34⁺ HSC after transplantation into irradiated NSG mice, only the viable cells from different samples were gated for analysis after excluding the propidium iodide (PI)-positive dead cells. The gated human leukocyte common antigen CD45-positive and mouse CD45.1-negative viable cells were further analyzed for characterization

with different human blood cell lineage-specific surface markers such as CD3 and CD4 for T cells, CD19 for B cells, CD41b for megakaryocytes/platelets, CD14, CD11b and CD11c for monocyte/macrophage, CD66b for granulocytes, and CD235a for erythroid cells. SYTO60 was utilized to stain the CD235a⁺ nucleated erythroid cells. To determine T cell population and remove CD4⁺ monocytes, anti-CD3 Ab was employed for gating out CD4⁺ monocytes, in addition to the consideration of cell-size difference. Isotype-matched IgGs served as controls for flow cytometry.

Statistics

Statistical analyses were performed with GraphPad Prism 8 (version 8.0.1) software. The normality test of samples was performed by the Shapiro-Wilk test. Statistical analyses of data were performed by the two-tailed paired Student's t-test to determine statistical significance between untreated and treated groups. The Mann-Whitney U test was utilized for non-parametric data. Values were given as mean $\pm$ SD (standard deviation). Statistical significance was defined as $P < 0.05$, with two sided.

References

1. Doulatov S, Notta F, Laurenti E, Dick JE. Hematopoiesis: a human perspective. *Cell Stem Cell*. 2012; 10: 120-136.
2. Wahlster L, Daley GQ. Progress towards generation of human haematopoietic stem cells. *Nat Cell Biol*. 2016; 18: 1111-1117.
3. Swart JF, Delemarre EM, van Wijk F, Boelens JJ, Kuball J, et al. Haematopoietic stem cell transplantation for autoimmune diseases. *Nat Rev Rheumatol*. 2017; 13: 244-256.
4. Gambell P, Herbert K, Dickinson M, Stokes K, Bressel M, et al. Peripheral blood CD34⁺ cell enumeration as a predictor of apheresis yield: an analysis of more than 1,000 collections. *Biol. Blood Marrow Transplant*. 2012; 18: 763-772.

5. Chabannon C, Kuball J, Bondanza A, Dazzi F, Pedrazzoli P, et al. Hematopoietic stem cell transplantation in its 60s: A platform for cellular therapies. *Sci. Transl. Med.* 2018; 10.
6. Kanakry CG, Fuchs EJ, Luznik L. Modern approaches to HLA-haploidentical blood or marrow transplantation. *Nat Rev Clin Oncol.* 2016; 13: 132.
7. Woods NB, Parker AS, Moraghebi R, Lutz MK, Firth AL, et al. Brief report: efficient generation of hematopoietic precursors and progenitors from human pluripotent stem cell lines. *Stem Cells.* 2011; 29: 1158-1164.
8. Gori JL, Butler JM, Chan YY, Chandrasekaran D, Poulos MG, et al. Vascular niche promotes hematopoietic multipotent progenitor formation from pluripotent stem cells. *J. Clin. Invest.* 2015; 125: 1243-1254.
9. Blaser BW, Zon LI. Making HSCs in vitro: don't forget the hemogenic endothelium. *Blood.* 2018; 132: 1372-1378.
10. Okita K, Yamakawa T, Matsumura Y, Sato Y, Amano N, et al. An efficient nonviral method to generate integration-free human-induced pluripotent stem cells from cord blood and peripheral blood cells. *Stem Cells.* 2013; 31: 458-466.
11. Sugimura R, Jha DK, Han A, Soria-Valles C, da Rocha EL, et al. Haematopoietic stem and progenitor cells from human pluripotent stem cells. *Nature.* 2017; 545: 432-438.
12. Doulatov S, Vo LT, Chou SS, Kim PG, Arora N, et al. Induction of multipotential hematopoietic progenitors from human pluripotent stem cells via respecification of lineage-restricted precursors. *Cell Stem Cell.* 2013; 13: 459-470.
13. Staal FJ, Baum C, Cowan C, Dzierzak E, Hacein-Bey-Abina S, et al. Stem cell self-renewal: lessons from bone marrow, gut and iPS toward clinical applications. *Leukemia.* 2011; 25: 1095-1102.
14. Zhao Y, Mazzone T. Human cord blood stem cells and the journey to a cure for type 1 diabetes. *Autoimmun. Rev.* 2010; 10: 103-107.
15. Zhao Y, Jiang Z, Zhao T, Ye M, Hu C, et al. Reversal of type 1 diabetes via islet beta cell regeneration following immune modulation by cord blood-derived multipotent stem cells. *BMC Med.* 2012; 10: 3.
16. Zhao Y, Jiang Z, Zhao T, Ye M, Hu C, et al. Targeting insulin resistance in type 2 diabetes via immune modulation of cord

- blood-derived multipotent stem cells (CB-SCs) in stem cell educator therapy: phase I/II clinical trial. *BMC. Med.* 2013; 11: 160.
17. Delgado E, Perez-Basterrechea M, Suarez-Alvarez B, Zhou H, Revuelta EM, et al. Modulation of Autoimmune T-Cell Memory by Stem Cell Educator Therapy: Phase 1/2 Clinical Trial. *EBioMedicine.* 2015; 2: 2024-2036.
18. Zhao Y, Jiang Z, Delgado E, Li H, Zhou H, et al. Platelet-Derived Mitochondria Display Embryonic Stem Cell Markers and Improve Pancreatic Islet beta-cell Function in Humans. *Stem Cells Transl. Med.* 2017; 6: 1684-1697.
19. Li Y, Yan B, Wang H, Li H, Li Q, et al. Hair regrowth in alopecia areata patients following Stem Cell Educator therapy. *BMC. Med.* 2015; 13: 87.
20. Zhao Y, Huang Z, Lazzarini P, Wang Y, Di A, et al. A unique human blood-derived cell population displays high potential for producing insulin. *Biochem. Biophys. Res. Commun.* 2007; 360: 205-211.
21. Yu H, Hu W, Song X, Zhao Y. Generation of Multipotent Stem Cells from Adult Human Peripheral Blood Following the Treatment with Platelet-Derived Mitochondria. *Cells.* 2020; 9.
22. Notta F, Zandi S, Takayama N, Dobson S, Gan OI, et al. Distinct routes of lineage development reshape the human blood hierarchy across ontogeny. *Science.* 2016; 351: aab2116.
23. Calloni R, Cordero EA, Henriques JA, Bonatto D. Reviewing and updating the major molecular markers for stem cells. *Stem Cells Dev.* 2013; 22: 1455-1476.
24. Krebsbach PH, Villa-Diaz LG. The Role of Integrin alpha6 (CD49f) in Stem Cells: More than a Conserved Biomarker. *Stem Cells Dev.* 2017; 26: 1090-1099.
25. Hao QL, Zhu J, Price MA, Payne KJ, Barsky LW, et al. Identification of a novel, human multilymphoid progenitor in cord blood. *Blood.* 2001; 97: 3683-3690.
26. Yamamoto R, Morita Y, Ooehara J, Hamanaka S, Onodera M, et al. Clonal analysis unveils self-renewing lineage-restricted progenitors generated directly from hematopoietic stem cells. *Cell.* 2013; 154: 1112-1126.

27. Butko E, Pouget C, Traver D. Complex regulation of HSC emergence by the Notch signaling pathway. *Dev Biol.* 2016; 409: 129-138.
28. De Obaldia ME, Bhandoola A. Transcriptional regulation of innate and adaptive lymphocyte lineages. *Annual review of immunology.* 2015; 33: 607-642.
29. Weber D, Wiese C, Gessler M. Hey bHLH transcription factors. *Curr. Top. Dev. Biol.* 2014; 110: 285-315.
30. Anjos-Afonso F, Currie E, Palmer HG, Foster KE, Taussig DC, et al. CD34(-) cells at the apex of the human hematopoietic stem cell hierarchy have distinctive cellular and molecular signatures. *Cell Stem Cell.* 2013; 13: 161-174.
31. Akashi K, Traver D, Miyamoto T, Weissman IL. A clonogenic common myeloid progenitor that gives rise to all myeloid lineages. *Nature.* 2000; 404: 193-197.
32. Nishimura K, Fukuda A, Hisatake K. Mechanisms of the Metabolic Shift during Somatic Cell Reprogramming. *International journal of molecular sciences.* 2019; 20.
33. Zhao Y, Glesne D, Huberman E. A human peripheral blood monocyte-derived subset acts as pluripotent stem cells. *Proc. Natl. Acad. Sci. U. S. A* 2003; 100: 2426-2431.
34. Zhao Y, Wang H, Mazzone T. Identification of stem cells from human umbilical cord blood with embryonic and hematopoietic characteristics. *Exp. Cell Res.* 2006; 312: 2454-2464.
35. Shultz LD, Lyons BL, Burzenski LM, Gott B, Chen X, et al. Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R gamma null mice engrafted with mobilized human hemopoietic stem cells. *J. Immunol.* 2005; 174: 6477-6489.

Supplementary Materials

Figure S1: Scheme of the animal protocol for in vivo multiple differentiation of $miCD34^+$ HSC after transplantation into irradiated NSG mice. The purified $miCD34^+$ HSC (3×10^5 cells/mouse in 200 μ l physiological saline, i.v. injection via tail vein, $n = 10$) were transplanted into NSG mice exposed to 300 cGy irradiation using a RS-2000 x-ray irradiator. Mice injected with an equal volume of saline served as a control group ($n = 6$).