

Book Chapter

Aphid Behavior on *Amaranthus hybridus* L. (Amaranthaceae) Associated with *Ocimum* spp. (Lamiaceae) as Repellent Plants

Boni Barthélémy Yarou^{1,2*}, Aimé H Bokonon-Ganta³, François J Verheggen², Georges C Lognay^{2*} and Frédéric Francis²

¹Institut National des Recherches Agricoles du Bénin, Sous-Programme Cultures Maraîchères, Recette Principale, Cotonou, Godomey, Route IITA, Cotonou 01 BP: 884, Benin

²Functional and Evolutionary Entomology, TERRA Research Center, Gembloux Agro-bio Tech, University of Liege (ULg) Passage des Déportés, 2 BE-5030 Gembloux, Belgium

³Laboratoire d'Entomologie Agricole, Faculté des Sciences Agronomiques, Université d'Abomey-Calavi, Cotonou 03 BP: 2819, Benin

***Corresponding Authors:** Boni Barthélémy Yarou, Functional and Evolutionary Entomology, TERRA Research Center, Gembloux Agro-bio Tech, University of Liege (ULg) Passage des Déportés, 2 BE-5030 Gembloux, Belgium

Georges C Lognay, Functional and Evolutionary Entomology, TERRA Research Center, Gembloux Agro-bio Tech, University of Liege (ULg) Passage des Déportés, 2 BE-5030 Gembloux, Belgium

Published **December 30, 2020**

This Book Chapter is a republication of an article published by Boni Barthélémy Yarou, et al. at Agronomy in May 2020. (Yarou, B.B.; Bokonon-Ganta, A.H.; Verheggen, F.J.; Lognay, G.C.; Francis, F. Aphid Behavior on *Amaranthus hybridus* L. (Amaranthaceae) Associated with *Ocimum* spp. (Lamiaceae) as Repellent Plants. *Agronomy* 2020, 10, 736.)

How to cite this book chapter: Boni Barthélémy Yarou, Aimé H Bokonon-Ganta, François J Verheggen, Georges C Lognay, Frédéric Francis. Aphid Behavior on *Amaranthus hybridus* L. (Amaranthaceae) Associated with *Ocimum* spp. (Lamiaceae) as Repellent Plants. In: Isaac Danso, editor. Prime Archives in Agronomy. Hyderabad, India: Vide Leaf. 2020.

© The Author(s) 2020. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author Contributions: Conceptualization, B.B.Y.; methodology, B.B.Y.; software, B.B.Y.; validation, G.C.L., F.F.; formal analysis, B.B.Y.; investigation, B.B.Y.; data curation, B.B.Y.; writing—original draft preparation, B.B.Y.; writing—review and editing, A.H.B.-G., G.C.L., F.J.V., G.C.L. and F.F.; supervision, F.F. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Erasmus Mundus Program.

Acknowledgments: The authors would like to thank Antoine Boullis for his assistance in setting up the experimental system; Félicien Tosso for his assistance in statistical data analysis; Thomas Bawin, Hervé Kombiéni and Sognigbé N’Danikou for reading the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

Abstract

Various plant species contain biocidal and/or semiochemical components. These can be used for managing insect pests, in order to reduce the use of synthetic pesticides and to improve the quality of vegetable crops. This study was conducted to assess the effect of repellent plants *Ocimum gratissimum* L. and *Ocimum basilicum* L. on aphids *Aphis craccivora* Koch, *Aphis*

fabae Scopoli and *Myzus persicae* Sulzer when they are associated with *Amaranthus hybridus* L. plants. The results have shown that in the two approaches tested—*Ocimum* sp. plants surrounded by *A. hybridus* plants and the dual-choice test—the number of aphids on the *A. hybridus* plant associated with either *O. gratissimum* or *O. basilicum* was significantly less significant compared to the *A. hybridus* alone. This first study on the association between *A. hybridus* and *Ocimum* spp. shows that the *Ocimum* species might be used as an alternative method for controlling aphids in order to avoid the use of synthetic pesticides on *Amaranthus*. The ability of *Ocimum* spp. to repel pests can make it an important companion plant for farmers, because those plants can not only be used to control pests, but they can also be harvested, providing a direct economic return.

Keywords

Aphids; Basil; Pest Control; Amaranth; Repellent Activity; Companion Plant

Introduction

Aphids are pests that cause economically significant crop losses on many vegetable crops. To control these aphid populations, synthetic pesticides are increasingly used, especially in developing countries [1]. However, there is, nowadays, a growing public concern about the harmful effects of pesticides on humans and non-target organisms, in addition to the resistance of many pests to some commonly used insecticides [2,3]. Together, these problems have motivated the development of environmentally friendly alternative means to manage pests. Biological control using natural enemies is one of these alternatives to control crop pests. However, although this form of control strategy can potentially be successful at a low cost [4], it remains difficult to implement by small-scale farmers in developing countries. In this context, cheap, plant-based solutions can play an important role. Using plants with pesticidal properties to protect crops against pests can have advantageous effects for humans, such as reduced environmental degradation and increased food safety [5]. Indeed, botanical

extracts have been reported to be effective on pests and pathogens [6,7]. These plants can be used to protect crops by keeping pests away and then avoiding potential damage [8,9]. The biocidal activity of these plants is generally attributed to volatile organic compounds (VOC) that are emitted and can directly affect herbivorous pests due to their toxic, repellent or dissuasive properties [10,11].

Ocimum spp (Lamiaceae), or basils, are important plants with pesticidal effects, whose biocidal effect has been reported mainly on stored-product pests and malaria vectors [12–14]. Some studies indicated that their qualities as repellent or companion plants decreased pest abundance on crops. In a semi-field experiment, *O. americanum* L. essential oil had a repellent effect on *Agrotis ipsilon* Hufnagel (Lepidoptera: Noctuidae) [15]. Testing *O. basilicum* as a banker plant in greenhouse tomato crops led to fewer recorded pests [16]. Similarly, near to the tomato plants, *Ocimum* spp plants (*O. gratissimum* and *O. basilicum*) or their essential oils (through diffusers) reduced the oviposition of *Tuta absoluta* Meyrick (Lepidoptera: Gelechiidae) [17]. Intercropping of *O. basilicum* with *Gossypium barbadense* L. (Malvaceae) was also found to reduce pest abundance [18]. Introducing *O. basilicum* in *Vicia faba* L. (Fabaceae) reduced infestation levels of the bean aphid–*Aphis fabae* Scopoli (Hemiptera: Aphididae) [19]. A repellent effect was also reported on cabbage pests *Phyllotreta sinuata* Steph. (Coleoptera: Chrysomelidae), *Hellula undalis* Fabricius (Lepidoptera: Crambidea), *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae), *Spodoptera littoralis* Fabricius (Lepidoptera: Noctuidae) and *Plutella xylostella* (Lepidoptera: Plutellidae) when this crop was intercropped with *Ocimum* species [9,20]. *Ocimum basilicum* reduced pest abundance by 23% on *Abelmoschus esculentus* L. (Malvaceae) compared to the control, and also reduced the use of synthetic pesticides. In an orchard ecosystem, it was reported that planting *Ocimum* spp between trees can reduce pest levels and also attract natural enemies, including Coccinellidae, Syrphidae, Chrysopidae and Phytoseiidae [21–23].

The aim of the present study was to evaluate the repellent effect of *O. basilicum* and *O. gratissimum* on the aphids (Hemiptera: Aphididae) *Aphis craccivora* Koch (Cowpea aphid), *A. fabae* Scop and *Myzus persicae* Sulzer (Green peach aphid). The experiments were carried out on *Amaranthus hybridus* L. (Amaranthaceae), one of the most popular vegetables crops in tropical Africa [24] and highly attacked by aphids [25–27]. The *Ocimum* species are also used as vegetables in several West African countries [28]. The outcome of these experiments shows that *Ocimum* could be used as a companion plant for pest management.

Materials and Methods

Plants and Aphids Rearing

Seeds of *Amaranthus hybridus* (Ah), *O. gratissimum* (Og) and *O. basilicum* (Ob) were provided by the Vegetable Crops Research Program of the National Institute of Agricultural Research of Benin (INRAB), West Africa. The plants were grown under greenhouse condition (25 ± 5 °C, 50%–70% relative humidity, 16:8-h light: dark) in plastic pots ($8 \times 8 \times 9$ cm) filled with potting soil (VP113BIO, Peltracom, Belgium). The plants used in these experiments were at the vegetative development stage of four weeks after seedling for *A. hybridus* and *O. basilicum* and six weeks after seedling for *O. gratissimum*.

Myzus persicae and *A. fabae* were maintained on a broad bean plant–*V. faba* L. *Aphis craccivora* were collected on an *A. hybridus* crop in Libreville, Gabon ($0^{\circ}27'30.46''$ N; $9^{\circ}25'6.30''$ E) and were subsequently reared on *A. hybridus* plants. Aphid species were separately reared in $45 \times 45 \times 45$ cm net cages (BugDorm, MegaView Science, Taichung, Taiwan) under the same laboratory conditions.

Impact of *Ocimum* on the Dispersion of Unwinged Aphids

Three treatments were tested for each aphid species: (1) *A. hybridus* alone (control), (2) *A. hybridus* + *O. gratissimum* (Ah + Og), and (3) *A. hybridus* + *O. basilicum* (Figure 1). For each

treatment, one plant (*A. hybridus*, *O. gratissimum* or *O. basilicum*) was set in the center of a tray. Two batches of four plants were distributed at 8 cm and 12 cm from the central plant (Figure 1). Twenty (wingless) adult aphids were transferred to the central plant (release plant) using a fine brush. The plants located at 8 and 12 cm from the central plant were considered as a potential selected plant by aphids. The trays were individually placed in the 45 × 45 × 45 cm net cages (BugDorm, MegaView Science, Taichung, Taiwan). Twelve days later, the number of aphids on the central and selected plants at 8 cm ($n = 4$) and 12 cm ($n = 4$) were counted. Six replicates were assessed per treatment and aphid species. The repellent index (RI) was calculated using the following formula adapted from [29]:

$$RI = (P_{sp} - P_{cp}) / (P_{sp} + P_{cp}), \quad (1)$$




Figure 1: Illustration of the different treatments. Ah: *A. hybridus*, Og: *O. gratissimum*, Ob: *O. basilicum*, control (*A. hybridus* alone), 8 cm and 12 cm: distance to the central plant (0 cm).

Psp: percentage of aphids on selected plant (plants at 8 cm and 12 cm), Pcp: percentage of aphids on central plant (*A. hybridus* or *Ocimum*). Repellent index ranging from -1.00 to 1.00. Negative value indicates attractant effect toward the central plant, and positive value indicates a repellent effect to the central plant.

Impact of *Ocimum* on the Behavior of Winged Aphids in the Dual-Choice Test

A flight tunnel (90 × 45 × 45 cm) was used to evaluate how *Ocimum* plants in the vicinity of *A. hybridus* plants impacted aphid behavior. The tested modalities were: (1) an *A. hybridus* plant associated with an *Ocimum* plant (*O. gratissimum* or *O. basilicum*) versus an *A. hybridus* alone, and (2) an *A. hybridus* plant versus an *A. hybridus* (black control). Ten replicates were assessed for each modality. For each replicate, twenty winged aphids, randomly sampled from the rearing population, were released in the central area of the tunnel. After 24 h, the number of aphids on each *A. hybridus* plant associated with and without the *Ocimum* plant was recorded.

Statistical Analysis

For the dispersal test, the aphid populations recorded on the selected (8 and 12 cm) and central (0 cm) plants were grouped by treatment: control (*A. hybridus*), *A. hybridus* + *O. gratissimum* (Ah + Og) and *A. hybridus* + *O. basilicum* (Ah + Ob). In order to compare the effect of the interactions between the different factors and their main effect, Generalized Linear Models (“glm”), Negative Binomial Generalized Linear Model (“glm.nb”) and Zero-Inflated Count Data Regression (“zeroinfl”) were compared using the value of their Akaike Information Criterion (AIC), in order to determine the best-fitted model to the data. Following this comparison, the Negative Binomial Generalized Linear Model (function “glm.nb”, “package MASS”; [30]) was used for each level of analysis with aphid species, plant position, plant species and treatment as fixed variables and aphid abundance as the response variable. Two levels of comparison were performed. Firstly, aphid abundance was compared between the selected plants and the central plant for each treatment, and then the number of aphids was compared between the treatments per selected plant. ANOVA test (function “Anova”, package “car”; [31]) was performed and the “SNK.test” function (“agricolae package”; [32]) was used for means comparison. For the Repellent Index, the function “aov” was performed on data. A

probability level lower than 0.05 was considered statistically significant. All statistical tests were performed using R software version 3.6.3 [33].

Binomial proportion tests (equal distribution hypothesized) were used to compare the number of aphids on the control plant (*A. hybridus* alone) and the associated treatments (*Amaranthus* versus *Ocimum*) during the dual-choice test, according to each modality.

Results

Effect of *Ocimum* Species on the Abundance of Unwinged Aphids and Repellent Activity

Analysis of the interactions between the factors studied on the aphids shows that only “Treatments”, “Distance”, the interaction “Treatments: Distance” and “Aphids: Treatments: Distance” were statistically significant ($p < 0.05$) (Table 1). Synthesis of analysis of the deviance (Anova) for fixed factors – plants, treatments and distance – and their interactions on abundance of each aphid species on amaranth plants is available in the supplementary materials (Table S1).

Table 1: Synthesis of analysis of the deviance (Anova) for fixed factors plants, aphids, treatment and distance and their interactions on aphid’s abundance on amaranth plants.

Factors	Df	Chisq	p-Value
Plants	1	0.17	0.680
Aphids	2	0.99	0.608
Treatments	2	12.74	0.001
Distance	2	35.63	<0.001
Aphids: Treatments	4	2.74	0.603
Aphids: Distance	4	1.63	0.803
Treatments: Distance	4	627.82	<0.001
Aphids: Treatments: Distance	8	23.43	0.002

Aphid abundance in each treatment is presented in Figure 2. For all aphid species, the number of individuals decreased from the central plant (0 cm) to the selected plants (8 and 12 cm) in the control (*A. hybridus* plant alone), while the opposite effect was observed in the association treatments (Ah + Og or Ah + Ob).

Additionally, no aphid was recorded on an *Ocimum* plant. The means comparison of aphid populations between distances (the central plant and selected plants) for each treatment showed an overall significant effect ($p < 0.05$) (Figure 2). The comparison of aphid abundance between treatments shows that on the central plant (0 cm) the number of aphids was significantly higher in the controls than in the associated treatments (Ah + Og or Ah + Ob). The selected plants closer to the central plant showed no significant difference in aphid numbers compared to the central plant. However, the plant at 12 cm away showed significantly higher numbers of aphids when the central plant was a basil species.




Figure 2: Aphid number (Mean \pm SE, six replicates) on the central plant (0 cm) and selected plants (8 and 12 cm) according to the tested treatments, twelve days after aphid release. Control: *A. hybridus* alone, Ah: *A. hybridus*, Og: *O. gratissimum*, Ob: *O. basilicum*. For each distance, the bars with the same letter do not differ significantly ($p > 0.05$). Within each treatment (control, Ah + Og and Ah + Ob), the bars with the same letter do not differ significantly ($p > 0.05$).

The repellent indexes of *Ocimum* plants on aphids are presented in Figure 3. In the presence of basil plants (Ah + Og and Ah + Ob), the repellent index is positive, and close to one,

irrespective of the distance from the released plant (central plant) and the aphid species. On the other hand, in the absence of the basil plant (control), the repellent index is negative overall, but varied with aphid species. Statistical analysis showed a significant difference between the control and treatments (Ah + Og and Ah + Ob) irrespective of the distance ($p < 0.001$). The synthesis of analysis of the *Ocimum* repellent index on aphids at 8 and 12 cm is available in the supplementary materials (Table S2).




Figure 3: Repellent index towards three aphid species according to the selected plants (8 and 12 cm) compared to the central plant (0 cm) in each modality. Control: *A. hybridus* (Ah) alone, Og: *O. gratissimum*, Ob: *O. basilicum*. *** $p < 0.001$.

Repellent Activity of *Ocimum* on Winged Aphids

Two hundred aphids were tested for each modality. In general, more than 75% of the aphids responded (Table 2). Aphids were evenly distributed among *A. hybridus* when tested alone (Figure 4). In comparison, the number of aphids observed on *A. hybridus* plants associated with *Ocimum* plants (*O. gratissimum* or *O. basilicum*) was significantly less significant ($p < 0.001$) (Figure 4).

Table 2: Responding aphids in choice test according to treatment. Control: Ah versus Ah, Ah: *A. hybridus*, Og: *O. gratissimum*, Ob: *O. basilicum*.

Aphids	Treatments					
	Control		Ah Versus Ah + Og		Ah Versus Ah + Ob	
	Responding aphids (%) ^a	<i>p</i> -value	Responding aphids (%) ^a	<i>p</i> -value	Responding aphids (%) ^a	<i>p</i> -value
<i>A. craccivora</i>	166 (83)	<0.001	182 (91)	<0.001	167 (84)	<0.001
<i>A. fabae</i>	155 (78)	<0.001	158 (79)	<0.001	163 (82)	<0.001
<i>M. persicae</i>	167 (84)	<0.001	166 (83)	<0.001	178 (89)	<0.001

^a Responding insects include living individuals present in one of the two side areas of the tunnel.




Figure 4: Dual-choice of aphids on an *A. hybridus* plant associated with *O. gratissimum* or *O. basilicum* plants or not 24 h after release in the flight tunnel (mean percentage \pm SE, 10 replicates). ns: no significant ($p > 0.05$), *** $p < 0.001$.

Discussion

Pesticidal plants are generally an environmentally friendly solution and are more sustainable in integrated pest management programs than synthetic pesticides. The repellent activity of *O. gratissimum* and *O. basilicum* plants on aphids was determined, as the number of aphids in the control was higher on the central plant. In addition, in the treatments (Ah + Og and Ah + Ob), no aphid was observed on the central plant (*Ocimum* plant) but only on *A. hybridus* plants. This absence of aphids on *Ocimum* plants could firstly indicate an antifeeding or irritant effect of these plants on the aphids, which is defined as a category of repellent effect based on the physiological process of the insect [34]. In this case, a higher number of aphids would probably be observed on *A. hybridus* plants located at 8 cm than at 12 cm from the *Ocimum* plants, because they would only look for a feeding resource. However, overall there were more aphids at 12 cm than 8 cm, with a significant difference in some cases. Additionally, the repellent indexes clearly indicated that this *Ocimum* species had a significant repellent activity on the tested aphids, and this is supported by the choice test results. A similar study reported that the *O. basilicum* plants reduced the infestation level of *Vicia faba* (Fabaceae) by *A. fabae* when these plants were intercropped [19]. The repellent activity of various *Ocimum* species has been reported on other pest families, including Chrysomelidae, Pyralidae, Psyllidae, Pseudococcidae, Tetranychidae, Gelechiidae, Noctuidae, and

Plutellidae [9,17,20,21,35]. Other studies have also demonstrated the repellent effect of *Ocimum* extracts or essential oils on many storage pests [12,36] and human disease vectors [13,37].

Host plant location by insects for feeding or reproduction is affected by their ability to perceive VOCs emitted by these plants [38]. However, when host plants are located near to non-host plants, they may be less attractive to insects [39]. Indeed, VOCs from non-host plants can mask or disrupt the chemical environment of the host plants, which would prevent their recognition by pests [34,40]. Thus, it can be assumed that volatiles emitted by *Ocimum* spp., which is considered as a non-host plant, had changed the chemical environment of the amaranth, and then caused the observed repellent effect. According to some studies, the biocidal activity of plant volatiles was generally attributed to their main compounds [41,42]. Therefore, the repellent activity on aphids could be attributed to the major compounds p-cymene, γ -terpinene, α -terpinene, α -thujene E- α -bergamotene, methyl eugenol, E- β -ocimene linalool, previously identified in the volatiles collected on the tested *Ocimum* species [17].

For example, the repellent effect of linalool has been shown on *Cavariella aegopodii* Scopoli, *M. persicae* and *Rhopalosiphum maidis* Fitch [42,43]. The two *Ocimum* species had a similar effect on the three tested aphid species. As they had quite different chemical profiles, according to the relative proportions [17], this could be advantageous in management schemes, as these could be used alternatively to avoid insect habituation.

Conclusions

Our results provide evidence that *Ocimum* plants have a repellent effect on aphids. This indicates that these plants have potential as natural pesticides. These plants could then be used as an agroecological alternative for aphid control, in order to avoid the use of synthetic pesticides on amaranth or other vegetable crops in West Africa especially. The ability of *Ocimum* spp. to repel pests and attract natural enemies can make them

important companion plants, especially for small-scale farmers. Not only can they be used to control pests, they can also be harvested and commercialized, which may provide a significant economic return for farmers. These benefits can also contribute to the food safety of producers and the consumers. However, additional experiments should be carried out in the field conditions to evaluate the real repellent activity of basil on aphids and how to maximize this effect.

References

1. Bhatia V, Uniyal PL, Bhattacharya R. Aphid resistance in Brassica crops: Challenges, biotechnological progress and emerging possibilities. *Biotechnol. Adv.* 2011; 29: 879–888.
2. Parker JE, Snyder WE, Hamilton GC, Saona CR. Companion planting and insect pest control. In: Soloneski S, Larramendy M, editors. *Weed and Pest Control—Conventional and New Challenges*. Rijeka: InTech. 2013; 1–30.
3. Fouad EA, Abou-Yousef HM, Abdallah IS, Kandil MA. Resistance monitoring and enzyme activity in three field populations of cowpea aphid (*Aphis craccivora*) from Egypt. *Crop Prot.* 2016, 81, 163–167.
4. Lundström, N.L.P., Zhang, H., Brännström, Å. Pareto-efficient biological pest control enable high efficacy at small costs. *Ecol. Model.* 2017; 364: 89–97.
5. Erdog̃ an P, Yildirim A. Insecticidal activity of three different plant extracts on the green peach aphid (*Myzus persicae* Sulzer) (Hemiptera: Aphididae). *J. Entomol. Res. Soc.* 2016; 18: 27–35.
6. Petrakis EA, Kimbaris AC, Perdikis DC, Lykouressis DP, Tarantilis PA, et al. Responses of *Myzus persicae* (Sulzer) to three Lamiaceae essential oils obtained by microwave-assisted and conventional hydrodistillation. *Ind. Crops Prod.* 2014; 62: 272–279.
7. Abtew A, Subramanian S, Cheseto X, Kreiter S, Garzia G, et al. Repellency of plant extracts against the legume flower Thrips *Megalurothrips sjostedti* (Thysanoptera: Thripidae). *Insects.* 2015; 6: 608–625.
8. Zhang Z, Zhou C, Xu Y, Huang X, Zhang L, et al. Effects of

- intercropping tea with aromatic plants on population dynamics of arthropods in Chinese tea plantations. *J. Pest Sci.* 2017; 90: 227–237.
9. Kianmatee S, Ranamukhaarachchi SL. Pest repellent plants for management of insect pests of chinese kale, *Brassica oleracea* L. *Int. J. Agric. Biol.* 2007; 9: 64–67.
 10. Adeniyi SA, Orjiekwe CL, Ehiagbonare JE, Arimah BD. Preliminary phytochemical analysis and insecticidal activity of ethanolic extracts of four tropical plants (*Vernonia amygdalina*, *Sida acuta*, *Ocimum gratissimum* and *Telfaria occidentalis*) against beans weevil (*Acanthoscelides obtectus*). *Int. J. Phys. Sci.* 2010; 5: 753–762.
 11. Ma YF, Xiao C. Push-pull effects of three plant secondary metabolites on oviposition of the potato tuber moth. *J. Insect Sci.* 2013; 13: 128.
 12. Kiradoo MM, Srivastava M. A comparative study on the efficacy of two Lamiaceae plants on egg-laying performance by the pulse beetle *Callosobruchus chinensis* Linn. (Coleoptera: Bruchidae). *J. Biopestic.* 2010; 3: 590–595.
 13. Koech P, Mwangi R. Repellent activities of *Ocimum basilicum*, *Azadirachta indica* and *Eucalyptus citriodora* extracts on rabbit skin against *Aedes aegypti*. *J. Entomol. Zool. Stud.* 2013; 1: 84–91.
 14. Akono Ntonga P, Baldovini N, Mouray E, Mambu L, Belong P, et al. Activity of *Ocimum basilicum*, *Ocimum canum*, and *Cymbopogon citratus* essential oils against *Plasmodium falciparum* and mature-stage larvae of *Anopheles funestus* s.s. *Parasite.* 2014; 21: 33.
 15. Shadia E, Abd E, Elsayed AO, Aly SS. Chemical composition of *Ocimum americanum* essential oil and its biological effects against, *Agrotis ipsilon*, (Lepidoptera: Noctuidae). *Res. J. Agric. Biol. Sci.* 2007; 3: 740–747.
 16. Parolin P, Bresch C, Poncet C, Suay-Cortez R, Van Oudenhove L. Testing basil as banker plant in IPM greenhouse tomato crops. *Int. J. Pest Manag.* 2015; 61: 235–242.
 17. Yarou BB, Bawin T, Boullis A, Heukin S, Lognay G, et al. Oviposition deterrent activity of basil plants and their essentials oils against *Tuta absoluta* (Lepidoptera: Gelechiidae). *Environ. Sci. Pollut. Res.* 2017; 25: 29880–

- 29888.
18. Schader C, Zaller JG, Köpke U. Cotton-Basil intercropping: Effects on pests, yields and economical parameters in an organic field in Fayoum, Egypt. *Biol. Agric. Hortic.* 2005; 23: 59–72.
 19. Basedow T, Hua L, Aggarwal N. The infestation of *Vicia faba* L. (Fabaceae) by *Aphis fabae* (Scop.) (Homoptera: Aphididae) under the influence of Lamiaceae (*Ocimum basilicum* L. and *Satureja hortensis* L.). *J. Pest Sci.* 2006; 79: 149–154.
 20. Yarou BB, Assogba-Komlan F, Tossou E, Mensah AC, Simon S, et al. Efficacy of Basil-Cabbage intercropping to control insect pests in Benin, West Africa. *Commun. Agric. Appl. Biol. Sci.* 2017; 82: 157–166.
 21. Beizhou S, Jie Z, Jinghui H, Hongying W, Yun K, et al. Temporal dynamics of the arthropod community in pear orchards intercropped with aromatic plants. *Pest Manag. Sci.* 2011; 67: 1107–1114.
 22. Beizhou S, Tang G, Sang X, Zhang J, Yao Y, et al. Intercropping with aromatic plants hindered the occurrence of *Aphis citricola* in an apple orchard system by shifting predator–prey abundances. *Biocontrol Sci. Technol.* 2013; 23: 381–395.
 23. Tang GB, Song BZ, Zhao LL, Sang XS, Wan HH, et al. Repellent and attractive effects of herbs on insects in pear orchards intercropped with aromatic plants. *Agrofor. Syst.* 2013; 87: 273–285.
 24. Das S. *Amaranthus: A Promising Crop of Future*. Singapore: Springer Nature. 2016; 42–46.
 25. Sæthre MG, Godonou I, Hofsvang T, Tèpa-Yotto G, James B. Aphids and their natural enemies in vegetable agroecosystems in Benin. *Int. J. Trop. Insect Sci.* 2011; 31: 103–117.
 26. Singh G, Singh R. Distribution and economic importance of *Aphis* (*Aphis*) *craccivora* Koch, 1854 (Aphidini: Aphidinae: Aphididae: Hemiptera) and its food plants in India. *Int. J. Recent Adv. Multidiscip.* 2017; 4: 2274–2286.
 27. Bayendi Loudit SM, Poligui RN, Verheggen F, Francis F. Occurrence of Aphids and their predators within vegetable crops in peri-urban areas in Libreville (Gabon). *Commun.*

- Agric. Appl. Biol. Sci. 2017; 82: 199–205.
28. Achigan-Dako EG, Pasquini MW, Assogba-Komlan F, N'danikou S, Dansi A, Ambrose-Oji B. Traditional Vegetables in Benin, Institut National des Recherches Agricoles du Bénin. Impr. Cotonou: du CENAP. 2010.
 29. Stefanazzi N, Stadler T, Ferrero A. Composition and toxic, repellent and feeding deterrent activity of essential oils against the stored-grain pests *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae). *Pest Manag. Sci.* 2011; 69: 639–646.
 30. Venables WN, Ripley BD. *Modern Applied Statistics with S*. New York: Springer. 2002.
 31. Fox J, Weisberg S. *An R Companion to Applied Regression*. Los Angeles: Sage publications. 2018.
 32. De Mendiburu F. *Agricolae: Statistical Procedures for Agricultural Research*. 2020. Available online at: https://www.researchgate.net/publication/303256192_Agricolae_Statistical_Procedures_for_Agricultural_Research
 33. R Core Team. *A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. Vienna, Austria. 2020. Available online at: <https://www.R-project.org/>
 34. Deletre E, Schatz B, Bourguet D, Chandre F, Williams L, et al. Prospects for repellent in pest control: Current developments and future challenges. *Chemoecology*. 2016; 26: 127–142.
 35. Ogayo KO, Ogwenjo JO, Nyaanga JG, Ogendo JO, Wagara IN, et al. Bioefficacy of lion's ear and African basil extracts in management of adult two-spotted spider mite on french beans. In *Proceedings of the RUFORUM Fourth Biennial Conference, Maputo, Mozambique*. 2014; 21: 199–203.
 36. Asawalam E, Emosairue S, Hassanali A. Essential oil of *Ocimum grattissimum* (Labiatae) as *Sitophilus zeamais* (Coleoptera: Curculionidae) protectant. *Afr. J. Biotechnol.* 2008; 7: 3771–3776.
 37. Del Fabbro S, Nazzi F. Repellent effect of sweet basil compounds on *Ixodes ricinus* ticks. *Exp. Appl. Acarol.* 2008; 45: 219–228.
 38. Webster B, Bruce T, Pickett J, Hardie J. Volatiles functioning

- as host cues in a blend become nonhost cues when presented alone to the black bean aphid. *Anim. Behav.* 2010; 79: 451–457.
39. Dahlin I, Vucetic A, Ninkovic V, Dahlin I, Vucetic ANV. Changed host plant volatile emissions induced by chemical interaction between unattacked plants reduce aphid plant acceptance with intermorph variation. *J. Pest Sci.* 2015; 88: 249–257.
 40. Zhang Z, Luo Z, Gao Y, Bian L, Sun X, et al. Volatiles from non-host aromatic plants repel tea green leafhopper *Empoasca vitis*. *Entomol. Exp. Appl.* 2014; 153: 156–169.
 41. Chang X, Alderson PG, Wright CJ. Variation in the essential oils in different leaves of basil (*Ocimum basilicum* L.) at day time. *Open Hortic. J.* 2009; 2: 13–16.
 42. Chapman RF, Bernays EA, Simpson SJ. Attraction and repulsion of the aphid, *Cavariella aegopodii*, by plant odors. *J. Chem. Ecol.* 1981; 7: 881–888.
 43. Gbrays B, Dancewicz K, Halarewicz-Pacan A, Janusz E. Effect of natural monoterpenes on the behavior of the peach potato aphid *Myzus persicae* (Sulz.). *IOBC Wprs Bull.* 2005; 28: 29–34.

Supplementary Materials

Table S1: Synthesis of analysis of the deviance (Anova) for fixed factors - plants, treatment and distance - and their interactions on abundance of each aphid species on amaranth plants

Variable	Factors	Df	Chisq	p-value
<i>A. craccivora</i>	Plants	1	1.51	0.218
	Treatments	2	3.63	0.162
	Distances	2	12.59	0.002
	Treatments: Distances	4	378.31	<0.001
<i>A. fabae</i>	Plants	1	0.00	0.990
	Treatments	2	4.70	0.096
	Distances	2	10.13	0.006
	Treatments: Distances	4	126.87	<0.001
<i>M. persicae</i>	Plants	1	0.19	0.665
	Treatments	2	4.45	0.108
	Distances	2	15.64	<0.001
	Treatments: Distances	4	260.63	<0.001

Table S2: Synthesis of analysis of the Ocimum repellent index on aphids at 8 and 12 cm

Aphids	Treatments	Distance	Df	F-value	p-value
<i>Aphis craccivora</i>	Control versus Ah+Og	8 cm	1	48.11	<0.001
		12 cm	1	135.90	<0.001
	Control versus Ah+Ob	8 cm	1	64.13	<0.001
		12 cm	1	819.00	<0.001
<i>Aphis fabae</i>	Control versus Ah+Og	8 cm	1	23.65	<0.001
		12 cm	1	40.43	<0.001
	Control versus Ah+Ob	8 cm	1	41.24	<0.001
		12 cm	1	61.44	<0.001
<i>Myzus persicae</i>	Control versus Ah+og	8 cm	1	92.74	<0.001
		12 cm	1	61.48	<0.001
	Control versus Ah+Ob	8 cm	1	25.57	<0.001
		12 cm	1	62.57	<0.001