

Book Chapter

Multifaceted Functions of Platelets in Cancer: From Tumorigenesis to Liquid Biopsy Tool and Drug Delivery System

Melania Dovizio^{1,2†}, Patrizia Ballerini^{2,3†}, Rosa Fullone^{1,2}, Stefania Tacconelli^{1,2}, Annalisa Contursi^{1,2} and Paola Patrignani^{1,2*}

¹Department of Neuroscience, Imaging and Clinical Science, “G. d’Annunzio” University, Italy

²Center for Advanced Studies and Technology (CAST), Italy

³Department of Innovative Technologies in Medicine and Dentistry, “G. d’Annunzio” University, Italy

†These authors contributed equally.

***Corresponding Author:** Paola Patrignani, Department of Neuroscience, Imaging and Clinical Science, “G. d’Annunzio” University, 66100 Chieti, Italy

Published **June 09, 2021**

This Book Chapter is a republication of an article published by Paola Patrignani, et al. at International Journal of Molecular Sciences in December 2020. (Dovizio, M.; Ballerini, P.; Fullone, R.; Tacconelli, S.; Contursi, A.; Patrignani, P. Multifaceted Functions of Platelets in Cancer: From Tumorigenesis to Liquid Biopsy Tool and Drug Delivery System. *Int. J. Mol. Sci.* 2020, 21, 9585. <https://doi.org/10.3390/ijms21249585>)

How to cite this book chapter: Melania Dovizio, Patrizia Ballerini, Rosa Fullone, Stefania Tacconelli, Annalisa Contursi, Paola Patrignani. Multifaceted Functions of Platelets in Cancer: From Tumorigenesis to Liquid Biopsy Tool and Drug Delivery System. In: Letizia Giampietro, editor. Prime Archives in Molecular Sciences: 2nd Edition. Hyderabad, India: Vide Leaf. 2021.

© The Author(s) 2021. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License(<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author Contributions: Conceptualization: MD, PB, PP; Writing-original draft preparation: MD, PB, ST, AC, PP; Writing-review and editing: MD, PB, ST, AC, PP, RF; Funding acquisition: PP, PB; All authors have read and agreed to the published version of the manuscript.

Funding: This review was funded by Associazione Italiana per la Ricerca sul Cancro (AIRC) [IG 2017-ID. 20365 Project; Principal Investigator PP]. In addition, this review was conducted on Behalf of the Aspirin for Cancer Prevention Group (AsCaP), Wolfson Institute of Preventive Medicine, Queen Mary School of Medicine and Dentistry, University of London, UK. This study was funded by Ministero dell'Istruzione, dell'Università e della Ricerca (MIUR) [Fondi per la Ricerca Scientifica di Ateneo, (ex 60%)] to PB.

Conflicts of Interest: The authors declare no conflict of interest.

Abstract

Platelets contribute to several types of cancer through plenty of mechanisms. Upon activation, platelets release many molecules, including growth and angiogenic factors, lipids, and extracellular vesicles, and activate numerous cell-types, including vascular and immune cells, fibroblasts, and cancer cells. Hence, platelets are a crucial component of cell-cell communication. In particular, their interaction with cancer cells can enhance their malignancy and facilitate the invasion and colonization of distant organs. These findings suggest the use of antiplatelet agents to restrain cancer development and progression. Another peculiarity of platelets is their capability to uptake proteins and transcripts from the circulation. Thus, cancer patient platelets show specific proteomic and transcriptomic expression patterns, a phenomenon called tumor-educated platelets (TEP). The

transcriptomic/proteomic profile of platelets can provide information for the early detection of cancer and disease monitoring. Platelet ability to interact with tumor cells and transfer their molecular cargo has been exploited to design platelet-mediated drug delivery systems to enhance the efficacy and reduce toxicity often associated with traditional chemotherapy. Platelets are extraordinary cells with many functions whose exploitation will improve cancer diagnosis and treatment.

Keywords

Platelets; Cancer Development; Tumor-Educated Platelets; Platelet-Derived Molecules; Drug-Delivery System; Extracellular Vesicles

Introduction

Platelets are known for the role played in hemostasis and thrombosis [1,2]. They are anucleated cells arising from cytoplasmic fragmentation of megakaryocytes, mainly in the bone marrow, and contain various mediators stored in at least three major types of granules— α -granules, dense granules, and lysosomes. Platelets represent an essential link between tissue damage/dysfunction and inflammatory response acting to repair the injury, but if platelet activation is uncontrolled, it leads to chronic inflammation associated with many pathological conditions, including atherothrombosis [3] and cancer [4,5]. The analysis of randomized clinical trials (RCTs) with aspirin performed to evaluate cardiovascular benefits has shown that the daily administration, even at low-doses (75-100 mg), is associated with a reduced incidence of colorectal cancer (CRC) and long-term mortality [6,7]. Low-dose aspirin can interfere with the molecular mechanisms involved in the early onset of CRC [8].

Aspirin is responsible for cardiovascular protection at low-doses through preferential inhibition of platelet cyclooxygenase(COX)-1 activity [9, 10]. The extent of inhibition of platelet COX-1-dependent thromboxane (TX)_A₂ generation and the clinical

benefit are saturable at daily low-doses of aspirin [9-12]. Since platelets do not have a nucleus, the irreversible inhibitory effect of aspirin towards platelet COX-1 persists for the lifespan of the platelet (between 5 to 10 days) [9], even though the drug exhibits a short half-life (approximately 20 minutes) [10,11]. Irreversible enzyme inactivation in an anucleated cell with a long lifespan in vivo explains the virtually complete inhibition of platelet COX-1 activity (>97%) by low-dose aspirin given once a day [9-11]. In contrast, in a nucleated cell, aspirin irreversible inhibition of COX activity is rapidly recovered through de novo protein synthesis; thus, multiple daily dosing is necessary to obtain an adequate inhibitory effect of prostanoid generation translating into therapeutic effects [10]. Patrignani et al. [13] have shown that aspirin inhibitory effect on colon cancer cells constitutively expressing COX-2 is completely recovered 24 hours after drug exposure. Prostaglandin (PG)E₂, the main product of COX-2 activity in epithelial and stromal cells, modulates apoptosis, cell proliferation, and migration [14]. It is unlikely that low-dose aspirin given once a day exerts anticancer effects by directly inhibiting COX-2-dependent PGE₂ biosynthesis in nucleated cells [13]. Low-dose aspirin can affect COX-2 expression in stromal cells and cancer cells by inhibiting platelet activation and the release of several molecules, such as growth factors and cytokines, and lipids, including TXA₂, which contribute to COX-2 induction in the tumor microenvironment and cancer cells [8,15-17]. All these lines of evidence support a pivotal role of platelets in promoting early cancerogenic signaling pathways restrained by aspirin [8,15,16].

Nonsteroidal anti-inflammatory drugs (NSAIDs), including aspirin, can inhibit the proliferation and induce apoptosis of colon cancer cells in vitro independently from their effects on COX-dependent prostanoid biosynthesis [Reviewed in #18]. However, it is unlikely that low-dose aspirin can act as an anticancer agent via these mechanisms because they occur at millimolar concentrations. It is noteworthy that aspirin's maximal systemic plasma concentration (C_{max}) after dosing with 100 mg of enteric-coated aspirin is approximately 4 μM [11].

The use of low-dose aspirin is recommended in CRC prevention strategies [19]. Nevertheless, a crucial challenge remains in identifying cancer patients who would have a more significant benefit from aspirin therapy; this would avoid the exposure of unresponsive patients to aspirin side-effects, including the risk of severe bleeding, such as bleeding of blood vessels in the brain or gut [8,20,21].

Platelets contain a changing repertoire of proteins and genetic material, including transcripts and microRNAs [22], i.e., small noncoding RNAs that act as regulators of gene expression by posttranscriptional mechanisms [23]. Upon platelet activation, biologically active molecules are secreted in a soluble form or packaged into extracellular vesicles (EVs) [24,25]. These mediators promote chronic inflammation implicated in different precancerous lesions [26].

Numerous experimental findings indicate that platelets contribute to tumor cell extravasation and metastasis [4,5]. The results of daily aspirin RCTs have shown a reduced distant metastasis frequency in patients who developed different-types of cancer [27-29]. Interestingly, a 75 mg controlled-release preparation of aspirin with low bioavailability, which inhibits platelet COX-1 in the presystemic circulation associated with marginal systemic effects [30], was used in the Thrombosis Prevention Trial (TPT) for the primary prevention of cardiovascular disease [31]. This trial was included in Rothwell's analysis of five large randomized trials of daily aspirin (≥ 75 mg daily) versus control [32]. The results of TPT contributed to show that aspirin reduced the risk of distant metastasis by 30–40% and reduced the risk of metastatic adenocarcinoma by almost half [32]. Altogether, the evidence obtained in experimental and clinical studies sustains antiplatelet agent benefit to restrain cancer cells from spreading to distant organs.

The activation of the coagulation system is associated with unfavorable patient survival. This issue is discussed in many excellent papers [4,33], and it is not included in the present review.

Another peculiarity of platelets is that they uptake circulating proteins and RNAs/microRNAs, thus acquiring a different intracellular molecular repertoire specific to the individual clinical condition. The analysis of platelet content, as a whole of transcripts (transcriptomics: the analysis of mRNAs and microRNAs) or proteins (proteomics), has the promise of being a novel tool for diagnosis and prognosis of several diseases, including cancers (known as the liquid biopsy) [34,35].

Platelets can also be proper as a drug delivery system since, upon activation, they release vesicles that deliver their content to other cells, including cancer cells. Loaded platelets with an anticancer agent administered to mouse models of cancer deliver the compound to the malignant cells via the release of vesicles containing it [36].

Role of Platelets in Early Events of Tumorigenesis

Platelets are activated in response to altered functions of vascular cells and intestinal mucosal cells, possibly dependent on lifestyle and aging [8,15,16]. Activated platelets release several lipid mediators, including the products of arachidonic acid (AA) metabolism [i.e., mainly TXA₂ and PGE₂], a-granule proteins (both angiogenic and antiangiogenic factors, growth factors, proteases, and many cytokines), and different types of vesicles rich in micro-ribonucleic acids (microRNAs) [8,16]. Due to this capacity, activated platelets trigger numerous signaling pathways in different cell types, such as vascular, immune, epithelial, tumor cells, and fibroblasts [37]. In colonic mucosa, these events induce a tissue microenvironment promoting intestinal neoplasia [8]. Enhanced levels of TXA₂ activate fibroblasts, which acquire proliferative and migratory properties [38]. Moreover, platelet-derived molecules can induce angiogenesis [39]. We have recently generated a mouse with specific deletion of COX-1 in platelets that allowed us to identify TXA₂-dependent platelet activation as a central mechanism in intestinal inflammation and fibrosis *in vivo* [38].

A key event in intestinal tumorigenesis is represented by enhanced biosynthesis of PGE₂ [40]. It can occur early in epithelial cells via COX-1 activity, associated with the repression of the prostaglandin-degrading enzyme 15-prostaglandin dehydrogenase (15-PGDH) expression [41,42]. Mucosal PGE₂ can phosphorylate ribosomal protein S6 (p-S6) via PKA [43]. S6 is a crucial regulator of the 40S ribosome biogenesis transcriptional program, and its phosphorylation is closely related to cell growth capacity and tumor progression [44]. We have found that the rectal mucosal p-S6/S6 ratio significantly correlated with PGE₂ levels [13]. Interestingly, we have shown that the administration of enteric-coated low-dose aspirin (100 mg daily) to individuals undergoing CRC screening can cause long-lasting acetylation of COX-1 and reduced phosphorylation of S6 in apparently healthy colorectal mucosa [13]. This effect may interfere with early colorectal carcinogenesis. Low-dose aspirin causes a significant inhibitory effect on platelet COX-1, and a lower inhibition of COX-1 expressed in colorectal mucosa associated with rectal mucosa changes phenotype [13].

A further increase of PGE₂ production occurs at later stages of tumorigenesis associated with induction of COX-2 and the downstream synthase, microsomal PGE₂ synthase-1 (mPGES-1; a major terminal PGE₂ synthase), in stromal cells and then in epithelial cells [45,46]. Enhanced PGE₂ production alters the normal apoptotic processes involved in preventing the accumulation of genetic mutations and leads ultimately to enhanced proliferation [40, 47]. Moreover, PGE₂ plays a role in suppressing immune functions and facilitating tumor immune escape [48].

Platelet-derived mediators also induce epithelial-mesenchymal transition (EMT) programs, where polarized, immobile epithelial cells become migratory mesenchymal cells promoting tumorigenesis and facilitate metastatic colonization [49].

All these findings sustained the benefit of low-dose aspirin, and possibly other antiplatelet agents, in CRC prevention. Inhibition of platelet function can restrain the cascade of molecular and

biological events associated with tumorigenesis occurring in the stromal compartment and epithelial cells.

Platelets and T Cell Immunity

Platelets can directly depress T cell function both in vitro and in vivo [50]. Upon activation, platelets release transforming growth factor (TGF) β , which exerts an immunosuppressive effect. Moreover, platelets express TGF β -docking receptor glycoprotein A repetitions predominant (GARP), which acts by increasing the activation of latent TGF β [50]. Specific deletion of GARP in platelets mitigated TGF β activity at the tumor site and potentiated protective immunity against both melanoma and colon cancer [50]. T cell therapy of B16 melanoma was improved by concurrent treatment with aspirin and clopidogrel (an antiplatelet agent acting as P2Y₁₂ antagonist) [50], suggesting that platelets can suppress antitumor T cell immunity.

Natural killer (NK) cells are cytotoxic lymphocytes that play an essential role in tumor immunosurveillance, preferentially eliminating targets with a low or absent expression of MHC class I and stress-induced expression ligands for activating NK receptors [51]. It has been shown that platelets promote metastasis by protecting disseminating tumor cells from NK cell immunosurveillance [52]. Tumor cells rapidly get coated in the presence of platelets in vitro and circulating tumor cells of cancer patients display co-expression of platelet markers [53]. The transfer of MHC class I onto the tumor cell surface results in high expression levels of platelet-derived normal MHC class I [53]. The resulting "phenotype of false pretenses" disrupts recognition of tumor cell missing self, thereby impairing cytotoxicity and IFN- γ production by NK cells [53]. This is a mechanism by which malignant cells may utilize platelets to impair NK cell ability to prevent metastasis. Improved knowledge of platelet role in immune evasion of tumor cells will open the way to develop new antimetastatic therapies.

Role of Platelets in Metastasis

Platelet-tumor cell interactions in the circulation contribute to the metastatic dissemination of epithelial tumors [4]. The formation of platelet aggregates surrounding tumor cells can support tumor cell survival and protection from immune elimination [4]. Moreover, platelets contribute to tumor cell adhesion to the endothelium, which translates to tumor cell arrest and extravasation [4].

Platelets and cancer cells interact reciprocally. Platelets directly associate with cancer cells using different molecules depending on specific receptors expressed in cancer cells [5,16] (**Table 1**). Thus, distinct pharmacological agents are necessary to interrupt the direct crosstalk between platelets and different cancer types. In **Table 1**, the known determinants of the interactions between platelets and cancer cells are reported with possible inhibitory pharmacological tools that have been described. These findings suggest that selecting a specific pharmacological agent to constrain metastasis development requires the characterization of the altered expression of tumor molecules involved in the interaction with platelets.

Events associated with cancer-dependent platelet activation is the release of TXA₂ and PGE₂ (as a minor product) and proteins from α-granules [such as TGF-β and platelet-derived growth factor (PDGF)], which contribute to tumor cell dissemination into the bloodstream [16].

Table 1: Molecular determinants involved in the interaction between platelets and cancer cells and their blockage by pharmacological tools.

<i>Platelet constituent</i>	<i>Cancer cell constituent</i>	<i>Cells responses</i>	<i>Pharmacological tools</i>	<i>References</i>
Integrin αIIbβ3 (or GPIIb/IIIa)	Integrin α v β 3	Cancer cell capacity to adhere to the endothelium was increased by platelet activation under flow	Abciximab	Lonsdorf et al., 2012; Felding-Habermann et al., 1996 [103, 209]
Integrin αIIbβ3 (or GPIIb/IIIa)	?	Platelet receptor Fc γ RIIa activation and ADP release induced platelet aggregation	Abciximab	Mitrugno et al., 2014 [210]
Integrin αIIbβ3 (or GPIIb/IIIa)	GPIIb/IIa-like complex	ADP release from cancer cells which induces platelet aggregation, degranulation and the formation of platelet-tumor cell aggregates	mAb LYP18 against GPIIb/IIIa	Boukerche et al., 1995 [211]
Integrin α6β1	ADAM9	Induction of platelet activation, granule secretion, and subsequent endothelial transmigration of tumor cells	GoH3 (Integrin α 6-blocking antibody)	Mammadova-Bach et al., 2016 [212]
P-selectin	Mucin-type glycoprotein	Direct interaction between cancer cells and activated platelets	Crizanlizumab	Mannori et al., 1995; Man et al., 2020 [213,214]
P-selectin	PSGL-1	Induction of platelet activation and platelet/cancer cell aggregates	Crizanlizumab	Man et al., 2020; Gong et al., 2012 [214,215]
P-selectin	CDC44	Induction of platelet/cancer cell aggregates under shear stress	Crizanlizumab	Man et al., 2020; Alves et al., 2008 [214,216]
P-selectin	PCLP1	Induction of platelet activation and formation of platelet/cancer cell aggregates	Crizanlizumab	Man et al., 2020; Larrucea et al., 2007 [214,217]
GPVI	Galectin-3	COX-2 overexpression and EMT induction in cancer cells	Revacept	Dovizio et al., 2013 [17]
CLEC-2	Podoplanin	Platelet aggregation and induction of platelet/cancer cell aggregates	mAb NZ-1 against CLEC-2	Suzuki-Inoue et al., 2007; Chang et al., 2015; Kato et al., 2006; [218-220]

Labelle et al. [55] showed that direct platelet interaction with tumor cells synergizes with platelet-released TGF- β to induce EMT, therefore, increasing tumor cell invasive potential and their capacity to colonize the lung.

Dovizio et al. [17] found that PDGF released by platelet/colon cancer cell interaction contributes to the COX-2 induction in tumor cells via a posttranscriptional mechanism. Moreover, the interaction of platelets with cancer cells induces changes in the mRNA levels of EMT markers, thus conferring greater migratory and metastatic capacities [56]. Platelets can adhere to the human colon adenocarcinoma cell line HT29 through the interaction of the platelet collagen receptor GPVI with galectin-3, which is highly expressed in tumor cells and contains a collagen-like domain [17]. The role of platelet GPVI was elucidated using Revacept, a fusion protein containing the extracellular domain of human GPVI [57], which binds specifically to collagen at the vascular injury site, thus inhibiting adhesion and platelet aggregation [57]. Revacept prevented platelet-induced changes in cancer cells, such as EMT-related genes and COX-2 expression [17].

Furthermore, Guillem-Llobat et al. [56] showed that the mesenchymal phenotype induced in colon cancer cells by exposure to platelets in vitro is characterized by enhanced prothrombotic and metastatic properties when injected in vivo into mice. These effects were mitigated by administering low-dose aspirin to mice, suggesting a pivotal role of platelet activation in metastasis development [56].

Platelet- HT29 cell interaction causes the release of PGE₂ from platelets [56], which promotes EMT and migration via the activation of the PGE₂ receptor subtype EP4 on cancer cells [56]. These effects were prevented by different antiplatelet agents: aspirin (a selective inhibitor of platelet COX-1), DG-041 (an antagonist of the EP3 receptor), and ticagrelor (a P2Y₁₂ receptor antagonist) [56].

Lucotti et al. [58] found that aspirin reduced lung metastasis through inhibition of platelet COX-1. The platelet COX-1/TXA₂

pathway was responsible for this antimetastatic effect. Its inhibition affected platelet aggregation on tumor cells, endothelial activation, tumor cell adhesion to the endothelium, and recruitment of metastasis-promoting monocytes/macrophages, and diminished the formation of a premetastatic niche [58].

Platelet-Derived Vesicles and their Implication to Cancer

Platelets release EVs, a heterogeneous population of membrane-enclosed vesicles characterized by different sizes. The medium size (100-1000 nm) EVs are also called microparticles (MPs) [59]. Platelet-derived MPs are shed from plasma membranes during platelet activation, stress, apoptosis, and necrosis, and they represent the most abundant population, accounting for 70-90% among circulating EVs in the peripheral blood of healthy people [60, 61]. MP generation requires the loss of the platelet membrane phospholipid asymmetry by the enzymatic pathway "flip-flop" and scramblase that, lastly, results in the phospholipid rearrangement between inner and outer leaflets. Notably, the exposure of anionic phosphatidylserine (PS) on the outer cell membrane of platelets is key to regulating thrombin generation [62].

Membrane-derived EVs are involved in cell-cell communication through different mechanisms: (i) directly stimulate cells as a kind of "signaling complex," (ii) transfer membrane receptors, proteins, mRNA, and organelles (e.g., mitochondria) between cells, and finally (iii) deliver infectious agents into cells (e.g., human immunodeficiency virus, prions) [reviewed in ref #63].

Platelet-derived MPs can transfer specific microRNAs to cancer cells in vitro and in vivo, thus promoting phenotypic changes [64,65].

Platelet-derived MPs can influence target cells via transcellular lipid metabolism, i.e., transferring AA to endothelial cells to induce the expression of COX-2, which, in turn, metabolizes AA to produce prostanoids [66].

Several studies focused on exploring the EV role as effectors of hemostasis, thrombosis, angiogenesis, immunity, inflammation, and cancer. In a recent study, higher circulating plasma levels of MPs were observed in patients with gastric cancer than in healthy controls and, more interestingly, plasma levels of MPs, along with interleukin (IL)-6, VEGF (Vascular-Endothelial Growth Factor), and RANTES, were the highest in the patients with more advanced stages of the disease [67]. These observations reveal a role of platelet MPs in tumorigenesis and also suggest the usefulness of evaluating their count and phenotype to identify metastatic development in patients with gastric cancer [67].

A study conducted in obese individuals has shown that platelet-derived MPs of obese women showed higher heterogeneity in size and contained different levels of proteins relevant to thrombosis and tumorigenesis [68]. Also, MPs from obese individuals presented the enhanced capacity to induce EMT and endothelial-to mesenchymal transition (EndMT) in cancer cells and endothelial cells, respectively [68]. These effects might contribute to the increased risk of developing thrombosis and multiple malignancies observed in obesity [68].

Recently, the analysis of the proteomic cargo of plasma EVs in cancer patients identified a panel of tumor-specific proteins able to detect cancer and classify its primary origin, thus proposing the analysis of EV as a promising for cancer detection [69].

Antiplatelet Drugs and Cancer

Results of clinical and experimental studies support that targeting platelet activation is involved in cancer development and the promotion of metastasis, opening the way to the possible chemopreventive use of antiplatelet agents [4,8,10,15,16]. The finding that low-dose aspirin (75–100 mg), which mainly targets platelets [9-11], reduces the incidence and mortality of several types of cancer, including CRC [6-8,27-29], led to recommend it for the primary prevention of cancer [19]. The US Preventive Services Task Force stated that low-dose aspirin should be used for the primary prevention of cardiovascular disease (CVD) and

CRC in adults aged 50–59 years "who have a 10% or greater 10-year CVD risk, are not at increased risk for bleeding, have a life expectancy of at least ten years, and are willing to take low-dose aspirin daily for at least ten years" [19]. However, we believe that some issues remain to be clarified before giving a conclusive recommendation on the use of aspirin for cancer chemoprevention. As discussed above, the knowledge of low-dose aspirin pharmacodynamics and pharmacokinetics supports that the drug acts as an anticancer agent by targeting platelet-dependent signalings of tumorigenesis [8-11,13,15-17]. Low-dose aspirin can indirectly restrain COX-2 induction in an adenomatous lesion by inhibiting platelet function. Clinical studies in CRC patients treated with low-dose aspirin are undergoing to obtain conclusive evidence of the proposed mechanism. To this aim, we have recently developed a direct biomarker of aspirin action by assessing the extent of acetylation of COX-2 in tissue biopsies [70]. The use of this novel biomarker of aspirin action will also be crucial to clarify whether higher doses of aspirin might translate into enhanced inhibition of COX-2-dependent PGE₂ biosynthesis in cancerous lesions. It is also necessary to clarify whether (i) other antiplatelet agents exert anticancer effects and (ii) their coadministration can improve the chemopreventive effect of low-dose aspirin.

Aspirin

Aspirin, acetylsalicylic acid (ASA), is a member of NSAIDs. Unlike other NSAIDs, aspirin inhibits prostanoid biosynthesis by an irreversible inactivation of COX-1 and COX-2, through the acetylation of a specific serine residue located in the cyclooxygenase active site, at position 529 and 516 of COX-1 and COX-2, respectively [reviewed in refs #8,10,16]. Despite the short half-life (i.e., 20 min) of the drug, aspirin administration at low-doses once a day causes an antiplatelet effect because of irreversible COX-1 inactivation (occurring both in the pre-systemic and systemic circulation) in the anucleated platelets characterized by a low rate of protein synthesis. The chronic administration with low-dose aspirin causes a virtually complete inhibition of platelet COX-1 activity (97%), maximal platelet

acetylation of COX-1 (75%), and inhibition of platelet function persisting throughout the dosing interval (i.e., 24 hours) [9,11].

The first evidence of aspirin chemopreventive effect against CRC was obtained in epidemiological studies. Both case-control and cohort studies have shown that the regular and continued use of aspirin is associated with an approximately 50% reduction in the incidence and mortality for CRC [71,72]. However, this benefit may not be evident until after at least a decade of regular aspirin consumption [73]. RCTs with different aspirin doses (81–600 mg daily) were carried out in different clinical conditions (**Table 2**) [74-87]. In two RCTs performed in healthy individuals [The Physicians' Health (PHS) and Women's Health (WHS) Studies], aspirin (325 and 100 mg, respectively) failed to reduce the risk of CRC (Table 2) [74,75], probably due to the inadequate treatment duration or follow-up and/or administration schedule (**Table 2**). Differently, in prospective, placebo-controlled RCTs performed in average/high-risk population, aspirin at low–medium doses (81–325 mg daily) significantly reduces the risk of colorectal adenoma recurrence [76-80] (**Table 2**). Up to date, limited information is available on aspirin's chemopreventive effect in Familial adenomatous polyposis (FAP) patients. In the study performed by Burn et al. [81], an international, multicenter, randomized, placebo-controlled trial of aspirin (600 mg/day) and/or resistant starch (RS) (30 g/day) (CAPP1) performed in young FAP patients (from ten to 21 years of age), the daily administration of aspirin 600 mg from one to 12 years did not meet the primary end-point consisting in the reduction of polyp number in the rectum and sigmoid colon. More recently, it was found that the same aspirin treatment, administered to Lynch syndrome patients (CAPP2 trial), was associated with a significant reduction of CRC development [82]. The results of the 10-year follow-up of the CAPP2 trial have been recently published [83]. The data show that adult carriers of a pathogenic mismatch repair gene defect (Lynch syndrome) should take 600 mg aspirin daily for at least two years to significantly reduce future cancer risk. However, this effect does not become apparent for at least four years. Importantly, adverse events during the intervention phase between aspirin and placebo groups were similar [83]. The

ongoing CaPP3 Study aims to define the optimal dose of aspirin (600 mg daily versus 300 mg and 100 mg daily) for cancer prevention versus adverse events in Lynch syndrome.

Recently, low-dose aspirin on cancer incidence and mortality was investigated in diabetic patients and elderly populations [84-87]. As reported in **Table 2**, Okada et al. showed data in type 2 diabetic patients [84] administered with aspirin at doses of 81 or 100 mg daily for 10.7 years, in which the drug caused a significant reduction of cancer incidence. Differently, the results of the ASCEND trial showed that the rate of cancer-related mortality in type 2 diabetics patients was unaffected by low-dose aspirin administration [85]. In the elderly population administered with aspirin 100 mg/die, the drug failed to satisfy the secondary end-point, i.e., cancer mortality rate [86,87]. All these studies have the limitation of the small cohort possibly inadequate to evaluate aspirin chemopreventive effect.

The ADD-Aspirin trial is an ongoing adjuvant trial in patients with newly diagnosed cancers, including colorectal, gastroesophageal, breast, and prostate cancer, administered with aspirin 100 mg or 300 mg /die [88]. The results of this trial will clarify whether regular aspirin use after treatment for early-stage cancer can prevent cancer recurrence and death. Moreover, it will define the appropriate dose of aspirin to be administered (**Table 2**).

Table 2: Randomized clinical trials which have compared the effect of aspirin in the prevention of colorectal adenoma or cancer versus comparator.

<i>Treatment</i>	<i>Patients</i>	<i>Primary end-point</i>	<i>Results (95%CI)</i>	<i>References</i>
Placebo or aspirin 325 mg every day for 5 years	Male physicians aged between 40 and 84 years	Incidence of total cancer	RR: 1.15 (0.80-1.65) for colorectal cancer	Gann et al.,1993. [74]
Placebo or aspirin 100mg every other day for an average of 10.1 years	Healthy women aged at least 45 years	Confirmed newly diagnosed invasive cancer at any site	RR: 1.01(0.94-1.08) for total cancer; RR: 0.97(0.77-1.24) for colorectal cancer	Cook et al.,2005 [75]
Placebo, aspirin 81 mg or 325 mg daily for 2.8 years	Patients with a recent history of histologically documented (removed) adenomas	Proportion of patients in whom one or more colorectal adenomas were detected	Any adenoma; RR: 0.81 (0.69–0.96) for ASA 81mg, P=0.04; RR: 0.96 (0.81–1.13) for ASA 325 mg Advanced lesion; RR: 0.59 (0.38–0.92) for ASA 81mg; RR: 0.83 (0.55–1.23) for ASA 325 mg	Baron et al., 2003. [76]
Placebo or enteric coated-aspirin 325 mg daily for 2.6 years	Patients who had histologically documented colon or rectal cancer with a low risk of disease recurrence	Detection of adenomas in the large bowel by either colonoscopy or sigmoidoscopy after randomization	RR: 0.65(0.46-0.91) P=0.004	Sandler et al., 2003. [77]
Placebo or soluble aspirin (160 or 300 mg daily) for 1 year	Patients with a history of colorectal adenomas	Adenoma recurrence after 1 year	RR: 0.73 (0.52–1.04) P=0.04 for both doses	Benamouzig et al.,2003 (APACC trial) [78]
Placebo or soluble aspirin (160 or 300 mg daily) for 4 years	Patients with a history of colorectal adenomas	Adenoma recurrence after 4 years	RR: 0.96(0.75-1.22) for both doses	Benamouzig et al., 2012 (APACC trial) [79]
Aspirin (300 mg daily) versus folate supplements (0.5 mg/day) for about 2.6 years	Patients with an adenoma removed in the 6 months before recruitment	A colorectal adenoma diagnosed after baseline	RR: 0.79(0.63–0.99) P=0.04	Logan et al., 2008. [80]
Aspirin (600 mg/day) and/or resistant starch (30 g/day) for 17 years	FAP young patients	Polyp number in the rectum and sigmoid colon	RR: 0.77 (0.54–1.10)	Burn et al., 2011. (CAPP1 trial) [81]
Aspirin (600 mg/day) and/or resistant starch (30 g/day), or placebo for a mean of 10 years	Lynch syndrome (hereditary non-polyposis colon cancer or HNPCC)	Development of colorectal cancer	HR: 0.65(0.43–0.97; p=0.035) Aspirin vs placebo	Burn et al., 2020. (CAPP2 trial). [83]
Aspirin 81 or 100 mg daily for 10.7years	Type 2 diabetic patients	Time to first cancer incidence	HR: 0.92 (0.73–1.14) P = 0.4; adjusted HR, 0.66 (0.43–0.99) P = 0.04	Okada et al., 2018. [84]
Aspirin 100 mg or placebo for 7.4 years	Subjects with a diagnosis of diabetes mellitus (any type) aged >40years	First serious vascular event	RR: 0.98 (0.84–1.15) for cancer-related mortality	Bowman et al., 2018. (ASCEND study) [85]
Aspirin 100 mg or placebo for 4.7 years	Healthy Elderly (>70years)	Composite of death, dementia, or persistent physical disability	HR: 1.35 (1.13 to 1.61) for cancer mortality; HR: 1.04 (0.95-1.14) for cancer incidence	McNeil et al., 2018; McNeil et al., 2020. [87,87]
Aspirin 100 mg or 300 mg or placebo daily for at least five years	Patients who have undergone potentially curative treatment for breast, colorectal, gastro-oesophageal or prostate cancer	Invasive disease-free survival for the breast cohort, disease-free survival for the colorectal cohort, overall survival for the gastro-oesophageal cohort	Ongoing	Coyle et al., 2016. (ADD-ASPIRIN trial) [88]

RR: Relative risk; HR: Hazard Ratio. Association pour la Prévention par l'Aspirine du Cancer Colorectal (APACC); Colorectal Adenoma/Carcinoma Prevention Programme (CAPP); A Study of Cardiovascular Events in Diabetes (ASCEND)

P2Y12 Receptor Antagonists

ADP, released from dense granules of platelets, induces platelet aggregation, but it is also involved in platelet granule secretion [2]. These effects occur via the activation of the three types of purinergic receptors expressed in platelets, the P2X-type ion channel-linked receptor and two P2Y-type G protein-coupled receptors, P2Y1 and P2Y12 [89]. P2Y1 and P2Y12 are coupled to Gq and Gi proteins, which activate PLC and inhibit adenylyl cyclase, respectively [89].

ADP-dependent activation of the P2Y1 receptor leads to an increase in intracellular calcium via Gq coupling; P2Y12 activation translates into Gi coupling, thus resulting in the inhibition of adenylyl cyclase and the prevention of the phosphorylation of vasodilator-stimulated phosphoprotein (VASP), whose phosphorylation correlates with the inhibition of platelet function [89]. Also, the stimulation of P2Y12 leads to the activation of phosphoinositide 3-kinase (PI3K) and Akt kinase [89].

The ADP platelet receptor P2Y12 is the target of a family of antithrombotic agents, which includes the thienopyridines (ticlopidine, clopidogrel, and prasugrel) that irreversibly inhibit the receptor and the novel direct and reversible antagonists (ticagrelor, cangrelor, and elinogrel) [91].

A possible role of the P2Y12 receptor in cancer has been shown. Wang et al. [92] demonstrated that tumor metastases are reduced in P2Y12-deficient mice. In animal models of spontaneous or experimentally induced lung metastasis, obtained by injecting Lewis lung carcinoma cells or B16 melanoma cells, respectively, the P2Y12 receptor deficiency was reported to be linked to a reduced weight of metastasis [93]. The coadministration of the antiplatelet drugs aspirin and clopidogrel prevented or delayed the development of hepatocellular carcinoma and improved survival in a mouse model of chronic immune-mediated hepatitis B [93].

Moreover, ticlopidine administration in a rat model of spontaneous pulmonary metastasis of Lewis lung carcinoma suppressed the dissemination process [94]. Ticagrelor reduced the lung colony-forming units and improved survival in an orthotopic 4T1 breast cancer model obtained by inoculating 4T1 mammary carcinoma cells into the mammary pad of female BALB/c mice [95]. In this study, ticagrelor significantly reduced the number of tumor cell-platelet aggregates present in the lung at 10, 30, and 60 min after the intravenous administration of 4T1 cells [95].

Up to date, no RCTs assessed the effect of P2Y₁₂ blockage in cancer patients. Recently, an epidemiological study performed by the analysis of a Spanish primary care database found that clopidogrel use reduced the risk of CRC similar to low-dose aspirin (20%–30% reduction in risk), starting at one year of treatment [96]. Interestingly, the dual antiplatelet therapy (aspirin and clopidogrel) had the same effect versus clopidogrel or low-dose aspirin monotherapy [96]. These data support the concept that inhibiting platelet activation signaling-pathways is an effective strategy for preventing CRC.

Thrombin Receptor Antagonists

Thrombin induces platelet activation by two PAR family members, PAR-1, the primary thrombin receptor, and PAR-4 [97]. These receptors may play a crucial role in regulating angiogenesis and, in turn, modulate the processes of wound healing and tumor growth [98]. PAR-1 and PAR-4 activation cause the secretion of different sets of α -granules from platelets [99]. Experimental pieces of evidence obtained *in vitro* and *in vivo* have suggested that the control of PAR1-mediated signaling may represent a promising strategy for malignancy treatment. The subcutaneous injection of MC38 cells, an aggressive C57Bl/6-derived colonic adenocarcinoma cell line, in PAR-1^{-/-} and wild-type mice caused the development of palpable tumors in both genotypes [100]. However, the tumors grew significantly more slowly in PAR-1^{-/-} mice than in control animals [100], suggesting a role for stromal-PAR-1 in tumor outgrowth. Moreover, the PAR-1 antagonist vorapaxar pretreatment of three

different epithelial ovarian cancer cells (SKOV-3, OVCAR-3, and CaOV-3) reduced the thrombin-induced cell proliferation, which returned to the baseline values [101].

Glycoprotein IIb/IIIa Antagonists

Glycoprotein IIb/IIIa (GPIIb/IIIa, also known as integrin α IIb β 3), formed via calcium-dependent association of GPIIb and GPIIIa, is a receptor for fibrinogen and von Willebrand factor [1]. GPIIb/IIIa plays a fundamental role in platelet functions, such as aggregation and adhesion. Many experimental pieces of evidence have demonstrated the contribution of platelet receptor GPIIb/IIIa on the crosstalk between platelets and cancer cells [5]. In particular, it was found that melanoma cells (M3Dau) interact with platelets by the platelet receptor GPIIb/IIIa and the GPIIb/IIIa-like complex expressed on tumor cells; also, the monoclonal antibody (LYP18) against the platelet GPIIb/IIIa complex inhibited platelet–melanoma cell interactions and platelet–platelet aggregation [102] (**Table 1**). Lonsdorf et al. [103] found that platelets are relevant for melanoma cell adhesion; they identified an interaction between the integrins GPIIb/IIIa of platelets and α v β 3 of murine B16 melanoma cells. In a mouse model of hematogenous metastasis to the lung, the treatment with mAb blocking the α v subunit of the α v β 3 integrin was associated with a decreased rate of lung metastasis formation by B16 melanoma cells [103].

Some in vitro pieces of evidence have shown that tirofiban, abciximab, and eptifibatide (three GPIIb/IIIa blockers currently used in cardiovascular prevention) inhibit cell proliferation and migration and cause apoptosis [104-106]. However, the enhanced risk for serious adverse events, such as bleeding, associated with their use [reviewed in ref#107] limits their long-term use as chemotherapeutic agents.

Platelet Omics as a Diagnostic and Prognostic Tool in Cancer

The discovery of platelets occurred in 1881 by Bizzozzero [108], and now, their physiology and the role in atherothrombosis are

clarified [1, 2]. However, the knowledge of novel platelet functions has shown their contribution to cancer development and progression.

Although antiplatelet drugs can be used to prevent CVD and cancer, their administration is associated with side-effects, including enhanced risk of bleeding [8]. Some risk factors (i.e., age over 65 years) of enhanced bleeding in long-term antiplatelet treatment after vascular events have been identified [109]. However, it remains to characterize a specific individual phenotype associated with enhanced susceptibility to harm by antiplatelet drug treatments. Furthermore, the patient phenotype receiving increased benefit from the treatment has to be defined. The use of omics technology and big data analysis using artificial intelligence will allow developing a personalized treatment with antiplatelet agents.

The omics analysis can capture dysregulated molecular processes and integrate them into a broader network of biological systems. For example, the advent of omics technology, such as proteomics, metabolomics, and lipidomics, using mass spectrometry, microarray- or RNA-sequencing-(RNA-seq)-based transcriptomics, and most recently ribosome footprint-based translomics has enabled a more detailed understanding of platelet biology. These new analyses will be of valuable help to identify individual fingerprints associated with disease susceptibility and intersubject variability to drug response.

Platelet Proteomics

In 1979, the first proteomic analysis of platelets was reported by Clemetson and colleagues [110]. The most abundant platelet proteins were identified through the two-dimensional gel electrophoresis and Coomassie staining of a platelet protein sample, i.e., actin, myosin, tubulin, fibrinogen, and several glycoproteins [110]. This methodology was applied for the assessment of the proteomic profile in resting [111] and activated platelets [112], platelet releasate [113], platelet MPs [114], alpha and dense granules [115,116]. Mass-spectrometry

technology and data analysis have led to a more in-depth quantitative analysis of platelet proteomic profile. For the first time, Burkhart and colleagues reported the quantitative proteomic analysis of resting platelets isolated from healthy volunteers, representing an essential reference for subsequent proteomic studies [117]. In 2013, Wijten et al. [118] published the protein dataset obtained from human activated platelets. They identified 124 proteins released upon activation, while Coppinger and colleagues identified more than 300 proteins released by human platelets following thrombin activation [113]. The discrepancy found in proteomic studies regarding the number of proteins identified may be due to the peptide identification criteria, sample prefractionation before mass spectrometry analysis, longer run times, and more sensitive mass spectrometry instruments.

The proteomic repertoire of platelets derived from (i) megakaryocyte inheritance, (ii) platelet uptake of circulating proteins, and (iii) *de novo* protein synthesis process [119]. Platelets contain all necessary protein synthesis components, such as the precursor messenger RNA (pre-mRNA), a spliceosome to process this pre-mRNA to mature mRNA endoplasmic reticulum, ribosomes, and transfer RNA [120]. Evangelista et al. [119] detected COX-1 mRNA in resting platelets, and after stimulation, *de novo* COX-1 synthesis occurred. This process may interfere with the complete and persistent suppression of TXA₂ biosynthesis by aspirin necessary for cardioprotection.

Recently, a new technology to study platelet translation, based on RNA-seq of ribosome-protected fragments (translatomics), was developed [121]. In 2017, Mills and collaborators performed the translatomics of thrombin-stimulated platelets [122]. Interestingly, they found a significant, but moderate, correlation between platelet transcriptome with the platelet translome and the platelet translome with the proteome. This data reinforced the concept that the platelet protein synthesis participates only marginally in platelet proteome composition [122].

The platelet proteome may be considered both as a diagnostic and predictive biomarker in cancer patients. Platelets store high levels of growth factors, such as VEGF, which is uptaken mainly from the circulation [33]. In cancer patients, VEGF is sequestered by platelets, and the total quantity of VEGF is 6.5 times higher than that in serum and 28.2 times higher than that in plasma [123]. Further studies are needed to clarify how platelet contribute to the increase of VEGF in cancer and whether the amount of VEGF sequestration by platelets could influence the response to anti-VEGF therapy.

Several human studies performed in cancer patients revealed elevated levels of circulating angiogenesis-regulator proteins versus healthy subjects, such as VEGF, ANGPT-1, MMP-2, platelet factor 4 (PF-4), and PDGF [124], which can be uptaken by platelets. Thus, fluctuations in the platelet proteome may serve as surrogate markers of tumor activity. Salgado et al. [125] have shown that platelet VEGF levels positively correlated with tumor growth kinetics in breast cancer patients. In patients with colorectal cancer, Peterson et al. [126] have found that the platelet concentrations of different angiogenesis regulatory proteins, i.e., VEGF, PDGF, and PF4, were higher than healthy controls. Using these data, they accurately discriminated patients with cancer from the control group [126]. More recently, Sabrkhany et al. [127] performed a proteome analysis in platelets isolated from early-stage lung or head of pancreas cancer and healthy matched-controls. In this study, the authors identified 4384 proteins, 85 significantly modulated in early-stage cancer than controls, and seven associated with early-stage cancer [127].

Platelet Transcriptome

Transcriptomics analysis consists of studying the entire landscape of RNA transcripts in a cell type or tissue. In addition to the transcriptome studies on coding mRNAs, the noncoding transcriptome (microRNAs, YRNAs, circular RNAs, long noncoding RNAs) is also gaining attention for its role as platelet function marker in various conditions [128,129]. Several studies have shown the association between the transcriptomic profile of

platelets with different pathological conditions, such as inflammatory [130], hematological [131], oncologic [132], autoimmune [133], metabolic [134], nephrological [135], and cardiovascular diseases. In particular, distinct platelet transcriptomes for patients with stable and acute coronary heart disease [136-138], coronary artery bypass graft surgery [139], peripheral artery disease [140], and atrial fibrillation [141] were described.

The platelet transcriptome analysis is of particular interest since it can define platelet-educated platelets and their use as a diagnostic and prognostic surrogate in different types of tumors.

Tumor-Educated Platelets as a Liquid Biopsy Tool in Cancer

Several studies have shown that tumor disease presence influences the platelet count, volume, activation status, proteins, and RNA content, and these changes happened already at the early stages of the disease [33].

Liquid biopsy represents an important tool in cancer for its diagnostic and prognostic values. It is a minimally invasive method for detecting and monitoring cancer. Liquid biopsies permit the early detection of cancer and prognosis for the individual patient for tumor stage and identification of new targets for personalized treatment. However, its clinical use requires validation in large-size clinical studies with outcomes. Up to date, several blood-based biosources are evaluated as liquid biopsies: the analysis of cancer cell-free DNA and RNA, proteins, and circulating tumor cells [34, 142-144]. Furthermore, EVs (exosomes) have been suggested as liquid biopsy since they harbor proteins and RNA molecules and surface membrane proteins correlated to organ tropism for cancer metastasis [145].

The first observation of the significant differences of the platelet genomic profiles in cancer patients versus healthy individuals was reported by Calverley et al. [146]. They found that 197 platelet-associated genes were downregulated in patients with metastatic lung cancer, and multiple genes were also spliced

differentially between the control and patient groups [146]. Moreover, Nilsson et al. [147] reported that platelets from cancer patients could actively absorb tumor-derived EVs and take up RNA from tumor cells. Best et al. [132] defined for the first time that the analysis of tumor-educated platelets (TEPs) can represent a potential tool for cancer diagnosis and screening. TEPs from cancer patients can release locally and systemically EVs, which can deliver their cargo, including RNAs [147-149] and proteins [150]; also, TEPs can alter their spliced RNA profile [34,133,146,151]. The analysis of the transcriptome of TEPs from cancer patients is a potential diagnostic tool for several types of cancer [34]. TEPs were evaluated in a study performed in patients with glioma [147]. It was shown that cancer cells transfer mutant RNA into blood platelets and TEPs isolated from glioma patients contained EGFRvIII mutant RNA molecules [147]. Similar evidence was found in platelets from non-small-cell lung carcinoma (NSCLC) or prostate cancer (PC) patients [148,149]. In NSCLC patients, the EML4-ALK rearrangement was found in platelets [148]. PC platelets contained the transcripts for the PC-associated biomarkers, such as kallikrein-related peptidase-2 and -3, folate hydrolase 1, and neuropeptide-Y [150]. Using an algorithm, the swarm-intelligence, to optimize the biomarker spliced RNA panel, it was possible to diagnose the late-stage NSCLC with great accuracy [151]. The recent analysis in patients with multiple myeloma identified differential platelet RNA profiles that discriminated between healthy individuals and patients with smoldering multiple myeloma [34]. Best et al. [132] developed the thromboSeq platform, in which the use of RNA sequencing-methodology identified the spliced-RNA profiles to discriminate patients with localized and metastasized cancer from healthy individuals with 84% to 96% accuracy, respectively. This technology can permit defining the organ-of-origin of the primary tumor with 71-89% accuracy [132]. Also, the spliced-RNA profile of NSCLC patient platelets identified the presence of tumor tissue-derived molecular biomarkers, such as the EGFR and KRAS mutations and HER2 and MET amplification with 85% to 95% accuracy [132]. The gene ontology analysis of RNA profiles found in platelets revealed that these RNAs are implicated in platelet activity, platelet vesiculation, and a

decrease in RNA maintenance and splicing was also observed [132]. These analysis results were corroborated by another study performed in NSCLC patients, which showed that the platelet-RNA panel containing 698 RNAs overlapped by 77% with an 1820-RNA gene panel associated with younger platelets, including the RNA for membrane marker P-selectin, previously correlated to younger reticulated platelets [150].

Interestingly, Clancy et al. [152] proposed a specific RNA profile that discriminates small (old) and large (young) platelets. In large platelets, an enrichment for gene ontologies associated with platelets/vesicles and hemostasis was found. The gene ontology studies on tumor-educated platelets suggest that cancer patients are characterized by younger platelets, possibly contributing to their enhanced thrombotic potential in cancer [152]. Also, it has been proposed that the older, smaller platelets accumulate a repertoire of vascular cell-derived RNAs during their lifespan [152], indicating that platelet subpopulations correlate with enrichment or depletion of specific transcripts.

Despite the great interest in the platelet molecule content analysis, it is necessary to validate how to collect and isolate platelets and EVs and the omics assessment within different platforms before recommending it as a diagnostic and prognostic tool. An international collaboration network of experts should be organized to discuss and define validated recommendations on this complex field.

Platelet as a Drug Delivery Tool in Cancer

Different cancer types can be successfully treated and even cured if treatments, including radiation, chemotherapy, immunotherapy, or surgery, are performed promptly following an early diagnosis. All these approaches aim to contain tumor growth or preventing metastatic formation. Chemotherapy, which uses anticancer drugs, represents one of the most common, standardized, and known therapeutic choices. However, chemotherapeutic agents have several limitations, such as the lack of selectivity, the development of severe side effects, limited efficacy, and drug resistance occurrence [153].

Moreover, the metastatic process, which represents the primary cause of cancer-associated mortality, is particularly challenging to be constrained or abolished by the traditional therapeutic approaches [154]. Thus, the need for novel strategies has opened new avenues, among which natural cell-based drug delivery has recently received considerable attention [155].

Several types of mammalian cells, including mesenchymal stem cells, red blood cells, macrophages, neutrophils, and T cells [156-160], have been tested as ideal carriers for drug delivery to potentiate the therapeutic efficacy minimizing toxic and side effects.

In this view, more recently also platelets have been exploited as carriers for cancer therapy, given their recognized role played in the complex, bidirectional interaction with malignant cells either in the blood or in the tumor microenvironment [16,25,107,161,162].

Besides the inherent tropism of platelets to vascular injury [163], particular interest for the drug delivery process is the recent finding relative to extravasated platelet presence in the microenvironment of different kinds of tumors, including pancreatic cancer, hepatocellular carcinoma, gastric and CRC [164]. In this peculiar environment, platelets interact with cancer cells without giving rise to the aggregation process. The mechanism underlining platelet extravasation and its role in tumor progression and the metastatic process is still poorly understood. A role has been ascribed to P-selectin, as shown by Qi and coworkers, who demonstrated that P-selectin deficiency abolishes platelet deposition within solid tumors and platelet focal adhesion kinase (FAK) protein [165]. In *in vivo* experiments using orthotopic models of ovarian cancer, Haemmerle et al. demonstrated that the administration of the FAK-inhibitor GSK2256098 was able to reduce platelet infiltration and to prevent tumor growth after the withdrawal of the antiangiogenic drugs bevacizumab or pazopanib [166]. The reported homing efficacy of platelets towards tumor cells is another critical characteristic that makes them a promising tool as drug delivery vehicles for anticancer agents.

Additional features make these anucleate circulating cells particularly attractive as a novel therapeutic cell-based strategy. Indeed they: i) present a relatively long lifespan in circulation (8-10days), which could improve the pharmacokinetics of drugs usually administered intravenously; ii) can be easily harvested from the patient with minimally invasive procedures and without ethical concerns, iii) can be engineered in vitro and re-infused to the patient, iv) can be targeted with nanoparticles and even with cells directly into the bloodstream thanks to the presence of specific surface binding sites such as the highly expressed GPIIb/IIIa [167]. Recently, a bispecific tandem single-chain antibody (Tand-scFvSca-1+GPIIb/IIIa) able to bind to both activated platelets via GPIIb/IIIa receptor and to a subset of peripheral blood mononuclear cells (PBMCs) expressing the stem cell antigen-1 (Sca-1) receptor has been developed [167]. When the targeted PBMCs were injected into a mouse model of myocardial infarction, their homing capability was increased, and a marked beneficial effect was found: decreased fibrosis, increased capillary density, and restored cardiac function. This innovative platelet-based cell-targeted therapy might have a relevant translational interest. Indeed, human CD34 might be equivalent to the murine Sca-1. CD34 is highly expressed in different types of promising regenerative cells. Thus its use in humans may allow the development of platelet-based targeted cell therapies similar to those obtained with Sca-1 in mice.

Platelet Loading

Platelets are also known to act as scavengers in the bloodstream where they take up small molecules and nucleic acids and particulate material, viruses, and bacteria [168,169]. Unlike a phagocytic process in which the engulfed material is metabolized inside the cells [170], it remains intact in platelets. This is why platelets are defined as covercytes, not phagocytes. The molecules can flow simply into the platelet; however, platelets can evaginate the open canalicular system, enhancing their surface [171].

Sarkar et al. [172] demonstrated that platelets loaded with doxorubicin (DOX), an anticancer drug that inhibits topoisomerase II, were able to inhibit cancer cell growth and significantly reduce the volume of ascitic fluid in Ehrlich Ascites Carcinoma (EAC) Tumor Bearing Mice (**Figure 1**). Interestingly, thanks to platelet target specificity towards cancer cells, the therapeutic effect was achieved by a drug dose lower than that usually prescribed, thus reducing toxicity.




Figure 1: Platelet-based drug delivery systems. Four different types are categorized: a) Platelet loading; b) platelet engineering; c) non-genetic platelet cell surface engineering; d) platelet camouflaged delivery systems. Their antitumoral effect in in vivo models are also reported.

More recently, DOX was loaded into platelets through the open canalicular system and was reported to effectively treat lymphoma in a mice model obtained by subcutaneous injection of Raji cells (Burkitt's Lymphoma; B lymphocyte cell line) [173]. The drug was efficiently loaded into platelets with minimal influence on their morphology and functionality and was stable. The DOX release from DOX-platelets was pH-dependent (the maximal rate was at pH 5.5, whereas it was much lower at pH 7.4) (**Figure 1**). This result is of particular interest given that, in tumor tissue environments and even more in cancer cells, pH values are less than 7. Tumor-bearing mice treated with DOX-platelets showed a significant reduction in the tumor size compared with both controls, and free DOX treated animals, suggesting a more significant therapeutic efficacy of the drug-delivery system. Bodyweight loss, considered an early side effect, was reported only in animals injected with the free drug. Moreover, myocardial injury, the most severe DOX-induced side effect, was not observed in the group treated with the DOX-platelet system, indicating a reduction in drug toxicity.

Platelet Engineering

In the last years, the genetic engineering of blood cells has been explored as promising drug delivery carriers based on their biocompatibility, circulation performance, and targetability [174].

In particular, there is a growing interest in exploring strategies for engineering cell-surface to manipulate cell-cell and cell-extracellular matrix interactions. This approach has led to the approval by the Food and Drug Administration (FDA) of autologous T cells engineered to express a chimeric antigen receptor (CAR) as first "living drugs" for the treatment of refractory pre-B cell acute lymphoblastic leukemia and diffuse large B cell lymphoma [175].

Genetic engineering is an approach that cannot be directly applied to platelets given their peculiar feature of anucleate and terminally differentiated cells. Thus, much effort has been put

into the genetic engineering of platelet precursor cells, i.e., megakaryocytes (MK).

The first studies were carried out mainly for targeted delivery of therapeutic agents at sites of vascular injury or the correction of genetic defects tested in animal models of Glanzmann Thrombasthenia, Bernard-Soulier Syndrome, or Hemophilia A [176-179]. However, the transfection of primary MK has proved to be a challenging process given the fragile nature of these cells and the difficulty of obtaining a sufficient cell number (in the bone marrow, they represent approximately 0.4% of the total nucleated cells) [180].

A strategy to overcome these difficulties was to manipulate progenitor or precursor cells to generate platelets for specific clinical applications [181].

Substantial progress in cancer treatment is the development of immune checkpoint inhibitors (ICI) such as an anti-cytotoxic T-lymphocyte antigen 4 (CTLA4) monoclonal antibodies (mAb) (ipilimumab) and anti-programmed death-1 (PD-1) mAbs (nivolumab and pembrolizumab), able to overcome the tumor-induced immune escape [182-184]. This pharmacological approach has led to long-term survival in patients with several kinds of cancer, including melanoma, cervical, breast, and non-small-cell lung cancer [186-188], and its use is increasing in clinical practice. Unfortunately, the ICI efficacy is observed in a small proportion of treated patients, and drug resistance development often occurs.

Recently, Zhang and coworkers [189] generated genetically engineered murine MK progenitor cells able to give rise to mature platelets expressing PD-1, which were also loaded with cyclophosphamide, a regulatory T cells (Treg) inhibitor [190](**Figure 1**).

These modified platelets were administered in mice transplanted with B16F10 melanoma tumor cells. Subsequently, the animals were subjected to partial surgery to mimic residual tumors in the surgical bed. The PD-1-platelets were able to accumulate within

the tumor microenvironment and to block the PDL-1, thus inhibiting the activity of the immune-suppressive Treg and favoring the activity of CD8⁺ T lymphocytes.

Moreover, the targeted release of their cyclophosphamide cargo potentiated the anticancer effects of CD8⁺ T lymphocytes (**Figure 1**). Thus, PD-platelets infusion significantly reduced and delayed the tumor growth, whereas the infusion of wild-type platelets or PBS did not.

The PD-1- platelet treated mice also displayed a longer lifespan surviving more than 60 days without showing signs of toxicity.

For anticancer therapy, platelets were also engineered to express surface-bound necrosis factor-related apoptosis-inducing ligand (TRAIL), a cytokine that induces apoptosis selectively in a wide range of cancer cells while sparing normal ones [191] (**Figure 1**).

Murine C57BL6-derived hematopoietic stem and progenitor cells (HSPCs) transduced with the lentiviral transgene and differentiated into platelets in vitro were used [192]. The insertion of α II β promoter in TRAIL cDNA of the genome of HSPCs was used to control TRAIL expression in platelet lineages following HSPC differentiation. The TRAIL-expressing platelets were able to kill cancer cells in vitro. The engineered HSPCs were transplanted into NOD SCID gamma (NSG) bone marrow to guarantee TRAIL-expressing platelet continual production in circulation (**Figure 1**). More interestingly, this drug delivery system significantly reduced liver metastases in a mouse model of prostate cancer metastasis compared with control mice receiving empty vector-transduced HSPCs. Additionally, this approach could represent a strategy to overcome the repeated administration problem for chemotherapeutic agents.

A novel approach to trigger the synthesis of therapeutic proteins in platelets without the need for precursor cell engineering comes from Chan and colleagues [193]. They developed a system aimed at synthesizing RNA and proteins encapsulated in

liposomes and called protocells. Platelets were able to take up the protocells as assessed by flow cytometry and confocal microscopy and, upon light control activation, a dramatic increase in platelet mRNA was observed. This represents a proof-of-concept that it is possible to induce a controlled transcription of exogenous RNA in anucleate cells, opening new avenues in the platelet-based therapeutic approach.

Non-Genetic Platelet Cell Surface Engineering

Platelets can also be directly modified through non-genetic cell surface engineering to act as drug delivery systems thanks to the abundance of different kinds of binding sites on their plasma membrane. Recently, monoclonal antibodies against programmed-death ligand 1 (aPDL1) were covalently linked to resting platelets via a bifunctional maleimide linker [194] (**Figure 1**). The complex (aPDL1-platelets) resulted in being stable, and aPDL1 was promptly released from the platelet surface upon thrombin activation. Once injected into mice models of experimental metastasis and cancer recurrence, aPDL1-platelets effectively prolonged animal survival by reducing the risk of cancer regrowth and metastatic dissemination.

In order to ameliorate the targetability of platelets as a drug delivery tool for the treatment of leukemia, Hu et al. [195] engineered platelets to express anti-PD-1 antibodies and subsequently decorated their surface with hematopoietic stem cells (HSCs) through a click reaction, which takes place in a water solution with a pH range of 6-8 and at a temperature of 37°C without generating toxic by-products [196](**Figure 1**). This method resulted in a lack of the aggregation of platelet-HSC assembly and the preservation of platelet function. The generated innovative drug delivery system determined a significantly increased efficacy of the checkpoint blockade therapy by taking advantage of the recognized homing capability of HSC, which drove platelets towards bone marrow. The treatment increased the number of active T cells, potentiated the immune reaction, and effectively inhibited leukemia progression. Moreover, it also

resulted in protection against a subsequent challenge with leukemia cells.

Platelet Camouflaged Delivery Systems

In the last decade, growing interest has emerged in applying nanotechnology to cancer, and different kinds of therapeutic nanoparticle (NP)-based drug delivery systems, including liposomes, polymeric, and albumin NPs have been proposed for approval in cancer treatment or are in the advanced experimental phase [197]. However, conventional NP-based systems present some critical issues in biodistribution and targetability so that after intravenous injection, they do not accumulate at the tumor site in a sufficient amount [198].

Nanoparticle characteristics such as surface features, composition, porosity, elasticity size, and geometry can influence these biological processes impairing cancer treatment efficacy.

Cell membrane-coated NPs (CMCNPs) have been developed as a novel pharmacological strategy. This "Trojan horse" technology allows NPs to increase their circulation time and reach the tumor cells [199].

Different kinds of CMCNPs have been developed using various cells, including erythrocytes, mesenchymal stem cells, white blood cells, red blood cells, platelets, cancer cells, and bacteria [200], and most of them are in the preclinical development stage.

Recently a CMCNP has been developed by using platelet membranes. The biomimetic platform was designed for site-specific and sequential delivery of two anticancer agents: TNF-related apoptosis-inducing ligand (TRAIL), known to induce cancer cell apoptosis by selectively interacting with the surface death receptors (DR4, DR5) [195] and DOX (TRAIL-Dox-PM-NV) [200] (**Figure 1**). When injected into MDA-MB-231 tumor-bearing nude mice, this delivery system significantly inhibited tumor growth. The histologic analysis showed a remarkable cancer cell remission in animals treated with TRAIL-Dox-PM-

NV compared with controls confirmed by situ TUNEL assay, which revealed the highest apoptosis in mice's tumors treated with this biomimetic drug delivery platform. Moreover, the hematoxylin and eosin staining showed that in mice treated with TRAIL-Dox-PM-NV, a significant reduction in metastatic lung nodules was achieved compared with animals treated with saline or TRAIL-Dox-NV, thus supporting, for the first time, the hypothesis that nanoparticles camouflaged with platelet membrane can have potential application in anticancer therapy.

Increasing evidence has been accumulating on the efficacy of platelet membrane coated delivery systems. The presence of surface marker proteins such as P-Selectin, CD47, and CD44 receptors on the platelet membranes protect from immunogenic clearance and favor the specific adhesion to injured tumor tissues of these camouflaged delivery platforms.

Jing and coworkers developed a platelet-camouflaged nano cocktail by encapsulating melanin nanoparticles (MNPs) and DOX inside RGD peptide (c(RGDyC))-modified nanoscale platelet vesicles (RGD-NPVs@MNPs/DOX) [201] (**Figure 1**). The biocompatible system was tested in a tumor model generated by injecting nude mice with 5×10^5 DOX-resistant breast cancer (MDA-MB-231/ADR) and being very effective in suppressing tumor growth.

It is noteworthy that RGD-NPVs@MNPs/DOX combined with laser (808 nm, 1.5W, and 10 min) improves the chemophotothermal elimination of resistant cells and tumor vasculature, thus, completely inhibiting metastasis formation in the lung.

A platelet-membrane coated NP delivery has been tested for its ability as a radiotherapy enhancer [202]. This is a very relevant clinical issue, given that radiotherapy is still one of the most efficient treatments for solid tumors. A mesoporous silica-coated bismuth nanorod (BMSNR) was camouflaged by a platelet membrane (BMSNR@PM). The biomimetic system was injected into BALB/c mice bearing 4T1 tumors, as a model of breast cancer, followed by near-infrared irradiation (1 W cm^{-2} , 10

min) (NIR) and by local low-dose X-ray irradiation (IR) (5 Gy, 1.0 Gy min⁻¹) (BMSNR@PM/NIR/IR). BMSNR@PM/NIR/IR caused a significant reduction in tumor mass and volume, confirming this NP-camouflaged system's capability to induce a synergistic therapeutic activity and benefit greater than phototherapy radiotherapy given alone.

A similar effect has been reported for a platelet coated delivery system in potentiating the therapeutic outcome of cancer immunotherapy by ferroptosis, a recently recognized iron- and reactive oxygen species-dependent form of regulated cell death [203].

Magnetic nanoparticles (Fe₃O₄) were loaded with Sulfasalazine (SAS), a drug primarily used in the treatment of rheumatoid arthritis and ulcerative colitis [204], which was also reported to suppress tumor growth and induce ferroptosis [205]. Magnetic nanoparticles loaded with SAS were camouflaged by platelet membranes (Fe₃O₄-SAS@PLT) and injected to 4T1 metastatic tumor-bearing mice alone or in combination with anti-PD-1 antibody after lung metastasis development. Their ability to target the metastatic lung lesions was monitored by in vivo imaging technique. The biomimetic system significantly enhanced the antitumor immune response causing an almost complete suppression of tumor metastasis. Moreover, tandem mass tags (TMT)-based proteomic analysis showed that Fe₃O₄-SAS@PLT treatment was able to induce M2 phenotype macrophage polarization towards the M1 phenotype with recognized antitumor properties.

The platelet-based biomimetic formulations need to be optimized for their use in cancer treatments. Several issues remain to be addressed: i) platelets have to be isolated and functionalized in vitro before being re-injected with a risk of contamination, ii) platelets contribute to thrombosis and tumor development. Thus their administration might also exacerbate pathological conditions, iii) platelets are highly reactive, and their unpredicted activation could cause drug-induced toxicity or reduced efficacy. The use of platelet membrane to obtain camouflaged drug delivery systems can overcome these critical aspects. However,

the integrity of both platelet membranes and NP core integrity and functionality in the bloodstream are not easy to characterize. Despite evidence of safety for platelet membrane camouflaged NPs, there is no information on their biocompatibility and lack of toxicity in long-term treatments in vivo; further careful investigations are still required.

Conclusive Remarks

Numerous pieces of evidence sustain the novel roles of platelets beyond hemostasis and thrombosis. Platelets are rapidly activated in response to vascular and epithelial damage and trigger numerous signaling, translating into an acute inflammatory response to repair the cellular damage. However, uncontrolled platelet activation contributes to chronic inflammation development, considered a hallmark of atherothrombosis and tissue fibrosis and cancer. The role of platelets in tumor metastasis is now fully appreciated, and the use of antiplatelet agents should be added to adjuvant antitumor therapies. However, numerous questions remain to be addressed, significantly associated with developing a personalized use of aspirin and other antiplatelet agents to improve anticancer efficacy and reduce enhanced bleeding associated with their use. The progress of omics technology associated with available bioinformatics tools will help identify individuals susceptible to cancer development (before any clinical symptoms detectable) and direct the appropriate anticancer therapy. The detailed analysis of platelet and EV content will help clarify the patient features. This information will contribute enormously to early cancer diagnosis, monitor drug treatments, and define the patient prognosis. Therefore, the next years will be exciting because the tremendous research efforts will lead to significant advances in cancer diagnosis and treatments.

The evidence of platelet role in tumor growth and metastasis has substantially increased since the first experimental in vivo findings by Gasic et al. (1968) [206]. Platelets can play a relevant function in primary tumor growth and all metastatic dissemination steps [4]. They can infiltrate the tumor microenvironment and directly interact with cancer cells, thus

modifying their phenotype toward a more malignant one [5,65]. Additionally, platelets interact with circulating tumor cells in the bloodstream, thus preventing the deadly attack of the immune system and promoting the adhesion to the endothelium, providing signals to establish a pre-metastatic niche [4]. The crosstalk between platelets and cancer cells accounts for the release of several soluble mediators and MPs by platelets [5, 25, 56]. Platelet-derived MPs can promote cell malignancy changes by delivering genomic (i.e., RNAs, miRNAs) and protein factors.

Besides biologically active mediators release, platelets can uptake and store many circulating molecules [150]. This peculiarity of platelet biology allows the development of novel biomarkers of disease status based on platelet signatures. Liquid biopsy is a minimally invasive tool helpful for cancer detection and disease or treatment monitoring. Up to date, several *in vivo* studies were performed to characterize the transcriptomic of TEPs and propose it as a liquid biopsy for cancer detection [132,151]. This methodological approach may offer advantages over other blood-based tests since platelets represent the most abundant cells in the blood, with high-quality RNA and the capacity to change their genomic and proteomic profile in response to the different pathological conditions, including cancer [34]. Further clinical validation studies are needed to progress toward blood-based screening programs for early cancer detection.

Platelet capacity to specifically interact with tumor cells suggests that platelets can act as a biomimetic drug delivery system for antitumor therapy [207,208]. Indeed, the rapidly developing field of cell-based drug carrier systems indicates platelets as smart candidates for anticancer agent delivery due to their peculiar features: (i) simple separation from blood and ready availability, (ii) long circulation time, (iii) natural selectivity towards damaged tissue and tumor microenvironment, (iv) weak antigenicity and immunogenicity [36]. Although growing evidence suggests translational feasibility and clinical value for *in vivo* application of platelet-based biomimetic formulations,

they remain to be optimized and further investigated for their use in cancer treatments.

References

1. Ruggeri ZM. Platelets in atherothrombosis. *Nat Med.* 2002; 8: 1227-1234.
2. van der Meijden PEJ, Heemskerk JWM. Platelet biology and functions: new concepts and clinical perspectives. *Nat Rev Cardiol.* 2019; 16: 166-179.
3. Gawaz M, Langer H, May AE. Platelets in inflammation and atherogenesis. *J Clin Invest.* 2005; 115: 3378–84.
4. Gay LJ, Felding-Habermann B. Contribution of platelets to tumour metastasis. *Nat Rev Cancer.* 2011; 11:123-134.
5. Contursi A, Sacco A, Grande R, Dovizio M, Patrignani P. Platelets as crucial partners for tumor metastasis: from mechanistic aspects to pharmacological targeting. *Cell. Mol. Life Sci.* 2017; 74: 3491–507.
6. Rothwell PM, Fowkes FG, Belch JF, Ogawa H, Warlow CP, et al. Effect of daily aspirin on long-term risk of death due to cancer: analysis of individual patient data from randomised trials. *Lancet.* 2011; 377: 31-41
7. Rothwell PM, Wilson M, Elwin CE, Norrving B, Algra A, et al. Long-term effect of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials. *Lancet.* 2010; 376: 1741-50.
8. Patrignani P, Patrono C. Aspirin and cancer. *J Am Coll Cardiol.* 2016; 68: 967–976.
9. Patrignani P, Filabozzi P, Patrono C. Selective cumulative inhibition of platelet thromboxane production by low-dose aspirin in healthy subjects. *J Clin Invest.* 1982; 69: 1366–1372.
10. Patrignani P, Patrono C. Cyclooxygenase inhibitors: From pharmacology to clinical read-outs. *Biochim Biophys Acta.* 2015; 1851: 422-32.
11. Patrignani P, Tacconelli S, Piazuolo E, Di Francesco L, Dovizio M, et al. Reappraisal of the clinical pharmacology of low-dose aspirin by comparing novel direct and traditional indirect biomarkers of drug action. *J Thromb Haemost.* 2014; 12: 1320–1330.

12. Baigent C, Sudlow C, Collins R, Peto R. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *British Medical Journal*. 2002; 324: 71–86
13. Patrignani P, Sacco A, Sostres C, Bruno A, Dovizio M, et al. Low-Dose Aspirin Acetylates Cyclooxygenase-1 in Human Colorectal Mucosa: Implications for the Chemoprevention of Colorectal Cancer. *Clin Pharmacol Ther*. 2017; 102: 52-61.
14. Cha YI, DuBois RN. NSAIDs and cancer prevention: targets downstream of COX-2. *Annu Rev Med*. 2007; 58: 239-52.
15. Patrono C, Patrignani P, García Rodríguez LA. Cyclooxygenase-selective inhibition of prostanoid formation: transducing biochemical selectivity into clinical read-outs. *J Clin Invest*. 2001; 108: 7-13.
16. Patrignani P, Patrono C. Aspirin, platelet inhibition and cancer prevention. *Platelets*. 2018; 29: 779-785.
17. Dovizio M, Maier TJ, Alberti S, Di Francesco L, Marcantoni E, et al. Pharmacological inhibition of platelet-tumor cell cross-talk prevents platelet-induced overexpression of cyclooxygenase-2 in HT29 human colon carcinoma cells. *Mol Pharmacol*. 2013; 84: 25-40.
18. Dovizio M, Bruno A, Tacconelli S, Patrignani P. Mode of action of aspirin as a chemopreventive agent. *Recent Results Cancer Res*. 2013; 191: 39-65.
19. Bibbins-Domingo K. U.S. Preventive Services Task Force. Aspirin Use for the Primary Prevention of Cardiovascular Disease and Colorectal Cancer: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med*. 2016; 836-845.8.
20. Dehmer SP, Maciosek MV, Flottemesch TJ, LaFrance AB, Whitlock EP. Aspirin for the Primary Prevention of Cardiovascular Disease and Colorectal Cancer: A Decision Analysis for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2016; 164: 777-86.
21. Patrono C, Baigent C. Role of aspirin in primary prevention of cardiovascular disease. *Nat Rev Cardiol*. 2019; 16: 675-686.
22. Nagalla S, Shaw C, Kong X, Kondkar AA, Edelstein LC, et al. Platelet microRNA-mRNA coexpression profiles

- correlate with platelet reactivity. *Blood*. 2011; 117: 5189–5197.
23. Fabian MR, Sonenberg N, Filipowicz W. Regulation of mRNA translation and stability by microRNAs. *Annu Rev Biochem*. 2010; 79: 351-379.
 24. Burnouf T, Goubran HA, Chou ML, Devos D, Radosevic M. Platelet microparticles: detection and assessment of their paradoxical functional roles in disease and regenerative medicine *Blood Reviews*. 2014; 28: 155–166.
 25. Dovizio M, Bruno A, Contursi A, Grande R, Patrignani P. Platelets and extracellular vesicles in cancer: diagnostic and therapeutic implications, *Cancer and Metastasis Reviews*. 2018; 37: 455-467.
 26. Colotta F, Allavena P, Sica A, Garlanda C, Mantovani A. Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability. *Carcinogenesis*. 2009; 30: 1073-1081.
 27. Rothwell PM, Wilson M, Price JF, Belch JF, Meade TW, et al. Effect of daily aspirin on risk of cancer metastasis: a study of incident cancers during randomised controlled trials. *Lancet*. 2012; 379: 1591-601.
 28. Algra AM, Rothwell PM. Effects of regular aspirin on long-term cancer incidence and metastasis: a systematic comparison of evidence from observational studies versus randomised trials. *Lancet Oncol*. 2012; 13: 518-27.
 29. Rothwell PM, Price JF, Fowkes FG, Zanchetti A, Roncaglioni MC, et al. Short-term effects of daily aspirin on cancer incidence, mortality, and non-vascular death: analysis of the time course of risks and benefits in 51 randomised controlled trials. *Lancet*. 2012; 379: 1602-12.
 30. Charman WN, Charman SA, Monkhouse DC, Frisbee SE, Lockhart EA, et al. Biopharmaceutical characterisation of a low-dose (75 mg) controlled-release aspirin formulation. *Br J Clin Pharmacol*. 1993; 36: 470-473.
 31. The Medical Research Council's General Practice Research Framework. Thrombosis Prevention Trial: randomised trial of lowintensity oral anticoagulation with warfarin and low-dose aspirin in the primary prevention of ischaemic heart disease in men at increased risk. *Lancet*. 1998; 351: 233–241.

32. Rothwell PM, Wilson M, Price JF, Belch JF, Meade TW, et al. Effect of daily aspirin on risk of cancer metastasis: a study of incident cancers during randomised controlled trials. *Lancet*. 2012; 379: 1591-1601.
33. Haemmerle M, Stone RL, Menter DG, Afshar-Kharghan V, Sood AK. The Platelet Lifeline to Cancer: Challenges and Opportunities. *Cancer Cell*. 2018; 33: 965-983.
34. Best MG, Wesseling P, Wurdinger T. Tumor-Educated Platelets as a Noninvasive Biomarker Source for Cancer Detection and Progression Monitoring. *Cancer Res*. 2018; 78: 3407-3412.
35. Sol N, In 't Veld SGJG, Vancura A, Tjerkstra M, Leurs C, et al. Tumor-Educated Platelet RNA for the Detection and (Pseudo)progression Monitoring of Glioblastoma. *Cell Rep Med*. 2020; 1: 100101.
36. Lu Y, Hu Q, Jiang C, Gu Z. Platelet for drug delivery. *Curr Opin Biotechnol*. 2019; 58: 81-91.
37. Dovizio M, Alberti S, Guillem-Llobat P, Patrignani P. Role of platelets in inflammation and cancer: novel therapeutic strategies. *Basic Clin Pharmacol Toxicol*. 2014; 114: 118-127.
38. Sacco A, Bruno A, Contursi A, Dovizio M, Tacconelli S, et al. Platelet-Specific Deletion of Cyclooxygenase-1 Ameliorates Dextran Sulfate Sodium-Induced Colitis in Mice. *J Pharmacol Exp Ther*. 2019; 370: 416-426.
39. Daniel TO, Liu H, Morrow JD, Crews BC, Marnett LJ. Thromboxane A2 is a mediator of cyclooxygenase-2-dependent endothelial migration and angiogenesis. *Cancer Res*. 1999; 59: 4574-4577.
40. Wang D, DuBois RN. An inflammatory mediator, prostaglandin E2, in colorectal cancer. *Cancer J*. 2013; 19: 502-510.
41. Myung SJ, Rerko RM, Yan M, Platzer P, Guda K, et al. 15-Hydroxyprostaglandin dehydrogenase is an in vivo suppressor of colon tumorigenesis. *Proc Natl Acad Sci U S A*. 2006; 103: 12098-12102.
42. Smartt HJ, Greenhough A, Ordóñez-Morán P, Talero E, Cherry CA, et al. β -catenin represses expression of the tumour suppressor 15-prostaglandin dehydrogenase in the

- normal intestinal epithelium and colorectal tumour cells. *Gut*. 2012; 61: 1306-1214.
43. Moore CE, Xie J, Gomez E, Herbert TP. Identification of cAMP-dependent kinase as a third in vivo ribosomal protein S6 kinase in pancreatic beta-cells. *J Mol Biol*. 2009; 389: 480-494.
 44. Ruvinsky I, Sharon N, Lerer T, Cohen H, Stolovich-Rain M, et al. Ribosomal protein S6 phosphorylation is a determinant of cell size and glucose homeostasis. *Genes Dev*. 2005; 19: 2199-2211.
 45. Prescott SM. Is cyclooxygenase-2 the alpha and the omega in cancer? *J Clin Invest*. 2000; 105: 1511-1513.
 46. Yoshimatsu K, Golijanin D, Paty PB, Soslow RA, Jakobsson PJ, et al. Inducible microsomal prostaglandin E synthase is overexpressed in colorectal adenomas and cancer. *Clin Cancer Res*. 2001; 7: 3971-3976.
 47. Wang D, Dubois RN. Eicosanoids and cancer. *Nat Rev Cancer*. 2010; 10: 181-93.
 48. Zelenay S, van der Veen AG, Böttcher JP, Snelgrove KJ, Rogers N, et al. Cyclooxygenase-Dependent Tumor Growth through Evasion of Immunity. *Cell*. 2015; 162: 1257-1270.
 49. Kalluri R, Weinberg RA. The basics of epithelial mesenchymal transition. *J Clin Invest*. 2009; 119: 1420-1428.
 50. Rachidi S, Metelli A, Riesenber B, Wu BX, Nelson MH, et al. Platelets subvert T cell immunity against cancer via GARP-TGF β axis. *Sci Immunol*. 2017; 2: eaai7911.
 51. Marcus A, Gowen BG, Thompson TW, Iannello A, Ardolino M, et al. Recognition of tumors by the innate immune system and natural killer cells. *Adv Immunol*. 2014; 122: 91-128.
 52. Maurer S, Kropp KN, Klein G, Steinle A, Haen SP, et al. Platelet-mediated shedding of NKG2D ligands impairs NK cell immune-surveillance of tumor cells. *Oncoimmunology*. 2017; 7: e1364827.
 53. Placke T, Örgel M, Schaller M, Jung G, Rammensee HG, et al. Platelet-derived MHC class I confers a pseudonormal phenotype to cancer cells that subverts the antitumor reactivity of natural killer immune cells. *Cancer Res*. 2012; 72: 440-448.

54. Tímár J, Tóvári J, Rásó E, Mészáros L, Bereczky B, et al. Platelet-mimicry of cancer cells: epiphenomenon with clinical significance. *Oncology*. 2005; 69: 185-201.
55. Labelle M, Begum S, Hynes RO. Direct signalling between platelets and cancer cells induces an epithelial-mesenchymal-like transition and promotes metastasis. *Cancer Cell*. 2011; 20: 576-590.
56. Guillem-Llobat P, Dovizio M, Bruno A, Ricciotti E, Cufino V, et al. Aspirin prevents colorectal cancer metastasis in mice by splitting the crosstalk between platelets and tumor cells. *Oncotarget*. 2016; 7: 32462-32477.
57. Ungerer M, Rosport K, Bültmann A, Piechatzek R, Uhland K, et al. Novel antiplatelet drug revacept (Dimeric Glycoprotein VI-Fc) specifically and efficiently inhibited collagen-induced platelet aggregation without affecting general hemostasis in humans. *Circulation*. 2011; 123: 1891-1899.
58. Lucotti S, Cerutti C, Soyer M, Gil-Bernabé AM, Gomes AL, et al. Aspirin blocks formation of metastatic intravascular niches by inhibiting platelet-derived COX-1/thromboxane A2. *J Clin Invest*. 2019; 129: 1845-1862.
59. van Niel G, D'Angelo G, Raposo G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol*. 2018; 19: 213–228.
60. Nieuwland R, van der Pol E, Gardiner C, Sturk A. Platelet-Derived Microparticles. In: *Platelets (Third Edition)*. Cambridge: Academic Press. 2013; 453-467.
61. Lynch SF, Ludlam CA. Plasma microparticles and vascular disorders. *Br J Haematol*. 2007; 137: 36–48.
62. Morel O, Toti F, Bakouboula B, Grunebaum L, Freyssinet JM. Procoagulant microparticles: 'criminal partners' in atherothrombosis and deleterious cellular exchanges. *Pathophysiol Haemost Thromb*. 2006; 35: 15-22.
63. Ratajczak J, Wysoczynski M, Hayek F. Membrane-derived microvesicles: important and underappreciated mediators of cell-to-cell communication. *Leukemia*. 2006; 20: 1487–1495.
64. Tang M, Jiang L, Lin Y, Wu X, Wang K, et al. Platelet microparticle-mediated transfer of miR- 939 to epithelial

- ovarian cancer cells promotes epithelial to mesenchymal transition. *Oncotarget*. 2017; 8: 97464–97475.
65. Michael JV, Wurtzel JGT, Mao GF, Rao AK, Kolpakov A, et al. Platelet microparticles infiltrating solid tumors transfer miRNAs that suppress tumor growth. *Blood*. 2017; 130: 567–580.
 66. Barry OP, Pratico D, Lawson JA, FitzGerald GA. Transcellular activation of platelets and endothelial cells by bioactive lipids in platelet microparticles. *J Clin Invest*. 1997; 99: 2118-2127.
 67. Kim HK, Song KS, Park YS, Kang YH, Lee YJ, et al. Elevated levels of circulating platelet microparticles, VEGF, IL-6 and RANTES in patients with gastric cancer: possible role of a metastasis predictor. *Eur J Cancer*. 2003; 39: 184–191.
 68. Grande R, Dovizio M, Marcone S, Szklanna PB, Bruno A, et al. Platelet-Derived Microparticles From Obese Individuals: Characterization of Number, Size, Proteomics, and Crosstalk With Cancer and Endothelial Cells. *Front Pharmacol*. 2019; 10: 7.
 69. Hoshino A, Kim HS, Bojmar L, Gyan KE, Cioffi M, et al. Extracellular Vesicle and Particle Biomarkers Define Multiple Human Cancers. *Cell*. 2020; 20: 182.
 70. Tacconelli S, Contursi A, Falcone L, Mucci M, D'Agostino I, et al. Characterization of cyclooxygenase-2 acetylation and prostanoid inhibition by aspirin in cellular systems. *Biochem Pharmacol*. 2020; 178: 114094.
 71. Thun MJ, Henley SJ, Patrono C. Nonsteroidal anti-inflammatory drugs as anticancer agents: mechanistic, pharmacologic, and clinical issues. *J Natl Cancer Inst*. 2002; 94: 252–266.
 72. Cuzick J, Otto F, Baron JA, Brown PH, Burn J, et al. Aspirin and non-steroidal anti-inflammatory drugs for cancer prevention: an international consensus statement. *Lancet Oncol*. 2009; 10: 501–507.
 73. Giovannucci E, Egan KM, Hunter DJ, Stampfer MJ, Colditz GA, et al. Aspirin and the risk of colorectal cancer in women. *N Engl J Med*. 1995; 333: 609–614.

74. Gann PH, Manson JE, Glynn RJ, Buring JE, Hennekens CH. Low-dose aspirin and incidence of colorectal tumors in a randomized trial. *J Natl Cancer Inst.* 1993; 85: 1220–1224.
75. Cook NR, Lee IM, Gaziano JM, Gordon D, Ridker PM, et al. Low-dose aspirin in the primary prevention of cancer: the Women's Health Study: a randomized controlled trial. *JAMA.* 2005; 294: 47–55.
76. Baron JA, Cole BF, Sandler RS, Haile RW, Ahnen D, et al. A randomized trial of aspirin to prevent colorectal adenomas. *N Engl J Med.* 2003; 348: 891–899.
77. Sandler RS, Halabi S, Baron JA, Budinger S, Paskett E, et al. A randomized trial of aspirin to prevent colorectal adenomas in patients with previous colorectal cancer. *N Engl J Med.* 2003; 348: 883–890.
78. Benamouzig R, Deyra J, Martin A, Girard B, Jullian E, et al. Daily soluble aspirin and prevention of colorectal adenoma recurrence: one-year results of the APACC trial. *Gastroenterology.* 2003; 125: 328–336.
79. Benamouzig R, Uzzan B, Deyra J, Martin A, Girard B, et al. Association pour la Prévention par l'Aspirine du Cancer Colorectal Study Group APACC). Prevention by daily soluble aspirin of colorectal adenoma recurrence: 4-year results of the APACC randomised trial. *Gut.* 2012; 61: 255-261.
80. Logan RF, Grainge MJ, Shepherd VC. Aspirin and folic acid for the prevention of recurrent colorectal adenomas. *Gastroenterology.* 2008; 134: 29-38.
81. Burn J, Bishop DT, Chapman PD, Elliott F, Bertario L, et al. A randomized placebo-controlled prevention trial of aspirin and/or resistant starch in young people with familial adenomatous polyposis. *Cancer Prev Res (Phila).* 2001; 4: 655-665.
82. Burn J, Gerdes AM, Macrae F, Mecklin JP, Moeslein G, et al. Long- term effect of aspirin on cancer risk in carriers of hereditary colorectal cancer: an analysis from the CAPP2 randomised controlled trial. *Lancet.* 2011; 378: 2081-2087.
83. Burn J, Sheth H, Elliott F, Reed L, Macrae F, et al. Cancer prevention with aspirin in hereditary colorectal cancer (Lynch syndrome), 10-year follow-up and registry-based 20-

- year data in the CAPP2 study: a double-blind, randomised, placebo-controlled trial. *Lancet*. 2020; 395: 1855-1863.
84. Okada S, Morimoto T, Ogawa H, Sakuma M, Matsumoto C, et al. Effect of Aspirin on Cancer Chemoprevention in Japanese Patients With Type 2 Diabetes: 10-Year Observational Follow-up of a Randomized Controlled Trial. *Diabetes Care*. 2018; 41: 1757-1764.
 85. ASCEND Study Collaborative Group, Bowman L, Mafham M, Wallendszus K, Stevens W, et al. Effects of Aspirin for Primary Prevention in Persons with Diabetes Mellitus. *N Engl J Med*. 2018; 379: 1529-1539.
 86. McNeil JJ, Wolfe R, Woods RL, Tonkin AM, Donnan GA, et al. Effect of Aspirin on Cardiovascular Events and Bleeding in the Healthy Elderly. *N Engl J Med*. 2018; 379: 1509-1518.
 87. McNeil JJ, Gibbs P, Orchard SG, Lockery JE, Bernstein WB, et al. Effect of aspirin on cancer incidence and mortality in older adults. *J Natl Cancer Inst*. 2020; djaa114.
 88. Coyle C, Cafferty FH, Rowley S, MacKenzie M, Berkman L, et al. ADD-ASPIRIN: A phase III, double-blind, placebo controlled, randomised trial assessing the effects of aspirin on disease recurrence and survival after primary therapy in common non-metastatic solid tumours. *Contemp Clin Trials*. 2016; 51: 56-64.
 89. Fitzgerald DJ, Fitzgerald GA. Historical lessons in translational medicine: cyclooxygenase inhibition and P2Y12 antagonism. *Circ Res*. 2013; 112: 174–194.
 90. Schwarz UR, Geiger J, Walter U, Eigenthaler M. Flow cytometry analysis of intracellular VASP phosphorylation for the assessment of activating and inhibitory signal transduction pathways in human platelets—definition and detection of ticlopidine/clopidogrel effects. *Thromb Haemost*. 1999; 82: 1145–1152.
 91. Ballerini P, Dovizio M, Bruno A, Tacconelli S, Patrignani P. P2Y12 Receptors in Tumorigenesis and Metastasis. *Front Pharmacol*. 2018; 9: 66.
 92. Wang Y, Sun Y, Li D, Zhang L, Wang K, et al. Platelet P2Y12 is involved in murine pulmonary metastasis. *PLoS One*. 2013; 8: e80780.
 93. Sitia G, Aiolfi R, Di Lucia P, Mainetti M, Fiocchi A, et al. Antiplatelet therapy prevents hepatocellular carcinoma and

- improves survival in a mouse model of chronic hepatitis B. *Proc Natl Acad Sci USA*. 2012; 109: E2165–E2172.
94. Kohga S, Kinjo M, Tanaka K, Ogawa H, Ishimara M, et al. Effects of 5-(2-chlorobenzyl)- 4,5,6,7-tetrahydrothieno[3,2-C]pyridine hydrochloride (Ticlopidine), a platelet aggregation inhibitor, on blood-borne metastasis. *Cancer Res*. 1981; 41: 4710–4714.
 95. Gareau AJ, Brien C, Gebremeskel S, Liwski RS, Johnston B, et al. Ticagrelor inhibits platelet-tumor cell interactions and metastasis in human and murine breast cancer. *Clin. Exp. Metastasis*. 2018; 35: 25–35.
 96. Rodríguez-Miguel A, García-Rodríguez LA, Gil M, Montoya H, Rodríguez-Martín S, et al. Clopidogrel and Low-Dose Aspirin, Alone or Together, Reduce Risk of Colorectal Cancer. *Clin Gastroenterol Hepatol*. 2019; 17: 2024-2033. e2.
 97. Andersen H, Greenberg DL, Fujikawa KW, Chung DW, Davie EW. Protease-activated receptor 1 is the primary mediator of thrombin-stimulated platelet procoagulant activity. *Proc. Natl. Acad. Sci. USA*. 1999; 96: 11189–11193.
 98. Wojtukiewicz MZ, Hempel D, Sierko E, Tucker SC, Honn KV. Protease-activated receptors (PARs)--biology and role in cancer invasion and metastasis. *Cancer Metastasis Rev*. 2015; 34: 775-796.
 99. Italiano JE Jr, Richardson JL, Patel-Hett S, Battinelli E, Zaslavsky A, et al. Angiogenesis is regulated by a novel mechanism: pro- and antiangiogenic proteins are organized into separate platelet alpha granules and differentially released. *Blood*. 2008; 111: 1227-1233.
 100. Adams GN, Rosenfeldt L, Frederick M, Miller W, Waltz D, et al. Colon Cancer Growth and Dissemination Relies upon Thrombin, Stromal PAR-1, and Fibrinogen. *Cancer Res*. 2015; 75: 4235–4243.
 101. Chanakira A, Westmark PR, Ong I.M, Sheehan JP. Tissue factor-factor VIIa complex triggers protease activated receptor 2-dependent growth factor release and migration in ovarian cancer. *Gynecol. Oncol*. 2017; 145: 167–175.
 102. Boukerche H, Berthier-Vergnes O, Tabone E, Dore´ JF, Leung LL, et al. Platelet–melanoma cell interaction is

- mediated by the glycoprotein IIb–IIIa complex. *Blood*. 1989; 74: 658–663.
103. Lonsdorf AS, Krämer BF, Fahrleitner M, Schönberger T, Gnerlich S, et al. Engagement of α IIb β 3 (GPIIb/IIIa) with α v β 3 integrin mediates interaction of melanoma cells with platelets: A connection to hematogenous metastasis. *J. Biol. Chem.* 2012; 287: 2168–2178.
104. Kononczuk J, Surazynski A, Czyzewska U, Prokop I, Tomczyk M, et al. α IIb β 3-integrin Ligands: Abciximab and Eptifibatide as Proapoptotic Factors in MCF-7 Human Breast Cancer Cells. *Curr. Drug Targets.* 2015; 16: 1429–1437.
105. Parikka M, Nissinen L, Kainulainen T, Bruckner-Tuderman L, Salo T, et al. Collagen XVII promotes integrin-mediated squamous cell carcinoma transmigration—A novel role for alphaIIb integrin and tirofiban. *Exp. Cell Res.* 2006; 312: 1431–1438.
106. Zhao F, Li L, Guan L, Yang H, Wu C, et al. Roles for GP IIb/IIIa and α v β 3 integrins in MDA-MB-231 cell invasion and shear flow-induced cancer cell mechanotransduction. *Cancer Lett.* 2014; 344: 62–73.
107. Bruno A, Dovizio M, Tacconelli S, Contursi A, Ballerini P, et al. Antithrombotic Agents and Cancer. *Cancers (Basel)*. 2018; 10: 253.
108. Bizzozero G. Su di un nuovo elemento morfologico del sangue dei mammiferi e della sua importanza nella trombosi e nella coagulazione, *L'Osservatore*. 1881; 17: 785–787.
109. Li L, Geraghty OC, Mehta Z, Rothwell PM. Oxford Vascular Study. Age-specific risks, severity, time course, and outcome of bleeding on long-term antiplatelet treatment after vascular events: a population-based cohort study. *Lancet*. 2017; 390: 490-499.
110. Clemetson KJ, Capitanio A, Lüscher EF. High resolution two-dimensional gel electrophoresis of the proteins and glycoproteins of human blood platelets and platelet membranes. *Biochim Biophys Acta*. 1979; 553: 11-24.
111. O'Neill EE, Brock CJ, von Kriegsheim AF, Pearce AC, Dwek RA, et al. Towards complete analysis of the platelet proteome. *Proteomics*. 2002; 2: 288-305.

112. García A, Prabhakar S, Hughan S, Anderson TW, Brock CJ, et al. Differential proteome analysis of TRAP- activated platelets: involvement of DOK-2 and phosphorylation of RGS proteins. *Blood*. 2004; 103: 2088-2095.
113. Coppinger JA, Cagney G, Toomey S, Kislinger T, Belton O, et al. Characterization of the proteins released from activated platelets leads to localization of novel platelet proteins in human atherosclerotic lesions. *Blood*. 2004; 103: 2096-104.
114. Garcia BA, Smalley DM, Cho H, Shabanowitz J, Ley K, et al. The platelet microparticle proteome. *J Proteome Res*. 2005; 4: 1516-1521.
115. Maynard DM, Heijnen HF, Horne MK, White JG, Gahl WA. Proteomic analysis of platelet alpha-granules using mass spectrometry. *J Thromb Haemost*. 2007; 5: 1945-55.
116. Hernandez-Ruiz L, Valverde F, Jimenez-Nuñez MD, Ocaña E, Sáez-Benito A, et al. Organellar proteomics of human platelet dense granules reveals that 14-3-3zeta is a granule protein related to atherosclerosis. *J Proteome Res*. 2007; 6: 4449-4457.
117. Burkhart JM, Vaudel M, Gambaryan S, Radau S, Walter U, et al. The first comprehensive and quantitative analysis of human platelet protein composition allows the comparative analysis of structural and functional pathways. *Blood*. 2012; 120: e73-82.
118. Wijten P, van Holten T, Woo LL, Bleijerveld OB, Roest M, et al. High precision platelet releasate definition by quantitative reversed protein profiling--brief report. *Arterioscler Thromb Vasc Biol*. 2013; 33: 1635-1638.
119. Evangelista V, Manarini S, Di Santo A, Capone ML, Ricciotti E, et al. De novo synthesis of cyclooxygenase-1 counteracts the suppression of platelet thromboxane biosynthesis by aspirin. *Circ Res*. 2006; 98: 593-595.
120. Denis MM, Tolley ND, Bunting M, Schwertz H, Jiang H, et al. Escaping the nuclear confines: signal-dependent pre-mRNA splicing in anucleate platelets. *Cell*. 2005; 122: 379-391.
121. Ingolia NT, Ghaemmaghami S, Newman JR, Weissman JS. Genome-wide analysis in vivo of translation with

- nucleotide resolution using ribosome profiling. *Science*. 2009; 324: 218-223.
122. Mills EW, Green R, Ingolia NT. Slowed decay of mRNAs enhances platelet specific translation. *Blood*. 2017; 129: e38-e48.
123. Kut C, Mac Gabhann F, Popel AS. Where is VEGF in the body? A meta-analysis of VEGF distribution in cancer. *Br J Cancer*. 2007; 97: 978-985.
124. Yan M, Lesyk G, Radziwon-Balicka A, jurasz P. Pharmacological regulation of platelet factors that influence tumor angiogenesis. *Semin Oncol*. 2014; 41: 370-377.
125. Salgado R, Benoy I, Bogers J. Platelets and vascular endothelial growth factor (VEGF): A morphological and functional study. *Angiogenesis*. 2001; 4: 37-43.
126. Peterson JE, Zurakowski D, Italiano JE Jr, Michel LV, Connors S, et al. VEGF, PF4 and PDGF are elevated in platelets of colorectal cancer patients. *Angiogenesis*. 2012; 15: 265-73.
127. Sabrkhany S, Kuijpers MJE, Knol JC, Olde Damink SWM, Dingemans AC, et al. Exploration of the platelet proteome in patients with early-stage cancer. *J Proteomics*. 2018; 177: 65-74.
128. Braza-Boïls A, Barwari T, Gutmann C, Thomas MR, Judge HM, et al. Circulating MicroRNA Levels Indicate Platelet and Leukocyte Activation in Endotoxemia Despite Platelet P2Y₁₂ Inhibition. *Int J Mol Sci*. 2020; 21: 2897.
129. Gutmann C, Joshi A, Mayr M. Platelet "-omics" in health and cardiovascular disease. *Atherosclerosis*. 2020; 307: 87-96.
130. McManus DD, Beaulieu LM, Mick E, Tanriverdi K, Larson MG, et al. Relationship among circulating inflammatory proteins, platelet gene expression, and cardiovascular risk. *Arterioscler Thromb Vasc Biol*. 2013; 33: 2666-73.
131. Raghavachari N, Xu X, Harris A, Villagra J, Logun C, et al. Amplified expression profiling of platelet transcriptome reveals changes in arginine metabolic pathways in patients with sickle cell disease. *Circulation*. 2007; 115: 1551-1562.
132. Best MG, Sol N, Kooi I, Tannous J, Westerman BA, et al. RNA-Seq of Tumor-Educated Platelets Enables Blood-

- Based Pan-Cancer, Multiclass, and Molecular Pathway Cancer Diagnostics. *Cancer Cell*. 2015; 28: 666-676.
133. Lood C, Amisten S, Gullstrand B, Jönsen A, Allhorn M, et al. Platelet transcriptional profile and protein expression in patients with systemic lupus erythematosus: up-regulation of the type I interferon system is strongly associated with vascular disease. *Blood*. 2010; 116: 1951-1957.
 134. Freedman JE, Larson MG, Tanriverdi K, O'Donnell CJ, Morin K, et al. Relation of platelet and leukocyte inflammatory transcripts to body mass index in the Framingham heart study. *Circulation*. 2010; 122: 119-129.
 135. Plé H, Maltais M, Corduan A, Rousseau G, Madore F, et al. Alteration of the platelet transcriptome in chronic kidney disease. *Thromb Haemost*. 2012; 108: 605-615.
 136. Healy AM, Pickard MD, Pradhan AD, Wang Y, Chen Z, et al. Platelet expression profiling and clinical validation of myeloid-related protein-14 as a novel determinant of cardiovascular events. *Circulation*. 2006; 113: 2278-2284.
 137. Colombo G, Gertow K, Marenzi G, Brambilla M, De Metrio M, et al. Gene expression profiling reveals multiple differences in platelets from patients with stable angina or non-ST elevation acute coronary syndrome. *Thromb Res*. 2011; 128: 161-168.
 138. Eicher JD, Wakabayashi Y, Vitseva O, Esa N, Yang Y, et al. Characterization of the platelet transcriptome by RNA sequencing in patients with acute myocardial infarction. *Platelets*. 2016; 27: 230-239.
 139. Reilly SJ, Li N, Liska J, Ekström M, Tornvall P. Coronary artery bypass graft surgery up-regulates genes involved in platelet aggregation. *J Thromb Haemost*. 2012; 10: 557-563.
 140. Saenz-Pipaon G, San Martín P, Planell N, Maillo A, Ravassa S, et al. Functional and transcriptomic analysis of extracellular vesicles identifies calprotectin as a new prognostic marker in peripheral arterial disease (PAD). *J Extracell Vesicles*. 2020; 9: 1729646.
 141. Wysokinski WE, Tafur A, Ammash N, Asirvatham SJ, Wu Y, et al. Impact of atrial fibrillation on platelet gene expression. *Eur J Haematol*. 2017; 98: 615-621.

142. Heitzer E, Haque IS, Roberts CES, Speicher MR. Current and future perspectives of liquid biopsies in genomics-driven oncology. *Nat Rev Genet.* 2019; 2: 71-88.
143. Babayan A, Pantel K. Advances in liquid biopsy approaches for early detection and monitoring of cancer. *Genome Med.* 2018; 10: 21.
144. Aravanis AM, Lee M, Klausner RD. Next-Generation Sequencing of Circulating Tumor DNA for Early Cancer Detection. *Cell.* 2017; 168: 571-574.
145. Hoshino A, Costa-Silva B, Shen TL, Rodrigues G, Hashimoto A, et al. Tumour exosome integrins determine organotropic metastasis. *Nature.* 2015; 527: 329-35.
146. Calverley DC, Phang TL, Choudhury QG, Gao B, Oton AB, et al. Significant downregulation of platelet gene expression in metastatic lung cancer. *Clin Transl Sci.* 2010; 3: 227-232.
147. Nilsson RJ, Balaj L, Hulleman E, van Rijn S, Pegtel DM, et al. Blood platelets contain tumor-derived RNA biomarkers. *Blood.* 2011; 118: 3680-3683.
148. Nilsson RJ, Karachaliou N, Berenguer J, Gimenez-Capitan A, Schellen P, et al. Rearranged EML4-ALK fusion transcripts sequester in circulating blood platelets and enable blood-based crizotinib response monitoring in non-small-cell lung cancer. *Oncotarget.* 2016; 7: 1066-1075.
149. Tjon-Kon-Fat LA, Lundholm M, Schröder M, Wurdinger T, Thellenberg-Karlsson C, et al. Platelets harbor prostate cancer biomarkers and the ability to predict therapeutic response to abiraterone in castration resistant patients. *Prostate.* 2018; 78: 48-53.
150. Klement GL, Yip TT, Cassiola F, Kikuchi L, Cervi D, et al. Platelets actively sequester angiogenesis regulators. *Blood.* 2009; 113: 2835-42
151. Best MG, Sol N, In 't Veld SGJG, Vancura A, Muller M, et al. Swarm Intelligence-Enhanced Detection of Non-Small-Cell Lung Cancer Using Tumor-Educated Platelets. *Cancer Cell.* 2017; 32: 238-252.e9.
152. Clancy L, Beaulieu LM, Tanriverdi K, Freedman JE. The role of RNA uptake in platelet heterogeneity. *Thromb Haemost.* 2017; 117: 948-961.

153. Navya PN, Kaphle A, Srinivas SP, Bhargava SK, Rotello VM, et al. Current trends and challenges in cancer management and therapy using designer nanomaterials. *Nano Converg.* 2019; 6: 23.
154. Wei Q, Qian Y, Yu J, Wong CC. Metabolic rewiring in the promotion of cancer metastasis: mechanisms and therapeutic implications. *Oncogene.* 2020; 39: 6139-6156.
155. Li T, Dong H, Zhang C, Mo R. Cell-Based Drug Delivery Systems for Biomedical Applications. *Nano Research.* 2018; 11: 5240-5257.
156. Zhang CL, Huang T, Wu BL, He WX, Liu D. Stem cells in cancer therapy: opportunities and challenges. *Oncotarget.* 2017; 8: 75756-75766.
157. Muzykantov VR. Drug delivery by red blood cells: vascular carriers designed by mother nature. *Expert Opin Drug Deliv.* 2010; 7: 403-27.
158. He W, Kapate N, Shields CW, Mitragotri S. Drug delivery to macrophages: A review of targeting drugs and drug carriers to macrophages for inflammatory diseases. *Adv Drug Deliv Rev.* 2019; 6: S0169-409X(19)30239-X.
159. Chu D, Dong X, Shi X, Zhang C, Wang Z. Neutrophil-Based Drug Delivery Systems. *Adv Mater.* 2018; 30: e1706245.
160. Schmid D, Park CG, Hartl CA, Subedi N, Cartwright AN, et al. T cell-targeting nanoparticles focus delivery of immunotherapy to improve antitumor immunity. *Nature Comm.* 2017; 8: 1747.
161. Xu XR, Yousef GM, Ni H. Cancer and platelet crosstalk: opportunities and challenges for aspirin and other antiplatelet agents. *Blood.* 2018; 131: 1777-1789.
162. Contursi A, Grande R, Dovizio M, Bruno A, Fullone R, et al. Platelets in cancer development and diagnosis. *Biochem Soc Trans.* 2018; 46: 1517-1527.
163. Morrell CN, Aggrey AA, Chapman LM, Modjeski KL. Emerging roles for platelets as immune and inflammatory cells. *Blood.* 2014; 123: 2759-2767.
164. Huong PT, Nguyen LT, Nguyen XB, Lee SK, Bach DH. The Role of Platelets in the Tumor-Microenvironment and the Drug Resistance of Cancer Cells. *Cancers (Basel).* 2019; 11: 240.

165. Qi C, Wei B, Zhou W, Yang Y, Li B, et al. P-selectin-mediated platelet adhesion promotes tumor growth. *Oncotarget*. 2015; 6: 6584-6596.
166. Haemmerle M, Bottsford-Miller J, Pradeep S, Taylor ML, Choi HJ, et al. FAK regulates platelet extravasation and tumor growth after antiangiogenic therapy withdrawal. *J. Clin. Invest*. 2016; 126: 1885–1896.
167. Ziegler M, Wang X, Lim B, Leitner E, Klingberg F, et al. Platelet-Targeted Delivery of Peripheral Blood Mononuclear Cells to the Ischemic Heart Restores Cardiac Function after Ischemia-Reperfusion Injury. *Theranostics*. 2017; 7: 3192-3206.
168. Banerjee M, Huang Y, Joshi S, Popa GJ, Mendenhall MD, et al. Platelets Endocytose Viral Particles and Are Activated via TLR (Toll-Like Receptor) Signaling. *Arterioscler Thromb Vasc Biol*. 2020; 40: 1635-1650.
169. Gaertner F, Ahmad Z, Rosenberger G, Fan S, Nicolai L, et al. Migrating Platelets Are Mechano- scavengers that Collect and Bundle Bacteria. *Cell*. 2017; 171: 1368-1382.e23.
170. White JG. Why human platelets fail to kill bacteria. *Platelets*. 2006; 17: 191–200.
171. White JG. Platelets are covercytes, not phagocytes: uptake of bacteria involves channels of the open canalicular system. *Platelets*. 2005; 16: 121-131.
172. Sarkar S, Alam MA, Shaw J, Dasgupta AK. Drug delivery using platelet cancer cell interaction. *Pharm Res*. 2013; 30: 2785-2794.
173. Xu P, Zuo H, Chen B, Wang R, Ahmed A, et al. Doxorubicin-loaded platelets as a smart drug delivery system: An improved therapy for lymphoma. *SciRep*. 2017; 7: 42632.
174. Han X, Wang C, Liu Z. Red Blood Cells as Smart Delivery Systems. *Bioconjug Chem*. 2018; 29: 852-860.
175. Feins S, Kong W, Williams EF, Milone MC, Fraietta JA. An introduction to chimeric antigen receptor (CAR) T-cell immunotherapy for human cancer. *Am J Hematol*. 2019; 94: S3-S9.

176. Lyde R, Sabatino D, Sullivan SK, Poncz M. Platelet-delivered therapeutics. *J Thromb Haemost.* 2015; 13: S143-150.
177. Fang J, Hodivala-Dilke K, Johnson BD, DU LM, Hynes RO, et al. Therapeutic expression of the platelet-specific integrin, α IIb β 3, in a murine model for Glanzmann thrombasthenia. *Blood.* 2005; 106: 2671–2679.
178. Shi Q, Wilcox DA, Morateck PA, Fahs SA, Kenny D, et al. Targeting platelet GPIIb/IIIa transgene expression to human megakaryocytes and forming a complete complex with endogenous GPIIb/IIIa and GPIIb/IIIa. *J Thromb Haemost.* 2004; 2: 1989–1997.
179. Shi Q, Kuether EL, Cooley BC, Fahs SA, Schroeder JA, et al. Sustained Phenotypic Correction of Murine Hemophilia A with Pre-Existing Anti-FVIII Immunity Using Lentivirus-Mediated Platelet-Specific FVIII Gene Transfer. *ASH Annual Meeting Abstracts.* 2009; 114: 29.
180. Levine RF. Isolation and characterization of normal human megakaryocytes. *Br J Haematol.* 1980; 45: 487–497.
181. Sim X, Jarocho D, Hayes V, Hanby HA, Marks MS, et al. Identifying and enriching platelet-producing human stem cell-derived megakaryocytes using factor V uptake. *Blood.* 2017; 130: 192-204.
182. Sharma P, Allison JP. Dissecting the mechanisms of immune checkpoint therapy. *Nat Rev Immunol.* 2020; 20: 75-76.
183. Wei SC, Levine JH, Cogdill AP, Zhao Y, Anang NAS, et al. Distinct Cellular Mechanisms Underlie Anti-CTLA-4 and Anti-PD-1 Checkpoint Blockade. *Cell.* 2017; 170: 1120-1133.e17.
184. Motzer RJ, Escudier B, McDermott DF, George S, Hammers HJ, et al. Nivolumab versus Everolimus in Advanced Renal-Cell Carcinoma. *N Engl J Med.* 2015; 373: 1803-1813.
185. Chitsike L, Duerksen-Hughes P. The Potential of Immune Checkpoint Blockade in Cervical Cancer: Can Combinatorial Regimens Maximize Response? A Review of the Literature. *Curr Treat Options Oncol.* 2020; 21: 95.

186. Kwapisz D. Pembrolizumab and atezolizumab in triple-negative breast cancer. *Cancer Immunol Immunother.* 2021; 70: 607-617.
187. Hallqvist A, Rohlin A, Raghavan S. Immune checkpoint blockade and biomarkers of clinical response in Non-small cell lung cancer (NSCLC). *Scand J Immunol.* 2020; e12980.
188. de Melo Gagliato D, Buzaid AC, Perez-Garcia J, Cortes J. Immunotherapy in Breast Cancer: Current Practice and Clinical Challenges. *BioDrugs.* 2020; 34: 611-623.
189. Zhang X, Wang J, Chen Z, Hu Q, Wang C, et al. Engineering PD-1-Presenting Platelets for Cancer Immunotherapy. *Nano Lett.* 2018; 18: 5716-5725.
190. Scurr M, Pembroke T, Bloom A, Roberts D, Thomson A, et al. Low-Dose Cyclophosphamide Induces Antitumor T-Cell Responses, which Associate with Survival in Metastatic Colorectal Cancer. *Clin Cancer Res.* 2017; 23: 6771-6780.
191. Deng D, Shah K. TRAIL of Hope Meeting Resistance in Cancer. *Trends Cancer.* 2020; S2405-8033(20)30190-4.
192. Li J, Sharkey CC, Wun B, Liesveld JL, King MR. Genetic engineering of platelets to neutralize circulating tumor cells. *J Control Release.* 2016; 228: 38-47.
193. Chan V, Novakowski SK, Law S, Klein-Bosgoed C, Kastrop CJ. Controlled Transcription of Exogenous mRNA in Platelets Using Protocells. *Angew Chem Int Ed Engl.* 2015; 54: 13590-13593.
194. Wang C, Sun W, Ye Y. In situ activation of platelets with checkpoint inhibitors for post-surgical cancer immunotherapy. *Nat Biomed Eng.* 2017; 1: 0011.
195. Hu Q, Sun W, Wang J, Ruan H, Zhang X, et al. Conjugation of haematopoietic stem cells and platelets decorated with anti-PD-1 antibodies augments anti-leukaemia efficacy. *Nat Biomed Eng.* 2018; 2: 831-840.
196. Kolb HC, Sharpless KB. The growing impact of click chemistry on drug discovery. *Drug Discov. Today.* 2003; 8: 1128-1137.
197. Shi J, Kantoff PW, Wooster R, Farokhzad OC. Cancer nanomedicine: progress, challenges and opportunities. *Nat Rev Cancer.* 2017; 17: 20-37.
198. Bose RJ, Paulmurugan R, Moon J, Lee SH, Park H. Cell membrane-coated nanocarriers: the emerging targeted

- delivery system for cancer theranostics. *Drug Discov Today*. 2018; 23:891-899.
199. Wu M, Le W, Mei T, Wang Y, Chen B, et al. Cell membrane camouflaged nanoparticles: a new biomimetic platform for cancer photothermal therapy. *Int J Nanomedicine*. 2019; 14: 4431-4448.
200. Hu Q, Sun W, Qian C, Wang C, Bomba HN, et al. Anticancer Platelet-Mimicking Nanovehicles. *Adv Mater*. 2015; 27: 7043-7050.
201. Jing L, Qu H, Wu D, Zhu C, Yang Y, et al. Platelet-camouflaged nanococktail: Simultaneous inhibition of drug-resistant tumor growth and metastasis via a cancer cells and tumor vasculature dual- targeting strategy. *Theranostics*. 2018; 8: 2683-2695.
202. Chen Y, Zhao G, Wang S, He Y, Han S, et al. Platelet-membrane-camouflaged bismuth sulfide nanorods for synergistic radio- photothermal therapy against cancer. *Biomater Sci*. 2019; 7: 3450-3459.
203. Jiang Q, Wang K, Zhang X, Ouyang B, Liu H, et al. Platelet Membrane- Camouflaged Magnetic Nanoparticles for Ferroptosis-Enhanced Cancer Immunotherapy. *Small*. 2020;16: e2001704.
204. Choi J, Fenando A. Sulfasalazine. 2020 Jun 28. In: *StatPearls* [Internet]. Treasure Island: StatPearls Publishing. 2020.
205. Xie Y, Hou W, Song X, Yu Y, Huang J, et al. Ferroptosis: process and Guillemin function. *Cell Death Differ*. 2016; 23: 369-379.
206. Gasic GJ, Gasic TB, Stewart CC, Antimetastatic effects associated with platelet reduction, *Proc Natl Acad Sci U S A*. 1968; 61: 46-52.
207. Palmblad J. To give and take - life of a platelet. *Blood*. 2009; 113: 2617.
208. Yang J, Wang S, Liu P, Dai L, Chen B, et al. Platelet-inspired medicine for tumor therapy. *Oncotarget*. 2017; 8: 115748-115753.
209. Felding-Habermann B, Habermann R, Saldívar E, Ruggeri ZM. Role of beta3 integrins in melanoma cell adhesion to activated platelets under flow. *J Biol Chem*. 1996; 271: 5892–5900.

210. Mitrugno A, Williams D, Kerrigan SW, Moran N. A novel and essential role for FcγRIIIa in cancer cell-induced platelet activation. *Blood*. 2014; 123: 249–260.
211. Boukerche H, Berthier-Vergnes O, Tabone E, Bailly M, Doré JF, et al. Thrombospondin modulates melanoma--platelet interactions and melanoma tumour cell growth in vivo. *Br J Cancer*. 1995; 72: 108-116.
212. Mammadova-Bach E, Zigrino P, Brucker C, Bourdon C, Freund M, et al. Platelet integrin controls lung metastasis through direct binding to cancer cell-derived ADAM9. *JCI Insight*. 2016; 1: e88245.
213. Mannori G, Crottet P, Cecconi O, Hanasaki K, Aruffo A, et al. Differential colon cancer cell adhesion to E-, P-, and L-selectin: role of mucintype glycoproteins. *Cancer Research*. 1995; 55: 4425–4431.
214. Man Y, Goreke U, Kucukal E, Hill A, An R, et al. Leukocyte adhesion to P-selectin and the inhibitory role of Crizanlizumab in sickle cell disease: A standardized microfluidic assessment. *Blood Cells Mol Dis*. 2020; 83102424.
215. Gong L, Cai Y, Zhou X, Yang H. Activated platelets interact with lung cancer cells through P-selectin glycoprotein ligand-1. *Pathol Oncol Res*. 2012; 18: 989–996.
216. Alves CS, Burdick MM, Thomas SN, Pawar P, Konstantopoulos K. The dual role of CD44 as a functional P-selectin ligand and fibrin receptor in colon carcinoma cell adhesion. *Am J Physiol Cell Physiol*. 2008; 294: C907–C916.
217. Larrucea S, Butta N, Rodriguez RB, Alonso-Martin S, Arias-Salgado EG, et al. Podocalyxin enhances the adherence of cells to platelets. *Cell Mol Life Sci*. 2007; 64: 2965–2974.
218. Suzuki-Inoue K, Kato Y, Inoue O, Kaneko MK, Mishima K, et al. Involvement of the snake toxin receptor CLEC-2, in podoplanin-mediated platelet activation, by cancer cells. *J Biol Chem*. 2007; 282: 25993–26001.
219. Chang YW, Hsieh PW, Chang YT, Lu MH, Huang TF, et al. Identification of a novel platelet antagonist that binds to CLEC-2 and suppresses podoplanin-induced platelet

- aggregation and cancer metastasis. *Oncotarget*. 2015; 6: 42733–42748.
220. Kato Y, Kaneko MK, Kuno A, Uchiyama N, Amano K, et al. Inhibition of tumor cell-induced platelet aggregation using a novel anti-podoplanin antibody reacting with its platelet-aggregation-stimulating domain. *Biochem Biophys Res Commun*. 2006; 349: 1301–1307.